

Physics 745 - Group Theory  
**Final, Spring 2009**

You may use (1) class notes, (2) the text, (3) former homeworks and solutions (available online), or (4) any math references, such as integral tables, Maple, etc, including any routines I provided for you. If you cannot solve an equation, try to go on, as if you knew the answer. Feel free to contact me with questions. Note that the final problem is more like a homework problem, you should feel free to ask me for help on it. Each question is worth 20 points out of a total of 100 points.

Work: 758-4994      Home: 724-2008      Cell: 407-6528

1. The proper icosohedral group is a 60 element group. Its character table is given at right. The irreps have been unimaginatly labeled as  $A, B, C, D,$  and  $E$ . For each of the five tensor products with  $E$ , break the tensor product into appropriate irreps; *i.e.*, work out  $A \otimes E, B \otimes E, \dots, E \otimes E$ .

$\mathcal{I}$	$E$	$20C_3$	$12C_5$	$12C_5^2$	$15C_2$
$A$	1	1	1	1	1
$B$	3	0	$\frac{1+\sqrt{5}}{2}$	$\frac{1-\sqrt{5}}{2}$	-1
$C$	3	0	$\frac{1-\sqrt{5}}{2}$	$\frac{1+\sqrt{5}}{2}$	-1
$D$	4	1	-1	-1	0
$E$	5	-1	0	0	1



2. The group  $SO(5)$  has ten generators, which can be labeled  $T_{ab}$ , where  $a, b \in \{1, 2, 3, 4, 5\}$ , and  $a \neq b$  (they are defined in such a way that  $T_{ab} = -T_{ba}$ , which is why there are only ten of them). They satisfy the commutation relations

$$[T_{ab}, T_{cd}] = i(\delta_{ac}T_{bd} + \delta_{bd}T_{ac} - \delta_{ad}T_{bc} - \delta_{bc}T_{ad})$$

where  $\delta_{ab}$  is the Kronecker delta function.

- (a) Let  $J_1 = T_{13}, J_2 = T_{23}, J_3 = T_{12}$ . Show that  $\mathbf{J}$  generates an  $SU(2)$  subgroup of  $SO(5)$ , *i.e.*, show all three of the commutators  $[J_a, J_b] = i\epsilon_{abc}J_c$ .

- (b) At right are given the weight diagram of two irreps of  $SO(5)$  in the basis of  $T_{12}$  and  $T_{34}$ , *i.e.*, the eigenvalues of these two generators are plotted.. The tick marks are at one unit. For each of these representations (which I'll call the "4" and the "5"), work out what irreps of  $SU(2)$  these break into.



3. An atom is in the state  $|njm\rangle$  with  $j = 1$  is about to decay via electric dipole radiation to the state  $|n'j'm'\rangle$ . As argued in homework set 25, the probability of it going from an initial state  $I$  to a final state  $F$  is proportional to

$$\Gamma(I \rightarrow F) = \sum_{q=-1}^1 |\langle F | r_q^{(1)} | I \rangle|^2$$

where  $r_q^{(1)}$  is the spherical tensor operator corresponding to the vector operator  $\mathbf{r}$ .

- (a) What are the possible final values of  $j'$ ?  
 (b) In fact, it is going to decay to a state with  $j' = 1$ . Using the Wigner Eckart Theorem, find the *relative* rate of decay

$$\Gamma(njm \rightarrow n'j'm')$$

for all non-vanishing possible values of  $m$  and  $m'$ .

4. Although the 1, 8, and 10 irreps of Gell-Mann SU(3) are all that are used when we look at combinations of up, down, and strange quarks, other possibilities occur with heavier quarks. For example, the six lightest spin 3/2 baryons containing a charm quark are listed with their mass in the table at right. These particles fit into the 6 irrep of SU(3), which can be written in the form (note: the  $c$  is not an index, it represents the presence of a charm quark)

$$|B_c^*\rangle = u^{ij} |B_{c,ij}^*\rangle$$

with the assignments:

$$\Sigma_c^{*0} : u^{22} = 1, \quad \Sigma_c^{*+} : u^{21} = u^{12} = \frac{1}{\sqrt{2}}, \quad \Sigma_c^{*++} : u^{11} = 1,$$

$$\Xi_c^{*0} : u^{32} = u^{23} = \frac{1}{\sqrt{2}}, \quad \Xi_c^{*+} : u^{31} = u^{13} = \frac{1}{\sqrt{2}}, \quad \Omega_c^{*0} : u^{33} = 1.$$

Name	Mass	$I$	$I_3$
$\Sigma_c^{*++}$	2518	1	+1
$\Sigma_c^{*+}$	2517	1	0
$\Sigma_c^{*0}$	2518	1	-1
$\Xi_c^{*+}$	2647	1/2	+1/2
$\Xi_c^{*0}$	2646	1/2	-1/2
$\Omega_c^{*0}$	????	0	0


All masses in MeV/ $c^2$

- (a) Work out a formula for the mass of these objects in terms of the  $u$ 's. Include not only an SU(3) respecting piece, but also include a piece where the symmetry is broken proportional to  $T_8$ . Include unknown parameters as needed.  
 (b) Write an expression for the mass of one of the  $\Sigma_c^*$ 's, one of the  $\Xi_c^*$ 's, and the  $\Omega_c^{*0}$  in terms of the parameters you chose in part (a). Find a linear relationship between them. Predict, on the basis of your relationship, the mass of the  $\Omega_c^{*0}$ .

*Note: This last problem is more like a homework problem; if you are having difficulty with it, come see me, and we will get you unstuck.*

5. The last day of class, I presented very abbreviated proofs that certain Dynkin diagrams are not allowed. You are going to elaborate one of them. The diagram at right is an illegal Dynkin diagram, as you will demonstrate.

- (a) Define one of the roots to have length  $r$ . Write the length of each of the five simple roots in terms of  $r$ .



- (b) For every pair of simple roots for which the dot product doesn't vanish, write the dot product between them in terms of  $r$ .
- (c) Show that an appropriate combination of the roots above vanishes; *i.e.*, that the square of the combination is zero, and hence this diagram is illegal.