

Test 1 Name _____
September 18, 2019

This test consists of three parts. Please note that in parts II and III, you can skip one question of those offered.

Possibly useful formulas:

$x' = \gamma(x - vt)$ $t' = \gamma(t - vx/c^2)$ $y' = y, \quad z' = z$	$E' = \gamma(E - vp_x)$ $p'_x = \gamma(p_x - vE/c^2)$ $p'_y = p_y, \quad p'_z = p_z$	$f = \frac{f_0}{\gamma(1 - v \cos \theta/c)}$ $e = 1.602 \times 10^{-19} \text{ C}$	$u'_x = \frac{u_x - v}{1 - vu_x/c^2}$ $u'_y = \frac{u_y}{\gamma(1 - vu_x/c^2)}$ $u'_z = \frac{u_z}{\gamma(1 - vu_x/c^2)}$
$\vec{F} = q\vec{E} + q\vec{u} \times \vec{B}$ $p = qRB$	$(1 + \varepsilon)^n = 1 + n\varepsilon + \frac{1}{2}n(n-1)\varepsilon^2 + \dots$	$\frac{\vec{u}}{c} = \frac{\vec{p}c}{E}$	

Part I: Multiple Choice [20 points]

For each question, choose the best answer (2 points each)

1. Two observers are moving relative to each other, and each tries to measure the other's meter sticks, which are oriented in the direction of their relative motion. What will they conclude about each other's meter sticks?
 - A) They will both agree that the one who is truly at rest will have the shorter meter stick
 - B) They will both agree that the one who is truly moving will have the shorter meter stick
 - C) Each one will see the other's stick as longer than their own
 - D) Each one will see the other's stick as shorter than their own
 - E) Insufficient information

2. When is the formula $E = mc^2$ valid?
 - A) Always
 - B) Only when the object is at rest
 - C) Only when the object is moving
 - D) Only on Friday the 13th
 - E) Never

3. The total energy of an object of mass m and velocity v is

A) $\frac{mc^2}{\sqrt{1-v^2/c^2}}$ B) $mc^2\sqrt{1-v^2/c^2}$ C) mc^2 D) $\frac{1}{2}mv^2$ E) $\frac{mvc}{\sqrt{1-v^2/c^2}}$

4. If you pushed on an object initially at rest with mass m with a force F , how long would you have to push to get the object to the speed of light c ?

A) $t = \frac{F}{mc}$ B) $t = \frac{Fm}{c}$ C) $t = \frac{Fc}{m}$ D) $t = \frac{mc}{F}$ E) Forever; you can never reach c


5. If events P_1 and P_2 have a timelike separation, what could be the relationship of P_2 to P_1 ?

- A) P_2 is in the absolute future of P_1 (only)
 B) P_2 is in the absolute past of P_1 (only)
 C) P_2 is elsewhere compared to P_1 (only)
 D) Both A and B are possible, but not C
 E) A, B, and C are all possibilities
6. It is possible to collide a proton and anti-proton (each mass m_p) to make three neutral pions (each mass m_π). From this information alone, what can we conclude about the pion mass?
 A) $m_\pi = 2m_p$ B) $m_\pi = \frac{2}{3}m_p$ C) $m_\pi = \frac{3}{2}m_p$ D) $m_\pi = 3m_p$ E) Nothing
7. If you tried to communicate using a perfectly rigid rod by pushing on one end while measuring the position of the other, you would find
 A) The other end moves simultaneously as viewed in all reference frames
 B) The other end moves simultaneously only in the frame where the rod is not moving
 C) The other end moves simultaneously only in the frame where the signaler is not moving
 D) The other end moves simultaneously only in the frame where the receiver is not moving
 E) Rigid rods are impossible in special relativity, so the question makes no sense
8. If you have two events (points in spacetime), which of the following will all observers agree on?
 A) Their separation in space (only)
 B) The difference in time between them (only)
 C) The proper distance squared, s^2 (only)
 D) The separation in space and in time, but not the proper distance squared
 E) The separation in space and in time as well as the proper distance
9. An electron has a mass squared that is _____ and a photon has a mass squared that is _____.
 A) Positive, positive
 B) Positive, zero
 C) Positive, negative
 D) Zero, positive
 E) Zero, zero
10. A particle of mass m initially pushed by a force F will experience a constant acceleration $a = F/m$, causing it to reach a speed $v = at$ and travel a distance $d = \frac{1}{2}at^2$, and reaching the speed of light after a time $t = c/a$ in a distance $d = c^2/2a$. Which of these formulas is the first one where I made a mistake, according to relativity?
 A) $a = F/m$ B) $v = at$ C) $d = \frac{1}{2}at^2$ D) $t = c/a$ E) $d = c^2/2a$

Part II: Short answer [20 points]

Choose **two** of the following questions and give a short answer (1-3 sentences) (10 points each).

11. A cylindrical spaceship is 4.00 m in length and 4.00 m in diameter. It is traveling so fast that it has a Lorentz factor of $\gamma = 2.00$. Describe its apparent shape, as observed by us, if it is moving (a) to the right \rightarrow at this speed or (b) moving up \uparrow at this speed.



12. The Tevatron was an approximately circular particle collider 1.00 km in radius that could accelerate protons up to a momentum of $0.980 \text{ TeV}/c$. The LHC uses slightly better magnets, is 4.30 km in radius, and can get the protons up to a momentum of $7.00 \text{ TeV}/c$. Explain what magnets have to do with it, and explain what advantage the LHC has that allows it to reach higher momenta. You should include at least one equation in your answer.
13. In a homework problem, you found that a “bunch” of protons, when it passes through the Atlas detector, as viewed by us is much shorter than the Atlas detector, but if you were moving along with them it is much longer than the Atlas detector. But surely the protons are either all in the detector at the same time or not. Resolve this apparent paradox.

Part III: Calculation: [60 points]

Choose **three** of the following four questions and perform the indicated calculations (20 points each)


14. A collection of 4.17×10^6 muons is traveling around a circle of radius $r = 26.9$ m, completing a circuit every $0.710 \mu\text{s}$.

- What is the speed of these particles?
- From the viewpoint of the muons, how long does it take them to go around once?
- The probability of a muon not decaying after a time τ as viewed in its own frame is given by $P = e^{-\tau/\tau_0}$, where $\tau_0 = 2.197 \mu\text{s}$ is the mean lifetime. How many muons are left after ten complete circuits?

15. A beam of light start at the origin, $(x, y, z, t) = (0, 0, 0, 0)$, traveling in the $+y$ direction. It is detected by two detectors each 2.00 m away from the initial event.

- What are all four coordinates (x, y, z, t) when the light beam is detected?
- An observer traveling at $v = 1.80 \times 10^8$ m/s in the $+x$ direction observes the emission and detection. Find the coordinates of the detection event (and emission event) (x', y', z', t') as viewed by this observer.
- Calculate the components of the velocity of light $\vec{v} = (v_x, v_y, v_z)$ using the locations and time you found in part (b).
- Calculate the speed (magnitude of the velocity) using the numbers you found in part (c). Comment on whether the answer makes sense or not.

16. The Δ^{++} baryons is produced by colliding positively charges pions π^+ ($m_\pi = 139 \text{ MeV}/c^2$) with an energy of $E_\pi = 328 \text{ MeV}$ with protons at rest ($m_p = 938 \text{ MeV}/c^2$).



- What is the momentum (in MeV/c) and speed (as a fraction of c) of the pions?
- What is the energy and momentum of the resulting Δ^{++} ?
- What is the mass (in MeV/c^2) and speed (as a fraction of c) of the resulting Δ^{++} ?

17. Neutral hydrogen atoms at rest emit radio waves at a frequency of 1420.4 MHz . A cloud of hydrogen atoms is moving directly away from us at $v = 172,000 \text{ km/s}$.

- At what frequency will we observe the radio waves from this source?
- If the cloud of gas has a mass $m = 1.989 \times 10^{30} \text{ kg}$, what is its momentum?
- If a constant force of $F = 1.56 \times 10^{26} \text{ N}$ were applied to slow down this cloud, how long would it take in years to bring it to rest? ($1 \text{ y} = 3.156 \times 10^7 \text{ s}$)