

Physics 741 – Graduate Quantum Mechanics 1
Solutions to Chapter 14

6. [25] A particle of mass m scatters from a potential $V(r) = F\delta(r-a)$, so that the potential exists only at the surface of a thin sphere of radius a .
(a) [4] What equation must the radial wave functions $R_l(r)$ satisfy? Solve this equation in the regions $r < a$ and $r > a$.

The radial wave functions must satisfy

$$\frac{1}{r} \frac{d^2}{dr^2} [rR_l(r)] = \left[\frac{l^2 + l}{r^2} + U(r) - k^2 \right] R_l(r) = \left[\frac{l^2 + l}{r^2} + \frac{2mF}{\hbar^2} \delta(r-a) - k^2 \right] R_l(r).$$

Away from the point $r = a$, the problem is simply that of a free particle, and the solution was worked out in class. The answer is spherical Bessel functions, and take the form

$$R_l(r) = \begin{cases} \gamma j_l(kr) - \delta n_l(kr) & r < a, \\ \alpha j_l(kr) - \beta n_l(kr) & r > a. \end{cases}$$

The constants will generally be different in the different regions.

- (b) [6] Apply appropriate boundary at $r = 0$. Deduce appropriate boundary conditions at $r = a$.

We want the radial function to be well-behaved at $r = 0$, which implies we only want the well-behaved $j_l(r)$. Hence we demand $\delta = 0$. At the boundary $r = a$, we must take our radial Schrödinger equation and integrate it across the boundary. We first multiply both sides by r , then we have

$$\int_{a-\varepsilon}^{a+\varepsilon} \frac{d^2}{dr^2} [rR_l(r)] dr = \int_{a-\varepsilon}^{a+\varepsilon} \left[\frac{l^2 + l}{r} + \frac{2mF}{\hbar^2} r\delta(r-a) - k^2 r \right] R_l(r) dr,$$

$$[rR_l(r)]' \Big|_{a-\varepsilon}^{a+\varepsilon} = \frac{2mF}{\hbar^2} aR_l(a).$$

Since the first derivative has a finite discontinuity, it follows that the wave function will be continuous at the boundary. This yields two boundary conditions on the wave function:

$$\alpha j_l(ka) - \beta n_l(ka) = \gamma j_l(ka),$$

$$\alpha [rj_l(kr)]'_{r=a} - \beta [rn_l(kr)]'_{r=a} = \frac{2mF}{\hbar^2} a\gamma j_l(ka) + \gamma [rj_l(kr)]'_{r=a}$$

Our goal, ultimately, will be to eliminate γ from these equations and deduce the ratio of β to α .

(c) [8] Assume now that $ka \ll 1$, so that the scattering will be dominated by the $l=0$ term. Find a formula for the phase shift δ_0 . Find the differential cross-section. Check that your formula agrees with the formula found in section C for the hard sphere in the case $F \rightarrow \infty$.

We substitute in the explicit form for $l=0$, namely $j_0(x) = \sin x/x$ and $n_0(x) = -\cos x/x$. Then our two boundary conditions become

$$\alpha \sin(ka)/(ka) + \beta \cos(ka)/(ka) = \gamma \sin(ka)/(ka),$$

$$\alpha \left[\sin(kr)/k \right]'_{r=a} + \beta \left[\cos(kr)/k \right]'_{r=a} = \frac{2mF}{\hbar^2} a \gamma \sin(ka)/(ka) + \gamma \left[\sin(kr)/k \right]'_{r=a}.$$

Clear the fractions from the first of these and work out the derivatives in the second.

$$\alpha \sin(ka) + \beta \cos(ka) = \gamma \sin(ka),$$

$$\alpha \cos(ka) - \beta \sin(ka) = \frac{2mF}{\hbar^2 k} \gamma \sin(ka) + \gamma \cos(ka).$$

Cross multiply these and then cancel the common factor of γ .

$$\left[\alpha \cos(ka) - \beta \sin(ka) \right] \sin(ka) = \left[2mF \sin(ka)/\hbar^2 k + \cos(ka) \right] \left[\alpha \sin(ka) + \beta \cos(ka) \right],$$

$$\beta \left[-\hbar^2 k \sin^2(ka) - \hbar^2 k \cos^2(ka) - 2mF \sin(ka) \cos(ka) \right] = \alpha 2mF \sin^2(ka),$$

$$\frac{\beta}{\alpha} = \frac{-2mF \sin^2(ka)}{\hbar^2 k + 2mF \sin(ka) \cos(ka)}.$$

We hence have the phase shift

$$\tan \delta_0 = \frac{\beta}{\alpha} = \frac{-2mF \sin^2(ka)}{\hbar^2 k + 2mF \sin(ka) \cos(ka)} \approx \frac{-2mF k^2 a^2}{\hbar^2 k + 2mF k a} = \frac{-2mF a^2 k}{\hbar^2 + 2mF a}.$$

where we used $ka \ll 1$ to approximate $\sin ka = ka$ and $\cos ka = 1$. It's then easy to see that the numerator is much smaller than the denominator, so we can also approximate $\sin \delta_0 = \tan \delta_0$, and hence the differential cross-section is

$$\frac{d\sigma}{d\Omega} = \frac{4\pi}{k^2} \sin^2 \delta_0 |Y_0^0|^2 = \frac{1}{k^2} \left(\frac{-2mF k a^2}{\hbar^2 + 2mF a} \right)^2 = \frac{4m^2 F^2 a^4}{(\hbar^2 + 2mF a)^2}.$$

In the limit of infinite potential, we ignore the first term in the denominator compared to the second, so $d\sigma/d\Omega = a^2$, the same as we found before.

(d) [7] Redo the problem using the first Born approximation. Again assume $ka \ll 1$ (effectively, this means $Ka = 0$). Check that the resulting differential cross-section in this case is identical with that found above in the limit $F \rightarrow 0$.

In the first Born approximation, the differential cross-section is given by

$$\begin{aligned} \frac{d\sigma}{d\Omega} &= \frac{m^2}{4\pi^2\hbar^4} \left| \int d^3\mathbf{r} V(\mathbf{r}) e^{-i\mathbf{K}\cdot\mathbf{r}} \right|^2 \approx \frac{m^2}{4\pi^2\hbar^4} \left| \int d^3\mathbf{r} V(\mathbf{r}) \right|^2 = \frac{m^2}{4\pi^2\hbar^4} \left[4\pi \int_0^\infty r^2 dr F \delta(r-a) \right]^2 \\ &= \frac{m^2}{4\pi^2\hbar^4} (4\pi a^2 F)^2 = \frac{4m^2 a^4 F^2}{\hbar^4}. \end{aligned}$$

In the limit of small F , the previous computation yields $d\sigma/d\Omega = 4m^2 F^2 a^4 / \hbar^4$, so they match.