

MTH 352/652

Homework #2

Due Date: January 31, 2025

1. Find all real valued solutions to the two-dimensional Laplace equation $u_{xx} + u_{yy} = 0$ that depend on the radial coordinate $r = \sqrt{x^2 + y^2}$.
2. Suppose $u(x, t)$ and $v(x, t)$ are infinitely differentiable functions in both x and t that satisfy the following system of equations:

$$u_t = v_x,$$

$$v_t = u_x.$$

- (a) Show that both u and v are solutions to the wave equation $u_{tt} = u_{xx}$, $v_{tt} = v_{xx}$. Which result from multivariable calculus do you need to justify the conclusion?
 - (b) Conversely, given a solution $u(x, t)$ to the wave equation, can you construct a function $v(x, t)$ such that $u(x, t), v(x, t)$ solve $u_t = v_x$ and $v_t = u_x$.
3. Classify the following differential equations as either (i) homogenous linear, (ii) inhomogeneous linear, or (iii) nonlinear.

(a) $u_t = x^2 u_{xx} + 2x u_x$

(b) $-u_{xx} - u_{yy} = \cos(u)$

(c) $u_{xx} + 2y u_{yy} = 3$

(d) $u_t + u u_x = 3u$

(e) $e^y u_x = e^x u_y$

(f) $u_t = 5u_{xxx} + x^2 u + x$

4. Suppose L and M are linear operators. Prove that the following are also linear operators:

(a) $L - M$.

(b) $3L$.

(c) fL , where f is an arbitrary function.

(d) $L \circ M$.

5. The displacement $u(x, t)$ of a forced violin string is modeled by the PDE $u_{tt} = 4u_{xx} + F(x, t)$. When $F(x, t) = \cos(x)$, the solution is $u(x, t) = \cos(x - 2t) + \frac{1}{4} \cos(x)$, while when $F(x, t) = \sin(x)$, the solution is $u(x, t) = \sin(x - 2t) + \frac{1}{4} \sin(x)$. Find a solution when the forcing function $F(x, t)$ is

(a) $\cos(x) - 5 \sin(x)$,

(b) $\sin(x - 3)$.

6. Find the general solution to the following partial differential equations
- $u_x = 0$
 - $u_t = 1$
 - $u_t = x - t$
 - $u_t + 3u = 0$
 - $u_x + tu = 0$
 - $u_{tt} + 4u = 0$
7. Solve the following initial value problems and graph the solutions at times $t = 0, 1, 2, 3$.
- $u_t - 3u_x = 0, u(x, 0) = e^{-x^2}$
 - $u_t + 2u_x = 0, u(x, -1) = x/(1 + x^2)$
 - $u_t + u_x + \frac{1}{2}u = 0, u(x, 0) = \tan^{-1}(x)$
 - $u_t - 4u_x + u = 0, u(x, 0) = 1/(1 + x^2)$
8. Let $c \neq 0$. Prove that if the initial data satisfies $u(x, 0) = v(x) \rightarrow 0$ as $x \rightarrow \pm\infty$, then, for each fixed x , the solution to the advection equation $u_t + cu_x = 0$ satisfies $u(x, t) \rightarrow 0$ as $t \rightarrow \infty$.
9. Solve the following initial value problem $u_t + 2u_x = \sin(x), u(x, 0) = \sin(x)$.
10. Consider the partial differential equation $u_t + u_x + u^2 = 0, u(x, 0) = f(x)$ for $x \in \mathbb{R}$ and $t \geq 0$.
- Find the general formula for the solution to this PDE.
 - Show that if $f(x)$ is positive and bounded, i.e., $0 \leq f(x) \leq M$, then the solution exists for all $t > 0$, and $u(x, t) \rightarrow 0$ as $t \rightarrow \infty$.
 - On the other hand, if $f(x)$ is negative somewhere, then show that the solution blows up in finite time: $\lim_{t \rightarrow \tau^-} u(y, t) = -\infty$ for some $\tau > 0$ and some $y \in \mathbb{R}$.
 - Find a formula for the earliest blow-up time $\tau^* > 0$.
11. Consider the initial value problem $u_t - xu_x = 0, u(x, 0) = (x^2 + 1)^{-1}$.
- Write down the differential equation satisfied by the characteristic curves and sketch the curves. Note, you don't have to find an explicit formula for the characteristic curves.
 - Sketch the solution $u(x, t)$ at times $t = 0, 1, 5, 10$.
 - Compute $\lim_{t \rightarrow \infty} u(x, t)$.
12. Consider the initial value problem $u_t + (1 + x^2)u_x = 0, u(0, x) = f(x)$.
- Write down the differential equation satisfied by the characteristic curves and find an explicit solution for the characteristic curves.
 - Sketch the characteristic curves.
 - Write down the general solution $u(x, t)$.
 - Discuss properties of your solution as t increases.