

# MTH 352/652

## Homework #3

Due Date: February 07, 2025

1. Consider the initial value problem  $u_t - xu_x = 0$ ,  $u(x, 0) = (x^2 + 1)^{-1}$ .
  - (a) Write down the differential equation satisfied by the characteristic curves and sketch the curves. Note, you don't have to find an explicit formula for the characteristic curves.
  - (b) Sketch the solution  $u(x, t)$  at times  $t = 0, 1, 5, 10$ .
  - (c) Compute  $\lim_{t \rightarrow \infty} u(x, t)$ .
2. Consider the initial value problem  $u_t + (1 + x^2)u_x = 0$ ,  $u(0, x) = f(x)$ .
  - (a) Write down the differential equation satisfied by the characteristic curves and find an explicit solution for the characteristic curves.
  - (b) Sketch the characteristic curves.
  - (c) Write down the general solution  $u(x, t)$ .
  - (d) Discuss properties of your solution as  $t$  increases.
3. Consider the advection equation with a speed that varies in time and space:

$$\begin{aligned}u_t + c(x, t)u_x &= 0 \\ u(x, 0) &= u_0(x),\end{aligned}$$

where  $x \in \mathbb{R}$  and  $t \in \mathbb{R}^+$ . The characteristic curves are solutions to the following differential equation:

$$\frac{dx}{dt} = c(x, t).$$


- (a) Show that any solution to the above advection equation is constant along the characteristic curves.
- (b) Suppose that the general form of the solutions to the characteristic equation can be written in the form  $g(x(t), t) = k$ , where  $k$  is an arbitrary constant. Show that  $u(x, t) = f(g(x, t))$  is a solution to the above advection equation for any smooth function  $f$ .

4. Consider the following advection equation

$$u_t + f(x)u_x = 0,$$

$$u(x, 0) = \frac{1}{1 + x^2},$$

where  $f(x)$  is plotted below:



- (a) Sketch a graph of  $u(x, 0)$  as a function of  $x$ .
- (b) Sketch a plot of the characteristic curves as a function of time  $t$ . Be sure to include enough of the curves to fully illustrate all possible types of curves.
- (c) Compute

$$u^*(x) = \lim_{t \rightarrow \infty} u(t, x)$$

and sketch a graph of  $u^*(x)$ .

5. Show that the advection-diffusion-decay equation

$$u_t = Du_{xx} - cu_x - \lambda u$$

can be transformed into the diffusion equation  $w_t = Dw_{xx}$  by the transformation

$$u(x, t) = w(x, t)e^{\alpha x - \beta t},$$

for some values of  $\alpha$  and  $\beta$ .

6. If  $z = x + iy$ , where  $x, y \in \mathbb{R}$ , find  $\operatorname{Re}(1/z)$  and  $\operatorname{Im}(1/z)$ .
7. If  $z$  is complex number, show that

$$\operatorname{Re}(iz) = -\operatorname{Im}(z) \text{ and } \operatorname{Im}(iz) = \operatorname{Re}(z).$$

8. Show that  $z\bar{z} = |z|^2$  for any complex number  $z$ .
9. Show that  $|z + w|^2 - |z - w|^2 = 4\operatorname{Re}(z\bar{w})$  for any complex numbers  $z, w$ .
10. For  $\theta \in \mathbb{R}$  prove that

$$\cos(\theta) = \frac{e^{i\theta} + e^{-i\theta}}{2} \text{ and } \sin(\theta) = \frac{e^{i\theta} - e^{-i\theta}}{2i}.$$

11. For  $n \in \mathbb{Z}$ , prove the following:

$$(\cos(\theta) + i \sin(\theta))^n = \cos(n\theta) + i \sin(n\theta).$$

12. Show that if  $m, n \in \mathbb{Z}$  then

$$\int_0^{2\pi} e^{in\theta} e^{-im\theta} d\theta = \begin{cases} 2\pi & \text{if } m = n \\ 0 & \text{if } m \neq n \end{cases}.$$

13. Calculate the Fourier transform of the “triangle function”:

$$f(x) = \begin{cases} x + 1 & \text{if } -1 \leq x \leq 0 \\ 1 - x & \text{if } 0 < x \leq 1 \\ 0 & \text{if } x < -1 \text{ or } x > 1 \end{cases}.$$

14. Find the Fourier transform of the function  $u$  defined by  $u(x) = e^{-ax}$  if  $x > 0$ , and  $u(x) = 0$  if  $x \leq 0$ .

15. Compute  $\mathcal{F}[xe^{-ax^2}](k)$ , where  $a > 0$  is a constant.

16. Given that  $\mathcal{F}[xe^{-|x|}](k) = \frac{4ik}{(1+k^2)^2}$ , find  $\mathcal{F}\left[\frac{x}{(1+x^2)^2}\right](k)$ .