

# Notes on quantum treatment of spin transitions

Reference: Charles P. Slichter, **Principles of Magnetic Resonance**, Harper & Row, 1963.

In the following, we will use the following Pauli spin matrices:

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (1)$$

In terms of these matrices, the electron magnetic moment can be written:

$$\vec{\mu} = -\frac{g\mu_B}{2}\vec{\sigma} \equiv \mu_e(\sigma_x\hat{x} + \sigma_y\hat{y} + \sigma_z\hat{z}), \quad (2)$$

where  $g$  denotes the electron g-factor,  $\mu_B = 5.788 \times 10^{-5}$  eV/T is the Bohr magneton, and  $\mu_e = -5.795 \times 10^{-5}$  eV/T is the electron magnetic moment. This same formalism can also treat nuclear spin-1/2 systems, with  $g_N\mu_N$  replacing  $-g\mu_B$ . The nuclear magneton  $\mu_N = 3.152 \times 10^{-8}$  eV/T and  $g_{proton} = 2.793$ ,  $g_{neutron} = -1.913$ . In the following, we write the explicit formulas in terms of the electron intrinsic magnetic moment, but the same formulation could be applied to other spin-1/2 systems as indicated. The Hamiltonian which represents the interaction between the electron magnetic moment and a magnetic field  $\mathbf{B}$  is given by

$$\mathcal{H} = -\vec{\mu} \cdot \mathbf{B}. \quad (3)$$

In magnetic resonance experiments, the magnetic field is generally composed of a constant component ( $B_0$ ) taken to be in the  $\hat{z}$  direction and a rotating component ( $B_1$ ) in the perpendicular direction taken to be in the  $x - y$  plane. Suppose that the rotation frequency is denoted by  $\Omega$ , the magnetic field can be written:

$$\mathbf{B} = B_1(\cos(\Omega t)\hat{x} + \sin(\Omega t)\hat{y}) + B_0\hat{z}, \quad (4)$$

where it is generally assumed that  $B_0 \gg B_1$ . For this field, the interaction Hamiltonian can be written:

$$\mathcal{H} = -\mu_e\mathbf{B} \cdot \vec{\sigma} \equiv -\mu_e \begin{pmatrix} B_0 & B_1e^{-i\Omega t} \\ B_1e^{i\Omega t} & -B_0 \end{pmatrix}. \quad (5)$$

We would like to solve the time-dependent Schrödinger equation:

$$i\hbar\frac{\partial\Psi(t)}{\partial t} = \mathcal{H}\Psi(t). \quad (6)$$

In order to simplify the mathematics, we notice that

$$\begin{pmatrix} B_0 & B_1e^{-i\Omega t} \\ B_1e^{i\Omega t} & -B_0 \end{pmatrix} = \begin{pmatrix} e^{-i\Omega t/2} & 0 \\ 0 & e^{i\Omega t/2} \end{pmatrix} \begin{pmatrix} B_0 & B_1 \\ B_1 & -B_0 \end{pmatrix} \begin{pmatrix} e^{i\Omega t/2} & 0 \\ 0 & e^{-i\Omega t/2} \end{pmatrix}. \quad (7)$$

Similarly, we can write,

$$i\hbar \frac{\partial \Psi(t)}{\partial t} = \begin{pmatrix} e^{-i\Omega t/2} & 0 \\ 0 & e^{i\Omega t/2} \end{pmatrix} \left\{ i\hbar \frac{\partial}{\partial t} - \begin{pmatrix} -\frac{\hbar\Omega}{2} & 0 \\ 0 & \frac{\hbar\Omega}{2} \end{pmatrix} \right\} \begin{pmatrix} e^{i\Omega t/2} & 0 \\ 0 & e^{-i\Omega t/2} \end{pmatrix} \Psi(t). \quad (8)$$

Defining a transformed wave function  $\Psi'$

$$\Psi' \equiv \begin{pmatrix} e^{i\Omega t/2} & 0 \\ 0 & e^{-i\Omega t/2} \end{pmatrix} \Psi(t), \quad (9)$$

the differential equation that must be solved to find the solutions to the Schrödinger equation can then be written:

$$i\hbar \frac{\partial \Psi'(t)}{\partial t} = \begin{pmatrix} -\mu_e B_0 - \frac{\hbar\Omega}{2} & -\mu_e B_1 \\ -\mu_e B_1 & -(-\mu_e B_0 - \frac{\hbar\Omega}{2}) \end{pmatrix} \Psi'(t) \equiv \mathcal{H}_{\text{eff}} \Psi'(t). \quad (10)$$

The transformed wave function  $\Psi'$  can be interpreted as representing the spin in a rotating coordinate system in which the effective Hamiltonian is now independent of time. Solving the differential equation (10), we find

$$\Psi'(t) = e^{-i\mathcal{H}_{\text{eff}}t/\hbar} \Psi'(0), \quad (11)$$

where  $\Psi'(0)$  denotes the initial value of the transformed wave function and where the exponential function must be evaluated by taking its Taylor series expansion. Consider a general  $2 \times 2$  matrix of the form

$$m \equiv \begin{pmatrix} a & b \\ b & -a \end{pmatrix}. \quad (12)$$

The exponential of  $m$  can be evaluated:

$$e^{-im} \equiv 1 - im - \frac{1}{2!}m^2 - \frac{1}{3!}m^2(-im) + \frac{1}{4!}(m^2)^2 \dots \quad (13)$$

For our form of  $m$ , all even terms are diagonal,

$$m^2 = \begin{pmatrix} a^2 + b^2 & 0 \\ 0 & a^2 + b^2 \end{pmatrix} \equiv (a^2 + b^2) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (14)$$

and all odd terms are proportional to  $m$  itself, so that we can simplify the expansion by summing the odd and even terms separately:

$$e^{-im} = \cos(\sqrt{a^2 + b^2}) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - i \frac{\sin(\sqrt{a^2 + b^2})}{\sqrt{a^2 + b^2}} \begin{pmatrix} a & b \\ b & -a \end{pmatrix}. \quad (15)$$

Defining the simplifying notation,  $\Omega_0 \equiv -\mu_e B_0/\hbar$ ,  $\Omega_1 \equiv -\mu_e B_1/\hbar$ ,  $\Omega_T \equiv \sqrt{(\Omega_0 - \Omega/2)^2 + \Omega_1^2}$ ,  $\cos(\theta_0) \equiv (\Omega_0 - \Omega/2)/\Omega_T$ , and  $\sin(\theta_0) \equiv \Omega_1/\Omega_T$ , we can write the full solution to the Schrödinger equation in the form

$$\Psi(t) = \begin{pmatrix} e^{-i\Omega t/2} & 0 \\ 0 & e^{i\Omega t/2} \end{pmatrix} \left\{ \cos(\Omega_T t) \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - i \sin(\Omega_T t) \begin{pmatrix} \cos(\theta_0) & \sin(\theta_0) \\ \sin(\theta_0) & -\cos(\theta_0) \end{pmatrix} \right\} \Psi(0). \quad (16)$$

For the special value of the rotational frequency  $\Omega = 2\Omega_0$ , the general result 16 simplifies to

$$\Psi(t) = \begin{pmatrix} e^{-i\Omega_0 t} & 0 \\ 0 & e^{i\Omega_0 t} \end{pmatrix} \begin{pmatrix} \cos(\Omega_1 t) & -i \sin(\Omega_1 t) \\ -i \sin(\Omega_1 t) & \cos(\Omega_1 t) \end{pmatrix} \Psi(0). \quad (17)$$

## PHY 741 – Assignment 17

1. Write down some of the intermediate steps of the above derivations. (Extra credit offered for any errors corrected.)
2. Suppose that  $\Psi(0) = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ , in other words, the system begins in the spin-up state. Assume that  $\Omega$  has been chosen at the resonant value (Eq. 17) and choose some reasonable value for  $\Omega_1$ . Evaluate and plot the probabilities as a function of time that the system is in the spin-up state ( $P_{\uparrow}(t)$ ), and that the system is in the spin-down state ( $P_{\downarrow}(t)$ ).