

PHY 113 A General Physics I
9-9:50 AM MWF Olin 101

Plan for Lecture 27:

Chapter 14: The physics of fluids

1. Review of static fluids

2. Bernoulli's equation

11/07/2012

PHY 113 A Fall 2012 -- Lecture 27

1

22	10/29/2012	Kepler's laws and satellite motion	13.4-13.6	13.28-13.34	10/31/2012
	10/31/2012	Review	10-13.15		
	11/02/2012	Exam	10-13.15		
23	11/05/2012	Fluid mechanics	14.1-14.4	14.8-14.24	11/07/2012
24	11/07/2012	Fluid mechanics	14.5-14.7	14.39-14.51	11/09/2012
25	11/09/2012	Temperature	19.1-19.5	19.1-19.20	11/12/2012
26	11/12/2012	Heat	20.1-20.4		11/14/2012
27	11/14/2012	First law of thermodynamics	20.5-20.7		11/16/2012
28	11/16/2012	Ideal gases	21.1-21.5		11/19/2012
29	11/19/2012	Engines	22.1-22.8		11/26/2012
	11/21/2012	Thanksgiving Holiday			
	11/23/2012	Thanksgiving Holiday			
	11/26/2012	Review	14.19-22		
	11/28/2012	Exam	14.19-22		
30	11/30/2012	Wave motion	18.1-18.6		12/03/2012
31	12/03/2012	Sound & standing waves	17.1-18.8		12/05/2012
	12/05/2012	Review	1-22		
	12/13/2012	Final Exam - 9 AM			

11/07/2012

PHY 113 A Fall 2012 -- Lecture 27

2

Review of equations describing static fluids in terms of pressure P and density ρ :

For all fluids near Earth's surface: $\frac{dP}{dy} = -\rho g$

For incompressible fluid, $\rho \equiv (\text{constant}) \Rightarrow P = P_0 - \rho g(y - y_0)$

For compressible fluid, equation is more complicated;

For an "ideal gas": $\rho = P \frac{\rho_0}{P_0} \Rightarrow \frac{dP}{dy} = -P \left(\frac{\rho_0 g}{P_0} \right)$

$$\Rightarrow P(y) = P_0 e^{-\frac{\rho_0 g}{P_0}(y-y_0)} \approx P_0 e^{-\frac{y-y_0}{8000m}} \approx P_0 e^{-\frac{y-y_0}{5mi}}$$

For small $y - y_0$, $P(y) \approx P_0 - \rho_0 g(y - y_0)$


11/07/2012

PHY 113 A Fall 2012 -- Lecture 27

3

Buoyant force

Buoyant force: $F_B = \rho_{\text{fluid}} g V_{\text{submerged}}$



Net force on cube from fluid :

$$F_B = F_{\text{bottom}} - F_{\text{top}} = \rho_{\text{fluid}} (A \Delta y) g$$

$$\Rightarrow F_B = \rho_{\text{fluid}} V_{\text{submerged}} g$$

11/07/2012

PHY 113 A Fall 2012 -- Lecture 27

4

Scale reading due to buoyant force:

Scale reading :

$$T + F_B - mg = 0$$

$$F_{\text{scale}} = T = mg - F_B$$


$$F_B = \rho_{\text{fluid}} V_{\text{submerged}} g$$

$$mg = \rho_{\text{solid}} V_{\text{solid}} g$$

If $\rho_{\text{fluid}} < \rho_{\text{solid}}$ $V_{\text{submerged}} = V_{\text{solid}}$

Measure solid density using :

$$\frac{\rho_{\text{solid}}}{\rho_{\text{fluid}}} = \frac{mg}{F_B} = \frac{mg}{mg - F_{\text{scale}}}$$



11/07/2012

PHY 113 A Fall 2012 -- Lecture 27

5

Scale reading due to buoyant force:

Scale reading :

$$T + F_B - mg = 0$$

$$F_{\text{scale}} = T = mg - F_B$$

$$F_B = \rho_{\text{fluid}} V_{\text{submerged}} g$$

$$mg = \rho_{\text{solid}} V_{\text{solid}} g$$

If $\rho_{\text{fluid}} > \rho_{\text{solid}}$ $V_{\text{submerged}} < V_{\text{solid}}$

$\Rightarrow T = 0$; Measure solid density using :


$$\frac{\rho_{\text{solid}}}{\rho_{\text{fluid}}} = \frac{V_{\text{submerged}}}{V_{\text{solid}}}$$

11/07/2012

PHY 113 A Fall 2012 -- Lecture 27

6

Mercury is poured into a U-tube as shown in Figure a. The left arm of the tube has cross-sectional area A_1 of 9.5 cm^2 , and the right arm has a cross-sectional area A_2 of 5.30 cm^2 . Four hundred grams of water are then poured into the right arm as shown in Figure b.




(a) Determine the length of the water column in the right arm of the U-tube.
_____ cm

(b) Given that the density of mercury is 13.6 g/cm^3 , what distance h does the mercury rise in the left arm?
_____ cm

11/07/2012

PHY 113 A Fall 2012 – Lecture 27

7



$$\begin{aligned} P_1 &= \rho_{H_2O} g \ell + \rho_{Hg} g \Delta z \\ &= \rho_{Hg} g (h + \Delta y + \Delta z) \\ \Rightarrow \rho_{H_2O} g \ell &= \rho_{Hg} g (h + \Delta y) \end{aligned}$$

$$\text{Note that: } h A_1 = \Delta y A_2$$

11/07/2012

PHY 113 A Fall 2012 – Lecture 27

8

$$\rho_{H_2O} g \ell = \rho_{Hg} g (h + \Delta y)$$

$$A_1 h = A_2 \Delta y$$

$$\Rightarrow \rho_{H_2O} g \ell = \rho_{Hg} g h \left(1 + \frac{A_1}{A_2} \right)$$

$$h = \ell \frac{\rho_{H_2O}}{\rho_{Hg} \left(1 + \frac{A_1}{A_2} \right)}$$

For $\ell = 0.2 \text{ m}$, $A_1 / A_2 = 2$:

$$h = 0.2 \text{ m} \frac{1000}{13600(1+2)} = 0.0049 \text{ m}$$


11/07/2012

PHY 113 A Fall 2012 – Lecture 27

9

Fluid dynamics (for incompressible fluids)

At $t = 0$, fluid in the blue portion is moving past point 1 at velocity $\vec{v}_1$.



After a time interval Δt , the fluid in the blue portion is moving past point 2 at velocity $\vec{v}_2$.

For an "incompressible" fluid :

$$\rho = (\text{constant})$$

Consider a given mass of fluid

$$M = \rho V_1 = \rho V_2$$

$$M = \rho A_1 \Delta x_1 = \rho A_2 \Delta x_2$$

$$M = \rho A_1 v_1 \Delta t = \rho A_2 v_2 \Delta t$$


$$\Rightarrow A_1 v_1 = A_2 v_2$$

10

11/07/2012

PHY 113 A Fall 2012 -- Lecture 27

10

Approximate energy conservation in fluid dynamics
→ **Bernoulli's equation**

Energy - work relationship on

"piece" of fluid M

$$K_i + U_i + W_{i \rightarrow f} = K_f + U_f$$

$$K_1 + U_1 + W_{1 \rightarrow 2} = K_2 + U_2$$

$$\frac{1}{2} M v_1^2 + M g y_1 - (P_2 - P_1) \Delta V = \frac{1}{2} M v_2^2 + M g y_2$$

$$M = \rho \Delta V$$

$$\frac{1}{2} \rho v_1^2 + \rho g y_1 - (P_2 - P_1) = \frac{1}{2} \rho v_2^2 + \rho g y_2$$

$$\frac{1}{2} \rho v_1^2 + \rho g y_1 + P_1 = \frac{1}{2} \rho v_2^2 + \rho g y_2 + P_2$$

11


11/07/2012

PHY 113 A Fall 2012 -- Lecture 27

11

Bernoulli's equation:

$$\frac{1}{2} \rho v_1^2 + \rho g y_1 + P_1 = \frac{1}{2} \rho v_2^2 + \rho g y_2 + P_2$$



$$y_1 = y_2 \quad A_1 v_1 = A_2 v_2$$

$$\frac{1}{2} \rho v_1^2 + P_1 = \frac{1}{2} \rho v_2^2 + P_2$$

$$\frac{1}{2} \rho v_2^2 \left(1 - \left(\frac{A_2}{A_1} \right)^2 \right) = P_1 - P_2$$

12

11/07/2012


PHY 113 A Fall 2012 -- Lecture 27

12

Bernoulli's equation:

$$\frac{1}{2} \rho v_1^2 + \rho g y_1 + P_1 = \frac{1}{2} \rho v_2^2 + \rho g y_2 + P_2$$

Point 2 is the surface of the liquid.



Point 1 is the exit point of the hole.

$$\frac{1}{2} \rho v_2^2 + \rho g y_2 + P =$$

$$\frac{1}{2} \rho v_1^2 + \rho g y_1 + P_0$$

$$A_2 v_2 = A_1 v_1$$

$$v_1 = \sqrt{\frac{2\rho g h + 2(P - P_0)}{\rho \left(1 - \left(\frac{A_1}{A_2}\right)^2\right)}}$$

11/07/2012

PHY 113 A Fall 2012 -- Lecture 27

13


Possibilities:**Fire extinguisher**

$$\rightarrow P \gg P_0$$

$$v_1 \approx \sqrt{\frac{2(P - P_0)}{\rho}}$$

$$\text{Example: } P = 10P_0$$

$$v_1 \approx \sqrt{\frac{18P_0}{\rho}} = \sqrt{\frac{18 \cdot 1.013 \times 10^5}{1000}} = 42.7 \text{ m/s}$$



11/07/2012

PHY 113 A Fall 2012 -- Lecture 27

14


Possibilities:**Open top:**

$$\rightarrow P = P_0$$

$$v_1 \approx \sqrt{2gh}$$

$$\text{Example: } h = 0.5 \text{ m}$$

$$v_1 \approx \sqrt{(2)(9.8)(0.5)} \text{ m/s} = 3.1 \text{ m/s}$$



11/07/2012

PHY 113 A Fall 2012 -- Lecture 27

15

$$v_1 = \sqrt{\frac{2\rho gh + 2(P - P_0)}{\rho \left(1 - \left(\frac{A_1}{A_2}\right)^2\right)}}$$

Point 2 is the surface of the liquid.

Point 1 is the exit point of the hole.

iclicker question:
Notice that if $P \ll P_0$, v_1 could become very small. How might this happen?

- Put an air-tight lid on the top.
- Remove the lid on the top.
- This will never happen.

11/07/2012 PHY 113 A Fall 2012 – Lecture 27 16

Siphon

$$\frac{1}{2} \rho v_A^2 + \rho gh + P_0 = \frac{1}{2} \rho v^2 + P_0$$

$$v = \sqrt{2gh}$$

11/07/2012 PHY 113 A Fall 2012 – Lecture 27 17

The air approaching from the right is deflected downward by the wing.

Simplistic statement:

$$\frac{1}{2} \rho v_1^2 + \rho g y_1 + P_1 = \frac{1}{2} \rho v_2^2 + \rho g y_2 + P_2$$

$$\frac{1}{2} \rho (v_1^2 - v_2^2) = P_2 - P_1$$

11/07/2012 PHY 113 A Fall 2012 – Lecture 27 18
