

**PHY 711 Classical Mechanics and
Mathematical Methods
10-10:50 AM MWF Olin 103**

Plan for Lecture 3:

**Chapter 1 – scattering theory
continued; center of mass versus
laboratory reference frame.**

9/3/2012

PHY 711 Fall 2012 -- Lecture 3

1

PHY 711 Classical Mechanics and Mathematical Methods

MWF 10 AM-10:50 PM OPL 103 <http://www.wfu.edu/~natalie/f12phy711/>

Instructor: Natalie Holzwarth | Phone: 758-5510 | Office: 300 OPL | e-mail: natalie@wfu.edu

Course schedule


(Preliminary schedule -- subject to frequent adjustment)			
Date	F&W Reading	Topic	Assignment
Wed. 8/29/2012	Chap. 1	Review of basic principles, Scattering theory	#1
Fri. 8/31/2012	Chap. 1	Scattering theory continued	#2
Mon. 9/03/2012	Chap. 1	Scattering theory continued	#3
Wed. 9/05/2012	Chap. 1	Scattering theory continued	#4

9/3/2012

PHY 711 Fall 2012 -- Lecture 3

2

Scattering geometry:



$$\left(\frac{d\sigma}{d\Omega} \right) = \frac{b}{\sin\theta} \left| \frac{db}{d\theta} \right|$$

9/3/2012

PHY 711 Fall 2012 -- Lecture 3

3

Relationship between scattering angle θ and impact parameter b for interaction potential $V(r)$:

$$\theta = \pi - 2b \int_{r_{\min}}^{\infty} dr \left(\frac{1/r^2}{\sqrt{1 - \frac{b^2}{r^2} - \frac{V(r)}{E}}} \right) \quad \text{where :}$$

$$1 - \frac{b^2}{r_{\min}^2} - \frac{V(r_{\min})}{E} = 0$$

$$\left(\frac{d\sigma}{d\Omega} \right) = \frac{b}{\sin \theta} \left| \frac{db}{d\theta} \right|$$

9/3/2012

PHY 711 Fall 2012 -- Lecture 3

4

Example of cross section analysis

Rutherford scattering :

$$\theta = 2 \sin^{-1} \left(\frac{1}{\sqrt{(b/\kappa)^2 + 1}} \right)$$

$$\frac{b}{\kappa} = \frac{|\cos(\theta/2)|}{|\sin(\theta/2)|}$$

$$\left(\frac{d\sigma}{d\Omega} \right) = \frac{b}{\sin \theta} \left| \frac{db}{d\theta} \right| = \frac{\kappa^2}{4} \frac{1}{\sin^4(\theta/2)}$$


9/3/2012

PHY 711 Fall 2012 -- Lecture 3

5

Example of cross section analysis

Hard sphere scattering:



For your homework you showed that

$$b = D \cos \left(\frac{\chi}{2} \right)$$

$$\left(\frac{d\sigma}{d\Omega} \right) = \frac{b}{\sin \chi} \left| \frac{db}{d\chi} \right| = \frac{D^2}{4}$$

9/3/2012


PHY 711 Fall 2012 -- Lecture 3

6


The results above were derived in the center of mass reference frame; relationship between normal laboratory reference and center of mass:

Laboratory reference frame:

Before




After




Center of mass reference frame:

Before



After



9/3/2012

PHY 711 Fall 2012 -- Lecture 3

7

Relationship between center of mass and laboratory frames of reference

Definition of center of mass \mathbf{R}_{CM}

$$m_1 \mathbf{r}_1 + m_2 \mathbf{r}_2 = (m_1 + m_2) \mathbf{R}_{CM}$$

$$m_1 \dot{\mathbf{r}}_1 + m_2 \dot{\mathbf{r}}_2 = (m_1 + m_2) \dot{\mathbf{R}}_{CM}$$

$$m_1 \mathbf{u}_1 + m_2 \mathbf{u}_2 = (m_1 + m_2) \mathbf{V}_{CM} = m_1 \mathbf{v}_1 + m_2 \mathbf{v}_2$$

In our case :

$$\mathbf{V}_{CM} = \frac{m_1}{m_1 + m_2} \mathbf{u}_1 = \frac{m_1 \mathbf{v}_1 + m_2 \mathbf{v}_2}{m_1 + m_2}$$

$$\mathbf{v}_1 = \mathbf{V}_1 + \mathbf{V}_{CM}$$

9/3/2012

PHY 711 Fall 2012 -- Lecture 3

8

Relationship between center of mass and laboratory frames of reference



Note that for an elastic collision

$$U_1 = V_1 \quad \text{and} \quad U_2 = V_2 = V_{CM}$$

Also note that : $m_1 U_1 = m_2 U_2$

$$\text{So that : } V_{CM}/V_1 = m_1/m_2$$

$$\begin{aligned} \mathbf{v}_1 &= \mathbf{V}_1 + \mathbf{V}_{CM} \\ v_1 \sin \psi &= V_1 \sin \theta \\ v_1 \cos \psi &= V_1 \cos \theta + V_{CM} \\ \tan \psi &= \frac{\sin \theta}{\cos \theta + V_{CM}/V_1} = \frac{\sin \theta}{\cos \theta + m_1/m_2} \end{aligned}$$

9/3/2012

PHY 711 Fall 2012 -- Lecture 3

9

Differential cross sections in different reference frames

$$\left(\frac{d\sigma_{LAB}(\psi)}{d\Omega_{LAB}} \right) = \left(\frac{d\sigma_{CM}(\theta)}{d\Omega_{CM}} \right) \frac{d\Omega_{CM}}{d\Omega_{LAB}}$$

$$\frac{d\Omega_{CM}}{d\Omega_{LAB}} = \left| \frac{\sin \theta}{\sin \psi} \frac{d\theta}{d\psi} \right| = \left| \frac{d \cos \theta}{d \cos \psi} \right|$$

From the previous result and/or conservation of momentum and energy, it is possible to show that :

$$\cos \psi = \frac{\cos \theta + m_1 / m_2}{\sqrt{1 + 2m_1 / m_2 \cos \theta + (m_1 / m_2)^2}}$$

9/3/2012

PHY 711 Fall 2012 -- Lecture 3

10

Differential cross sections in different reference frames – continued:

$$\left(\frac{d\sigma_{LAB}(\psi)}{d\Omega_{LAB}} \right) = \left(\frac{d\sigma_{CM}(\theta)}{d\Omega_{CM}} \right) \left| \frac{d \cos \theta}{d \cos \psi} \right|$$

$$\left(\frac{d\sigma_{LAB}(\psi)}{d\Omega_{LAB}} \right) = \left(\frac{d\sigma_{CM}(\theta)}{d\Omega_{CM}} \right) \frac{(1 + 2m_1 / m_2 \cos \theta + (m_1 / m_2)^2)^{3/2}}{(m_1 / m_2) \cos \theta + 1}$$

$$\text{where : } \tan \psi = \frac{\sin \theta}{\cos \theta + m_1 / m_2}$$

9/3/2012

PHY 711 Fall 2012 -- Lecture 3

11

$$\left(\frac{d\sigma_{LAB}(\psi)}{d\Omega_{LAB}} \right) = \left(\frac{d\sigma_{CM}(\theta)}{d\Omega_{CM}} \right) \frac{(1 + 2m_1 / m_2 \cos \theta + (m_1 / m_2)^2)^{3/2}}{(m_1 / m_2) \cos \theta + 1}$$

$$\text{where : } \tan \psi = \frac{\sin \theta}{\cos \theta + m_1 / m_2}$$

Example: suppose $m_1 = m_2$

$$\text{In this case : } \tan \psi = \frac{\sin \theta}{\cos \theta + 1} \Rightarrow \psi = \frac{\theta}{2}$$

$$\text{note that } 0 \leq \psi \leq \frac{\pi}{2}$$

$$\left(\frac{d\sigma_{LAB}(\psi)}{d\Omega_{LAB}} \right) = \left(\frac{d\sigma_{CM}(2\psi)}{d\Omega_{CM}} \right) \cdot 4 \cos \psi$$

9/3/2012

PHY 711 Fall 2012 -- Lecture 3

12
