

**PHY 711 Classical Mechanics and
Mathematical Methods**
10-10:50 AM MWF Olin 103

Plan for Lecture 5:

Start reading Chapter 3 –

**First focusing on the “calculus of
variation”**

9/4/2015

PHY 711 Fall 2015 – Lecture 5

1

Course schedule

(Preliminary schedule -- subject to frequent adjustment.)


Date	F&W Reading	Topic	Assignment
1 Wed, 8/26/2015	Chap. 1	Review of basic principles	#1
2 Fri, 8/28/2015	Chap. 1	Scattering theory	#2
3 Mon, 8/31/2015	Chap. 1	Scattering theory continued	#3
4 Wed, 9/02/2015	Chap. 2	Accelerated coordinate systems	#4
5 Fri, 9/04/2015	Chap. 3	Calculus of variations	#5
6 Mon, 9/07/2015	Chap. 3	Calculus of variations	#6
7 Wed, 9/09/2015	Chap. 3	Hamilton's principle	
8 Fri, 9/11/2015	Chap. 3 & 6	Hamilton's principle	
9 Mon, 9/14/2015	Chap. 3 & 6	Lagrangians with constraints	
10 Wed, 9/16/2015	Chap. 3 & 6	Lagrangians and constants of motion	
11 Fri, 9/18/2015	Chap. 3 & 6	Hamiltonian formalism	

9/4/2015

PHY 711 Fall 2015 – Lecture 5

2


In Chapter 3, the notion of Lagrangian dynamics is developed; reformulating Newton's laws in terms of minimization of related functions. In preparation, we need to develop a mathematical tool known as “the calculus of variation”.





9/4/2015

PHY 711 Fall 2015 – Lecture 5

3







Calculus of variation example for a pure integral functions

Find the function $y(x)$ which extremizes $L\left(\left\{y(x), \frac{dy}{dx}\right\}, x\right)$

where $L\left(\left\{y(x), \frac{dy}{dx}\right\}, x\right) \equiv \int_{x_i}^{x_f} f\left(\left\{y(x), \frac{dy}{dx}\right\}, x\right) dx$.

Necessary condition : $\delta L = 0$

At any x , let $y(x) \rightarrow y(x) + \delta y(x)$

$$\frac{dy(x)}{dx} \rightarrow \frac{dy(x)}{dx} + \delta \frac{dy(x)}{dx}$$

Formally :

$$\delta L = \int_{x_i}^{x_f} \left[\left(\frac{\partial f}{\partial y} \right)_{x, \frac{dy}{dx}} \delta y + \left[\left(\frac{\partial f}{\partial (dy/dx)} \right)_{x, y} \delta \left(\frac{dy}{dx} \right) \right] \right] dx.$$

9/4/2015

PHY 711 Fall 2015 – Lecture 5

7

After some derivations, we find

$$\begin{aligned} \delta L &= \int_{x_i}^{x_f} \left[\left(\frac{\partial f}{\partial y} \right)_{x, \frac{dy}{dx}} \delta y + \left[\left(\frac{\partial f}{\partial (dy/dx)} \right)_{x, y} \delta \left(\frac{dy}{dx} \right) \right] \right] dx \\ &= \int_{x_i}^{x_f} \left[\left(\frac{\partial f}{\partial y} \right)_{x, \frac{dy}{dx}} - \frac{d}{dx} \left[\left(\frac{\partial f}{\partial (dy/dx)} \right)_{x, y} \right] \right] \delta y dx = 0 \quad \text{for all } x_i \leq x \leq x_f \\ \Rightarrow \left(\frac{\partial f}{\partial y} \right)_{x, \frac{dy}{dx}} - \frac{d}{dx} \left[\left(\frac{\partial f}{\partial (dy/dx)} \right)_{x, y} \right] &= 0 \quad \text{for all } x_i \leq x \leq x_f \end{aligned}$$

9/4/2015

PHY 711 Fall 2015 – Lecture 5

8

Example : End points -- $y(0) = 0; y(1) = 1$

$$L = \int_0^1 \sqrt{1 + \left(\frac{dy}{dx} \right)^2} dx \Rightarrow f\left(\left\{y(x), \frac{dy}{dx}\right\}, x\right) = \sqrt{1 + \left(\frac{dy}{dx} \right)^2}$$

$$\left(\frac{\partial f}{\partial y} \right)_{x, \frac{dy}{dx}} - \frac{d}{dx} \left[\left(\frac{\partial f}{\partial (dy/dx)} \right)_{x, y} \right] = 0$$

$$\Rightarrow -\frac{d}{dx} \left(\frac{dy/dx}{\sqrt{1 + (dy/dx)^2}} \right) = 0$$

Solution :

$$\left(\frac{dy/dx}{\sqrt{1 + (dy/dx)^2}} \right) = K \quad \frac{dy}{dx} = K' \equiv \frac{K}{\sqrt{1 - K^2}}$$

$$\Rightarrow y(x) = x$$

9/4/2015

PHY 711 Fall 2015 – Lecture 5


9

Example :

$$A = 2\pi \int_{x_i}^{x_f} x \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx \Rightarrow f\left(\left\{y(x), \frac{dy}{dx}\right\}, x\right) = x \sqrt{1 + \left(\frac{dy}{dx}\right)^2}$$

$$\left(\frac{\partial f}{\partial y}\right)_{x, \frac{dy}{dx}} - \frac{d}{dx} \left[\left(\frac{\partial f}{\partial (dy/dx)}\right)_{x, y} \right] = 0$$

$$\Rightarrow -\frac{d}{dx} \left(\frac{x dy/dx}{\sqrt{1 + (dy/dx)^2}} \right) = 0$$



9/4/2015

PHY 711 Fa

$$-\frac{d}{dx} \left(\frac{x dy/dx}{\sqrt{1 + (dy/dx)^2}} \right) = 0$$

$$\frac{x dy/dx}{\sqrt{1 + (dy/dx)^2}} = K_1$$

$$\frac{dy}{dx} = -\frac{1}{\left(\frac{x}{K_1}\right)^2 - 1}$$

$$\Rightarrow y(x) = K_2 - K_1 \ln\left(x + \sqrt{x^2 - K_1^2}\right)$$

9/4/2015

PHY 711 Fall 2015 – Lecture 5

11

Another example:

(Courtesy of F. B. Hildebrand, Methods of Applied Mathematics)

Consider all curves $y(x)$ with $y(0)=0$ and $y(1)=1$ that minimize the integral :

$$I = \int_0^1 \left(\left(\frac{dy}{dx}\right)^2 - ay^2 \right) dx \quad \text{for constant } a > 0$$

Euler - Lagrange equation :

$$\frac{d^2 y}{dx^2} + ay = 0$$

$$\Rightarrow y(x) = \frac{\sin(\sqrt{a}x)}{\sin(\sqrt{a})}$$

9/4/2015

PHY 711 Fall 2015 – Lecture 5

12

Review: for $f\left(y(x), \frac{dy}{dx}, x\right)$,

a necessary condition to extremize $\int_{x_i}^{x_f} f\left(y(x), \frac{dy}{dx}, x\right) dx$:

$$\left(\frac{\partial f}{\partial y}\right)_{x, \frac{dy}{dx}} - \frac{d}{dx} \left[\left(\frac{\partial f}{\partial (dy/dx)}\right)_{x, y} \right] = 0$$

Note that for $f\left(y(x), \frac{dy}{dx}, x\right)$,

$$\begin{aligned} \frac{df}{dx} &= \left(\frac{\partial f}{\partial y}\right) \frac{dy}{dx} + \left(\frac{\partial f}{\partial (dy/dx)}\right) \frac{d}{dx} \frac{dy}{dx} + \left(\frac{\partial f}{\partial x}\right) \\ &= \left(\frac{d}{dx} \left(\frac{\partial f}{\partial (dy/dx)}\right)\right) \frac{dy}{dx} + \left(\frac{\partial f}{\partial (dy/dx)}\right) \frac{d}{dx} \frac{dy}{dx} + \left(\frac{\partial f}{\partial x}\right) \\ \Rightarrow \frac{d}{dx} \left(f - \frac{\partial f}{\partial (dy/dx)} \frac{dy}{dx} \right) &= \left(\frac{\partial f}{\partial x}\right) \end{aligned}$$


9/4/2015

PHY 711 Fall 2015 – Lecture 5

13

Brachistochrone problem: (solved by Newton in 1696)

<http://mathworld.wolfram.com/BrachistochroneProblem.html>



A particle of weight mg travels frictionlessly down a path of shape $y(x)$. What is the shape of the path $y(x)$ that minimizes the travel time from $y(0)=0$ to $y(\pi)=-2$?

9/4/2015

PHY 711 Fall 2015 – Lecture 5

14

$$T = \int_{x_i, y_i}^{x_f, y_f} \frac{ds}{v} = \int_{x_i}^{x_f} \frac{\sqrt{1 + \left(\frac{dy}{dx}\right)^2}}{\sqrt{2gy}} dx \quad \text{because } \frac{1}{2}mv^2 = mgy$$

$$f\left(y(x), \frac{dy}{dx}, x\right) = \sqrt{\frac{1 + \left(\frac{dy}{dx}\right)^2}{y}}$$

$$\frac{d}{dx} \left(f - \frac{\partial f}{\partial (dy/dx)} \frac{dy}{dx} \right) = 0$$

$$\frac{d}{dx} \left(\frac{1}{\sqrt{y \left(1 + \left(\frac{dy}{dx}\right)^2 \right)}} \right) = 0$$

Note that for the original form of Euler - Lagrange equation :

$$\left(\frac{\partial f}{\partial y}\right)_{x, \frac{dy}{dx}} - \frac{d}{dx} \left[\left(\frac{\partial f}{\partial (dy/dx)}\right)_{x, y} \right] = 0,$$

differential equation is more complicated :

$$-\frac{1}{2} \sqrt{\frac{1 + \left(\frac{dy}{dx}\right)^2}{y^3}} - \frac{d}{dx} \left(\frac{\frac{dy}{dx}}{\sqrt{y \left(1 + \left(\frac{dy}{dx}\right)^2 \right)}} \right) = 0$$

9/4/2015

PHY 711 Fall 2015 – Lecture 5

15

$$T = \int_{x_i, y_i}^{x_f, y_f} \frac{ds}{v} = \int_{x_i}^{x_f} \sqrt{\frac{1 + \left(\frac{dy}{dx}\right)^2}{-2gy}} dx \quad \text{because } \frac{1}{2}mv^2 = -mgy$$

$$f\left(\left\{y(x), \frac{dy}{dx}\right\}, x\right) = \sqrt{\frac{1 + \left(\frac{dy}{dx}\right)^2}{-y}}$$

Alternative relationships for extremization:

$$\left(\frac{\partial f}{\partial y}\right) - \frac{d}{dx} \left[\left(\frac{\partial f}{\partial (dy/dx)} \right) \right] = 0$$

$$\frac{d}{dx} \left(f - \frac{\partial f}{\partial (dy/dx)} \frac{dy}{dx} \right) = \left(\frac{\partial f}{\partial x} \right)$$

9/4/2015

PHY 711 Fall 2015 – Lecture 5

16

$$f\left(\left\{y(x), \frac{dy}{dx}\right\}, x\right) = \sqrt{\frac{1 + \left(\frac{dy}{dx}\right)^2}{-y}}$$

$$\frac{d}{dx} \left(f - \frac{\partial f}{\partial (dy/dx)} \frac{dy}{dx} \right) = \left(\frac{\partial f}{\partial x} \right)$$

$$\Rightarrow \frac{d}{dx} \left(\frac{1}{\sqrt{-y \left(1 + \left(\frac{dy}{dx} \right)^2 \right)}} \right) = 0 \quad -y \left(1 + \left(\frac{dy}{dx} \right)^2 \right) = K \equiv 2a$$

9/4/2015

PHY 711 Fall 2015 – Lecture 5

17

$$-y \left(1 + \left(\frac{dy}{dx} \right)^2 \right) = K \equiv 2a \quad \text{Let } y = -2a \sin^2 \frac{\theta}{2} = a(\cos \theta - 1)$$

$$\frac{dy}{dx} = -\frac{\sqrt{2a}}{\sqrt{-y}} \quad -\frac{dy}{\sqrt{\frac{2a}{-y}} - 1} = \frac{2a \sin \frac{\theta}{2} \cos \frac{\theta}{2} d\theta}{\sqrt{\frac{2a}{2a \sin^2 \frac{\theta}{2}} - 1}} = dx$$

$$-\frac{dy}{\sqrt{\frac{2a}{-y}} - 1} = dx \quad x = \int_0^\theta a(1 - \cos \theta') d\theta' = a(\theta - \sin \theta)$$

Parametric equations for Brachistochrone:

$$x = a(\theta - \sin \theta)$$

$$y = a(\cos \theta - 1)$$

9/4/2015

PHY 711 Fall 2015 – Lecture 5

18

