

**PHY 711 Classical Mechanics and
Mathematical Methods
11-11:50 AM MWF Olin 107**

Plan for Lecture 11:

Continue reading Chapter 3 & 6

- 1. Hamiltonian formalism**
- 2. Phase space & Liouville's theorem**
- 3. Modern applications**

9/26/2016

PHY 711 Fall 2016 -- Lecture 11

1

Course schedule

(Preliminary schedule -- subject to frequent adjustment.)

Date	F&W Reading	Topic	Assignment Due	
1 Wed, 8/31/2016	Chap. 1	Review of basic principles	#1	9/7/2016
2 Fri, 9/02/2016	Chap. 1	Scattering theory	#2	9/7/2016
Mon, 9/05/2016		Labor day -- no class		
3 Wed, 9/07/2016	Chap. 1	Scattering theory	#3	9/9/2016
4 Fri, 9/09/2016	Chap. 1 & 2	Scattering theory and rotations	#4	9/12/2016
5 Mon, 9/12/2016	Chap. 3	Calculus of variations	#5	9/14/2016
6 Wed, 9/14/2016	Chap. 3	Calculus of variations	#6	9/16/2016
7 Fri, 9/16/2016	Chap. 3	Lagrangian mechanics	#7	9/19/2016
8 Mon, 9/19/2016	Chap. 3 and 6	Lagrangian mechanics and constraints	#8	9/21/2016
9 Wed, 9/21/2016	Chap. 3 and 6	Constants of the motion	#9	9/23/2016
10 Fri, 9/23/2016	Chap. 3 and 6	Hamiltonian and canonical equations of motion	#10	9/26/2016
11 Mon, 9/26/2016	Chap. 3 and 6	Phase space	#11	9/28/2016
12 Wed, 9/28/2016				
13 Fri, 9/30/2016				
14 Mon, 10/03/2016				
15 Wed, 10/05/2016				

9/26/2016

PHY 711 Fall 2016 -- Lecture 11

2

Hamiltonian formalism

$$H = H(\{q_\sigma(t)\}, \{p_\sigma(t)\}, t)$$

Canonical equations of motion

$$\frac{dq_\sigma}{dt} = \frac{\partial H}{\partial p_\sigma}$$

$$\frac{dp_\sigma}{dt} = -\frac{\partial H}{\partial q_\sigma}$$

9/26/2016

PHY 711 Fall 2016 -- Lecture 11

3

$$\frac{dD}{dt} = 0$$

Importance of Liouville's theorem to statistical mechanical analysis:

In statistical mechanics, we need to evaluate the probability of various configurations of particles. The fact that the density of particles in phase space is constant in time, implies that each point in phase space is equally probable and that the time average of the evolution of a system can be determined by an average of the system over phase space volume.

9/26/2016

PHY 711 Fall 2016 -- Lecture 11

4

Modern usage of Lagrangian and Hamiltonian formalisms

J. Chem. Physics **72** 2384-2393 (1980)

Molecular dynamics simulations at constant pressure and/or temperature¹⁾

Hans C. Andersen

Department of Chemistry, Stanford University, Stanford, California 94305
(Received 10 July 1979; accepted 31 October 1979)

In the molecular dynamics simulation method for fluids, the equations of motion for a collection of particles in a fixed volume are solved numerically. The energy, volume, and number of particles are constant for a particular simulation, and it is assumed that time averages of properties of the simulated fluid are equal to microcanonical ensemble averages of the same properties. In some situations, it is desirable to perform simulations of a fluid for particular values of temperature and/or pressure or under conditions in which the energy and volume of the fluid can fluctuate. This paper proposes and discusses three methods for performing molecular dynamics simulations under conditions of constant temperature and/or pressure, rather than constant energy and volume. For these three methods, it is shown that time averages of properties of the simulated fluid are equal to averages over the isenthalpic-isobaric, canonical, and isothermal-isobaric ensembles. Each method is a way of describing the dynamics of a certain number of particles in a volume element of a fluid while taking into account the influence of surrounding particles in changing the energy and/or density of the simulated volume element. The influence of the surroundings is taken into account without introducing unwanted surface effects. Examples of situations where these methods may be useful are discussed.

9/26/2016

PHY 711 Fall 2016 -- Lecture 11

5

"Molecular dynamics" is a subfield of computational physics focused on analyzing the motions of atoms in fluids and solids with the goal of relating the atomistic and macroscopic properties of materials. Ideally molecular dynamics calculations can numerically realize the statistical mechanics viewpoint.

Imagine that the generalized coordinates $q_{\sigma}(t)$ represent N atoms, each with 3 spacial coordinates:

$$L = L(\{q_{\sigma}(t)\}, \{\dot{q}_{\sigma}(t)\}, t) = T - U$$


For simplicity, it is assumed that the potential interaction is a sum of pairwise interactions:

$$U(\mathbf{x}^N) = \sum_{\langle i,j \rangle} u(r_{ij}) . \quad (2.1)$$

9/26/2016

PHY 711 Fall 2016 -- Lecture 11

6



$$L = L(\{\mathbf{r}_i(t)\}, \{\dot{\mathbf{r}}_i(t)\}) = \sum_i \frac{1}{2} m_i |\dot{\mathbf{r}}_i|^2 - \sum_{i < j} u(|\mathbf{r}_i - \mathbf{r}_j|)$$

→ From this Lagrangian, can find the 3N coupled 2nd order differential equations of motion and/or find the corresponding Hamiltonian, representing the system at constant energy, volume, and particle number N (N,V,E ensemble).

9/26/2016 PHY 711 Fall 2016 -- Lecture 11 7

Lagrangian and Hamiltonian forms

$$L = L(\{\mathbf{r}_i(t)\}, \{\dot{\mathbf{r}}_i(t)\}) = \sum_i \frac{1}{2} m_i |\dot{\mathbf{r}}_i|^2 - \sum_{i < j} u(|\mathbf{r}_i - \mathbf{r}_j|)$$

$$\mathbf{p}_i = m_i \dot{\mathbf{r}}_i$$


$$H = \sum_i \frac{|\mathbf{p}_i|^2}{2m_i} + \sum_{i < j} u(|\mathbf{r}_i - \mathbf{r}_j|)$$

Canonical equations :

$$\frac{d\mathbf{r}_i}{dt} = \frac{\mathbf{p}_i}{m_i} \quad \frac{d\mathbf{p}_i}{dt} = - \sum_{i < j} u'(|\mathbf{r}_i - \mathbf{r}_j|) \frac{\mathbf{r}_i - \mathbf{r}_j}{|\mathbf{r}_i - \mathbf{r}_j|}$$

9/26/2016 PHY 711 Fall 2016 -- Lecture 11 8

H. C. Andersen wanted to adapt the formalism for modeling an (N,V,E) ensemble to one which could model a system at constant pressure (P).



V constant

P constant, V variable

9/26/2016 PHY 711 Fall 2016 -- Lecture 11 9

Andersen's clever transformation :

Let $\mathbf{p}_i = \mathbf{r}_i / Q^{1/3}$

$L = L(\{\mathbf{r}_i(t)\}, \{\dot{\mathbf{r}}_i(t)\}) = \sum_i \frac{1}{2} m_i |\dot{\mathbf{r}}_i|^2 - \sum_{i < j} u(|\mathbf{r}_i - \mathbf{r}_j|)$

$L = L(\{\mathbf{p}_i(t)\}, \{\dot{\mathbf{p}}_i(t)\}, Q, \dot{Q}) = Q^{2/3} \sum_i \frac{1}{2} m_i |\dot{\mathbf{p}}_i|^2 - \sum_{i < j} u(Q^{1/3} |\mathbf{p}_i - \mathbf{p}_j|) + \frac{1}{2} M \dot{Q}^2 - \alpha Q$

PV contribution to potential energy

kinetic energy of "balloon"

9/26/2016 PHY 711 Fall 2016 -- Lecture 11 10

$L = L(\{\mathbf{p}_i(t)\}, \{\dot{\mathbf{p}}_i(t)\}, Q, \dot{Q}) = Q^{2/3} \sum_i \frac{1}{2} m_i |\dot{\mathbf{p}}_i|^2 - \sum_{i < j} u(Q^{1/3} |\mathbf{p}_i - \mathbf{p}_j|) + \frac{1}{2} M \dot{Q}^2 - \alpha Q$

$\boldsymbol{\pi}_i = \frac{\partial L}{\partial \dot{\mathbf{p}}_i} = m Q^{2/3} \dot{\mathbf{p}}_i$

$\Pi = \frac{\partial L}{\partial \dot{Q}} = M \dot{Q}$

$H = \sum_i \frac{|\boldsymbol{\pi}_i|^2}{2 m_i Q^{2/3}} + \sum_{i < j} u(Q^{1/3} |\mathbf{p}_i - \mathbf{p}_j|) + \frac{\Pi^2}{2M} + \alpha Q$

$\frac{d\mathbf{p}_i}{dt} = \frac{\boldsymbol{\pi}_i}{2 m_i Q^{2/3}} \quad \frac{dQ}{dt} = \frac{\Pi}{M}$

$\frac{d\boldsymbol{\pi}_i}{dt} = -Q^{1/3} \sum_{i < j} u' (Q^{1/3} |\mathbf{p}_i - \mathbf{p}_j|) \frac{\mathbf{p}_i - \mathbf{p}_j}{|\mathbf{p}_i - \mathbf{p}_j|}$

$\frac{d\Pi}{dt} = \frac{2}{3Q} \sum_i \frac{|\boldsymbol{\pi}_i|^2}{2 m_i Q^{2/3}} - \frac{1}{3Q^{2/3}} \sum_{i < j} u' (Q^{1/3} |\mathbf{p}_i - \mathbf{p}_j|) |\mathbf{p}_i - \mathbf{p}_j| - \alpha$

9/26/2016 PHY 711 Fall 2016 -- Lecture 11 11

Relationship between system representations

Scaled	=	Original
$Q(t)$	=	$V(t)$
$Q^{1/3} \mathbf{p}_i(t)$	=	$\mathbf{r}_i(t)$
$\boldsymbol{\pi}_i / Q^{1/3}$	=	\mathbf{p}_i

Equations of motion in "original" coordinates:

$\frac{d\mathbf{r}_i}{dt} = \frac{\mathbf{p}_i}{m_i} + \frac{1}{3} \mathbf{r}_i \frac{d \ln V}{dt}$

$\frac{d\mathbf{p}_i}{dt} = - \sum_{j < i} \frac{\mathbf{r}_i - \mathbf{r}_j}{|\mathbf{r}_i - \mathbf{r}_j|} u'(|\mathbf{r}_i - \mathbf{r}_j|) - \frac{1}{3} \mathbf{p}_i \frac{d \ln V}{dt}$

$M \frac{d^2 V}{dt^2} = -\alpha + \frac{1}{V} \left(\frac{2}{3} \sum_i \frac{\mathbf{p}_i \cdot \mathbf{p}_i}{m_i} - \frac{1}{3} \sum_{j < i} |\mathbf{r}_i - \mathbf{r}_j| u'(|\mathbf{r}_i - \mathbf{r}_j|) \right)$

9/26/2016 PHY 711 Fall 2016 -- Lecture 11 12

Physical interpretation:

$\alpha \Leftrightarrow$ Imposed (target) pressure

$$\frac{1}{V} \left(\frac{2}{3} \sum_i \frac{\mathbf{p}_i \cdot \mathbf{p}_i}{m_i} - \frac{1}{3} \sum_{j < i} |\mathbf{r}_i - \mathbf{r}_j| u'(|\mathbf{r}_i - \mathbf{r}_j|) \right) \Leftrightarrow \text{Internal pressure of system}$$

Time dependence

$$M \frac{d^2 V}{dt^2} = -\alpha + \frac{1}{V} \left(\frac{2}{3} \sum_i \frac{\mathbf{p}_i \cdot \mathbf{p}_i}{m_i} - \frac{1}{3} \sum_{j < i} |\mathbf{r}_i - \mathbf{r}_j| u'(|\mathbf{r}_i - \mathbf{r}_j|) \right)$$

9/26/2016 PHY 711 Fall 2016 -- Lecture 11 13

Digression on numerical evaluation of differential equations

Example differential equation (one dimension);

$$\frac{d^2 x}{dt^2} = f(t) \quad \text{Let } t = nh \quad (n = 1, 2, 3, \dots)$$

$$x_n \equiv x(nh); \quad f_n \equiv f(nh)$$

Euler's method :

$$x_{n+1} = x_n + hv_n + \frac{1}{2} h^2 f_n$$

$$v_{n+1} = v_n + hf_n$$


Velocity Verlet algorithm :

$$x_{n+1} = x_n + hv_n + \frac{1}{2} h^2 f_n$$

$$v_{n+1} = v_n + \frac{1}{2} h(f_n + f_{n+1})$$

9/26/2016 PHY 711 Fall 2016 -- Lecture 11 14

Example simulation for NPT molecular dynamics
simulation of Li₂O using 1500 atoms with $\theta=0$




Pair interaction potential

$$u_{ij}(r_{ij}) = A_{ij} e^{\rho_{ij}/r_{ij}} - \frac{C_{ij}}{r_{ij}^6} + \frac{q_i q_j}{r_{ij}}$$

Use LAMMPS code
<http://LAMMPS.sandia.gov>

9/26/2016 PHY 711 Fall 2016 -- Lecture 11 15



MOLECULAR PHYSICS, 1984, VOL. 52, No. 2, 255-268

A molecular dynamics method for simulations in the canonical ensemble†

by SHŪICHI NOSÉ†

Division of Chemistry, National Research Council Canada,
Ottawa, Ontario, Canada K1A 0R6

(Received 3 October 1983 ; accepted 28 November 1983)

A molecular dynamics simulation method which can generate configurations belonging to the canonical (T, V, N) ensemble or the constant temperature constant pressure (T, P, N) ensemble, is proposed. The physical system of interest consists of N particles (f degrees of freedom), to which an external, macroscopic variable and its conjugate momentum are added. This device allows the total energy of the physical system to fluctuate. The equilibrium distribution of the energy coincides with the

Nose's Lagrangian:

$$L(\{\mathbf{r}_i\}, s, \{\dot{\mathbf{r}}_i\}, \dot{s}) = \frac{1}{2} \sum_i m_i s^2 \dot{\mathbf{r}}_i^2 + \frac{1}{2} Q \dot{s}^2 - \phi(\{\mathbf{r}_i\}) - (f+1)kT_{eq} \ln s$$

↑
↑
 velocity scaling fictitious mass

Equations of motion:

$$\frac{d}{dt} (m_i s^2 \dot{\mathbf{r}}_i) = -\frac{\partial \phi}{\partial \mathbf{r}_i}$$

$$\ddot{\mathbf{r}}_i = -\frac{1}{m_i s^2} \frac{\partial \phi}{\partial \mathbf{r}_i} - \frac{2\dot{s}}{s} \dot{\mathbf{r}}_i$$

$$Q \ddot{s} = \sum_i m_i s \dot{\mathbf{r}}_i^2 - \frac{(f+1)kT_{eq}}{s}$$

Time averaged relationships

$$\left\langle \frac{\sum_i m_i s^2 \dot{r}_i^2}{s} \right\rangle = (f+1)kT_{eq} \left\langle \frac{1}{s} \right\rangle$$

Hamiltonian

$$\mathcal{H}_1 = \sum_i \frac{\mathbf{P}_i^2}{2m_i s^2} + \phi(\mathbf{r}) + \frac{\dot{p}_s^2}{2Q} + (f+1)kT_{eq} \ln s,$$

9/26/2016

PHY 711 Fall 2016 -- Lecture 11

19

In statistical mechanics, the thermodynamic functions can be analyzed in terms of a partition function. A canonical partition function for a system with N particles at a temperature T_{eq} can be determined from the phase space integral:

$$Z = \frac{1}{N!} \int d^{3N}r d^{3N}p e^{-\mathcal{H}(\mathbf{r}, \mathbf{p})/kT_{eq}}$$


Nose' was able to show that his effective Hamiltonian well approximates such a canonical distribution.

9/26/2016

PHY 711 Fall 2016 -- Lecture 11

20

From LAMMPS simulation (using modified Nose' algorithm)



9/26/2016

PHY 711 Fall 2016 -- Lecture 11

21
