

PHY 711 Classical Mechanics and Mathematical Methods

11-11:50 AM MWF Olin 107

Plan for Lecture 22:

Motions of elastic membranes (Chap. 8)

1. Review of standing waves on a string
2. Standing waves on a two dimensional membrane.
3. Boundary value problems

10/24/2016

PHY 711 Fall 2016 -- Lecture 22

1



20 Mon, 10/17/2016	Chap. 5	Mechanics of rigid bodies	Exam due	
21 Wed, 10/19/2016	Chap. 5	Mechanics of rigid bodies	#16	10/24/2016
Fri, 10/21/2016		Fall break -- no class		
22 Mon, 10/24/2016	Chap. 8	Mechanics of Elastic Membranes	#17	10/28/2016
23 Wed, 10/26/2016				
24 Fri, 10/28/2016				
25 Mon, 10/31/2016				
26 Wed, 11/02/2016				
27 Fri, 11/04/2016				
28 Mon, 11/07/2016				
29 Wed, 11/09/2016				
30 Fri, 11/11/2016				
31 Mon, 11/14/2016				
32 Wed, 11/16/2016				
33 Fri, 11/18/2016				
34 Mon, 11/21/2016				
Wed, 11/23/2016		Thanksgiving Holiday -- no class		
Fri, 11/25/2016		Thanksgiving Holiday -- no class		
Wed, 12/07/2016		Presentations I		
Fri, 12/09/2016		Presentations II		

10/24/2016

PHY 711 Fall 2016 -- Lecture 22

2

Elastic media in two or more dimensions --

Review of wave equation in one-dimension -- here $\mu(x,t)$ can describe either a longitudinal or transverse wave.

Traveling wave solutions --

$$\frac{\partial^2 \mu}{\partial t^2} - c^2 \frac{\partial^2 \mu}{\partial x^2} = 0$$

Note that for any function $f(q)$ or $g(q)$:

$$\mu(x,t) = f(x-ct) + g(x+ct)$$

satisfies the wave equation.

10/24/2016

PHY 711 Fall 2016 -- Lecture 22

3

Initial value problem : $\mu(x,0) = \phi(x)$ and $\frac{\partial \mu}{\partial t}(x,0) = \psi(x)$

then : $\mu(x,0) = \phi(x) = f(x) + g(x)$

$$\frac{\partial \mu}{\partial t}(x,0) = \psi(x) = -c \left(\frac{df(x)}{dx} - \frac{dg(x)}{dx} \right)$$

$$\Rightarrow f(x) - g(x) = -\frac{1}{c} \int \psi(x') dx'$$

For each x , find $f(x)$ and $g(x)$:

$$f(x) = \frac{1}{2} \left(\phi(x) - \frac{1}{c} \int \psi(x') dx' \right)$$


$$g(x) = \frac{1}{2} \left(\phi(x) + \frac{1}{c} \int \psi(x') dx' \right)$$

$$\Rightarrow \mu(x,t) = \frac{1}{2} (\phi(x-ct) + \phi(x+ct)) + \frac{1}{2c} \int_{x-ct}^{x+ct} \psi(x') dx'$$

10/24/2016

PHY 711 Fall 2016 -- Lecture 22

4



10/24/2016

PHY 711 Fall 2016 -- Lecture 22

5

Standing wave solutions of wave equation:

$$\frac{\partial^2 \mu}{\partial t^2} - c^2 \frac{\partial^2 \mu}{\partial x^2} = 0$$

with $\mu(0,t) = \mu(L,t) = 0$.

Assume: $\mu(x,t) = \Re(e^{-i\omega t} \rho(x))$

where $\frac{d^2 \rho(x)}{dx^2} + k^2 \rho(x) = 0$ $k = \frac{\omega}{c}$


$$\rho_v(x) = A \sin\left(\frac{v\pi x}{L}\right)$$

$$k_v = \frac{v\pi}{L} \quad \omega_v = ck_v$$

10/24/2016

PHY 711 Fall 2016 -- Lecture 22

6



Wave motion on a two-dimensional surface – elastic membrane (transverse wave; linear regime).

Two-dimensional wave equation:

$$\frac{\partial^2 u}{\partial t^2} - c^2 \nabla^2 u = 0 \quad \text{where } c^2 = \frac{\tau}{\sigma}$$


Standing wave solutions:

$$u(x, y, t) = \Re(e^{-i\omega t} \rho(x, y))$$

$$(\nabla^2 + k^2)\rho(x, y) = 0$$

$$\text{where } k = \frac{\omega}{c}$$

$u(x, y, t)$



Lagrangian density: $\mathcal{L}\left(u, \frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial u}{\partial t}; x, y, t\right)$

$$L = \int \mathcal{L}\left(u, \frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial u}{\partial t}; x, y, t\right) dx dy$$

Hamilton's principle:

$$\delta \int_{t_1}^{t_2} L dt = 0$$

$$\frac{\partial \mathcal{L}}{\partial u} - \frac{\partial}{\partial t} \frac{\partial \mathcal{L}}{\partial (\partial u / \partial t)} - \frac{\partial}{\partial x} \frac{\partial \mathcal{L}}{\partial (\partial u / \partial x)} - \frac{\partial}{\partial y} \frac{\partial \mathcal{L}}{\partial (\partial u / \partial y)} = 0$$

10/24/2016

PHY 711 Fall 2016 -- Lecture 22

9

Lagrangian density for elastic membrane with constant σ and τ :

$$\mathcal{L}\left(u, \frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial u}{\partial t}; x, y, t\right) = \frac{1}{2}\sigma\left(\frac{\partial u}{\partial t}\right)^2 - \frac{1}{2}\tau(\nabla u)^2$$

$$\frac{\partial \mathcal{L}}{\partial u} - \frac{\partial}{\partial t} \frac{\partial \mathcal{L}}{\partial(\partial u / \partial t)} - \frac{\partial}{\partial x} \frac{\partial \mathcal{L}}{\partial(\partial u / \partial x)} - \frac{\partial}{\partial y} \frac{\partial \mathcal{L}}{\partial(\partial u / \partial y)} = 0$$

$$\frac{\partial^2 u}{\partial t^2} - c^2 \nabla^2 u = 0 \quad \text{where } c^2 = \frac{\tau}{\sigma}$$

Two - dimensional wave equation :

$$\frac{\partial^2 u}{\partial t^2} - c^2 \nabla^2 u = 0 \quad \text{where } c^2 = \frac{\tau}{\sigma}$$

Standing wave solutions :

$$u(x, y, t) = \Re(e^{-i\omega t} \rho(x, y))$$

$$(\nabla^2 + k^2)\rho(x, y) = 0 \quad \text{where } k = \frac{\omega}{c}$$

10/24/2016

PHY 711 Fall 2016 -- Lecture 22

10

Consider a rectangular boundary:



Clamped boundary conditions :

$$\rho(0, y) = \rho(a, y) = \rho(x, 0) = \rho(x, b) = 0 \quad (\nabla^2 + k^2)\rho(x, y) = 0$$


$$\Rightarrow \rho_{mn}(x, y) = A \sin\left(\frac{m\pi x}{a}\right) \sin\left(\frac{n\pi y}{b}\right) \quad \text{where } k = \frac{\omega}{c}$$

$$k_{mn}^2 = \left(\frac{m\pi}{a}\right)^2 + \left(\frac{n\pi}{b}\right)^2 \quad \omega_{mn} = ck_{mn}$$

10/24/2016

PHY 711 Fall 2016 -- Lecture 22

11



10/24/2016

PHY 711 Fall 2016 -- Lecture 22

12

More general boundary conditions:

$\tau \nabla u|_b = \kappa u|_b$ represents boundary side constrained with spring

$\tau \nabla u|_b = 0$ represents "free" side


Mixed boundary conditions:

$$\rho(x,0) = \rho(x,b) = \frac{\partial \rho(0,y)}{\partial x} = \frac{\partial \rho(a,y)}{\partial x} = 0$$

$$\Rightarrow \rho_{mn}(x,y) = A \cos\left(\frac{m\pi x}{a}\right) \sin\left(\frac{n\pi y}{b}\right)$$

$$k_{mn}^2 = \left(\frac{m\pi}{a}\right)^2 + \left(\frac{n\pi}{b}\right)^2 \quad \omega_{mn} = ck_{mn}$$

$$k_{11}^2 = \left(\frac{\pi}{a}\right)^2 + \left(\frac{\pi}{b}\right)^2$$




10/24/2016

PHY 711 Fall 2016 -- Lecture 22

Consider a circular boundary:

Clamped boundary conditions for $\rho(r, \varphi)$:

$$\rho(R, \varphi) = 0$$



$$(\nabla^2 + k^2)\rho(r, \varphi) = 0 \quad \text{where } k = \frac{\omega}{c}$$

In cylindrical coordinate system

$$\nabla^2 = \frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \frac{\partial^2}{\partial \varphi^2}$$

Assume: $\rho(r, \varphi) = f(r)\Phi(\varphi)$

Let: $\Phi(\varphi) = e^{im\varphi}$

Note: $\Phi(\varphi) = \Phi(\varphi + 2\pi)$

$$\Rightarrow m = \text{integer}$$

10/24/2016

PHY 711 Fall 2016 -- Lecture 22

14

Consider circular boundary -- continued


Differential equation for radial function:

$$\left(\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{m^2}{r^2} + k^2\right)f(r) = 0$$

$\Rightarrow$ Bessel equation of integer order with transcendental solutions

Cylindrical Bessel function $J_m(z)$

Cylindrical Neumann function $N_m(z)$ also called $Y_m(z)$



10/24/2016

PHY 711 Fall 2016 -- Lecture 22

15

Some properties of Bessel functions

Ascending series: $J_m(z) = \left(\frac{z}{2}\right)^m \sum_{j=0}^{\infty} \frac{(-1)^j}{j!(j+m)!} \left(\frac{z}{2}\right)^{2j}$

Recursion relations: $J_{m-1}(z) + J_{m+1}(z) = \frac{2m}{z} J_m(z)$

$$J_{m-1}(z) - J_{m+1}(z) = 2 \frac{dJ_m(z)}{dz}$$

Asymptotic form: $J_m(z) \xrightarrow{z \gg 1} \sqrt{\frac{2}{\pi z}} \cos\left(z - \frac{m\pi}{2} - \frac{\pi}{4}\right)$

Zeros of Bessel functions $J_m(z_{mn}) = 0$

$m=0$: $z_{0n} = 2.406, 5.520, 8.654, \dots$


$m=1$: $z_{1n} = 3.832, 7.016, 10.173, \dots$

$m=2$: $z_{2n} = 5.136, 8.417, 11.620, \dots$

10/24/2016

PHY 711 Fall 2016 -- Lecture 22

16

<http://dlmf.nist.gov/>


NIST Digital Library of Mathematical Functions

Project News

2014-08-29 DLMF Update, Version 1.0.9
 2014-04-21 DLMF Update, Version 1.0.8, errata & improved MathML
 2014-03-31 DLMF Update, Version 1.0.7, New Functions, Improved Math & 3D Graphics
 2013-08-16 DLMF Update, Version 1.0.6, New Functions, Improved Math & 3D Graphics
[More news](#)

Foreword

Preface	19 Elliptic Integrals
Mathematical Introduction	20 Theta Functions
1 Algebraic and Analytic Methods	21 Multidimensional Theta Functions
2 Asymptotic Approximations	22 Jacobian Elliptic Functions
3 Numerical Methods	23 Weierstrass Elliptic and Modular Functions
4 Elementary Functions	24 Bernoulli and Euler Polynomials
5 Gamma Function	25 Zeta and Related Functions
6 Exponential, Logarithmic, Sine, and Cosine Integrals	26 Combinatorial Analysis
7 Error Functions, Dawson's and Fresnel Integrals	27 Functions of Number Theory
8 Incomplete Gamma and Related Functions	28 Mathieu Functions and Hill's Equation
9 Airy and Related Functions	29 Lamé Functions
10 Bessel Functions	30 Spheroidal Wave Functions
11 Struve and Related Functions	31 Heun Functions
12 Parabolic Cylinder Functions	32 Painlevé Transcendents
	33 Coulomb Functions

10/24/2016

PHY 711 Fall 2016 -- Lecture 22

17

Series expansions of Bessel and Neumann functions

$$J_\nu(z) = \left(\frac{1}{2}z\right)^\nu \sum_{k=0}^{\infty} \frac{(-1)^k}{k! \Gamma(\nu + k + 1)} \left(\frac{1}{4}z^2\right)^k$$

$$Y_n(z) = -\frac{\left(\frac{1}{2}z\right)^{-n}}{\pi} \sum_{k=0}^{n-1} \frac{(n-k-1)!}{k!} \left(\frac{1}{4}z^2\right)^k + \frac{2}{\pi} \ln\left(\frac{1}{2}z\right) J_n(z) - \frac{\left(\frac{1}{2}z\right)^n}{\pi} \sum_{k=0}^{\infty} \left(\psi(k+1) + \psi(n+k+1)\right) \frac{\left(-\frac{1}{4}z^2\right)^k}{k!(n+k)!}$$

10/24/2016

PHY 711 Fall 2016 -- Lecture 22

18

Some properties of Bessel functions -- continued

Note : It is possible to prove the following

identity for the functions $J_m\left(\frac{z_{mn}}{R}r\right)$:

$$\int_0^R J_m\left(\frac{z_{mn}}{R}r\right) J_m\left(\frac{z_{mn'}}{R}r\right) r dr = \frac{R^2}{2} (J_{m+1}(z_{mn}))^2 \delta_{nn'}$$

Returning to differential equation for radial function :

$$\left(\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{m^2}{r^2} + k^2 \right) f(r) = 0$$


$$\Rightarrow f_{mn}(r) = A J_m\left(\frac{z_{mn}}{R}r\right); \quad k_{mn} = \frac{z_{mn}}{R}$$

10/24/2016

PHY 711 Fall 2016 -- Lecture 22


19

$$\rho_{01}(r, \varphi) = f_{01}(r) = A J_0\left(\frac{z_{01}}{R}r\right)$$



$$k_{01} = \frac{2.406}{R}$$

$$\rho_{02}(r, \varphi) = f_{02}(r) = A J_0\left(\frac{z_{02}}{R}r\right)$$



$$k_{02} = \frac{5.520}{R}$$


10/24/2016

PHY 711 Fall 2016 -- Lecture 22

20

$$\rho_{11}(r, \varphi) = f_{11}(r) \cos(\varphi)$$


$$= A J_1\left(\frac{z_{11}}{R}r\right) \cos(\varphi)$$



$$k_{11} = \frac{3.832}{R}$$

$$\rho_{12}(r, \varphi) = f_{12}(r) \cos(\varphi)$$

$$= A J_1\left(\frac{z_{12}}{R}r\right) \cos(\varphi)$$




$$k_{12} = \frac{7.016}{R}$$

10/24/2016

PHY 711 Fall 2016 -- Lecture 22

21

More complicated geometry – annular membrane



In cylindrical coordinate system

$$\nabla^2 = \frac{\partial^2}{\partial r^2} + \frac{1}{r} \frac{\partial}{\partial r} + \frac{1}{r^2} \frac{\partial^2}{\partial \varphi^2}$$

Assume: $\rho(r, \varphi) = f(r)\Phi(\varphi)$ Let: $\Phi(\varphi) = e^{im\varphi}$ Note: $\Phi(\varphi) = \Phi(\varphi + 2\pi)$
 $\Rightarrow m = \text{integer}$

10/24/2016


PHY 711 Fall 2016 -- Lecture 22

22

Consider circular boundary -- continued

Differential equation for radial function:

$$\left(\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{m^2}{r^2} + k^2 \right) f(r) = 0$$


 $\Rightarrow$ Bessel equation of integer order with transcendental solutionsCylindrical Bessel function $J_m(z)$ Cylindrical Neumann function $N_m(z)$ 

10/24/2016

PHY 711 Fall 2016 -- Lecture 22

23

Normal modes of an annular membrane -- continued



Differential equation for radial function:

$$\left(\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{m^2}{r^2} + k^2 \right) f(r) = 0$$


General form of radial function: $f(r) = AJ_m(kr) + BN_m(kr)$

10/24/2016

PHY 711 Fall 2016 -- Lecture 22

24

Normal modes of an annular membrane -- continued



Boundary conditions:
 $f(a) = 0$ $f(b) = 0$

$$AJ_m(ka) + BN_m(ka) = 0$$

$$AJ_m(kb) + BN_m(kb) = 0$$

$\Rightarrow$ 2 equations and 2 unknowns -- k and $\frac{B}{A}$


$$\frac{B}{A} = \frac{-J_m(ka)}{N_m(ka)} = \frac{-J_m(kb)}{N_m(kb)} \quad (\text{transcendental equation for } k)$$

10/24/2016

PHY 711 Fall 2016 -- Lecture 22

25

Normal modes of an annular membrane -- continued



Boundary conditions:
 $f(a) = 0$ $f(b) = 0$

$$\frac{B}{A} = \frac{-J_m(ka)}{N_m(ka)} = \frac{-J_m(kb)}{N_m(kb)} \quad \text{-- in terms of solution } k_{mn}:$$

$$f(r) = A \left(J_m(k_{mn}r) - \frac{J_m(k_{mn}a)}{N_m(k_{mn}a)} N_m(k_{mn}r) \right)$$


10/24/2016

PHY 711 Fall 2016 -- Lecture 22

26

Analysis for $m=0$ and $a=0.1, b=0.2$:

```
> plot( [ [-BesselJ(0, 0.1*k), -BesselJ(0, 0.2*k)], k=25..33, color=[red, blue] ];
```



10/24/2016

PHY 711 Fall 2016 -- Lecture 22

27

