

**PHY 711 Classical Mechanics and
Mathematical Methods**
11-11:50 AM MWF Olin 107

Plan for Lecture 2:

1. Brief comment on quiz
2. Particle interactions
3. Notion of center of mass
reference frame
4. Introduction to scattering theory

9/02/2016

PHY 711 Fall 2016 -- Lecture 2

1

PHY 711 Classical Mechanics and Mathematical Methods

MWF 11 AM-11:50 AM OPL 107 <http://www.wfu.edu/~natalie/f16phy711/>

Instructor: [Natalie Hotzwarth](#) Phone: 758-5510 Office: 300 OPL e-mail: natalie@wfu.edu

Course schedule

(Preliminary schedule -- subject to frequent adjustment.)



Date	F&W Reading	Topic	Assignment Due
1 Wed, 8/31/2016	Chap. 1	Review of basic principles	#1 9/7/2016
2 Fri, 9/02/2016	Chap. 1	Scattering theory	#2 9/7/2016
Mon, 9/05/2016		Labor day -- no class	
3 Wed, 9/07/2016	Chap. 1	Scattering theory	9/9/2016
4 Fri, 9/09/2016			
5 Mon, 9/12/2016			
6 Wed, 9/14/2016			
7 Fri, 9/16/2016			

9/02/2016

PHY 711 Fall 2016 -- Lecture 2

2

PHY 711 -- Assignment #2

Sept. 2, 2016

Read Chapter 1 in **Fetter & Walecka**.

1. In class, we "derived" the differential cross section for the scattering of two hard spheres of mutual radius D in the center of mass frame. Find the differential cross section for this system in the lab frame in which m_{target} is initially at rest and evaluate the expression for the following cases.

- a. $m_1/m_{\text{target}}=0.1$
- b. $m_1/m_{\text{target}}=1$
- c. $m_1/m_{\text{target}}=1000$

9/02/2016

PHY 711 Fall 2016 -- Lecture 2

3

Fall 2016 Schedule for <u>N. A. W. Holzwarth</u>					
	Monday	Tuesday	Wednesday	Thursday	Friday
9:00-11:00	Lecture Preparation/ Office Hours		Lecture Preparation/ Office Hours		Lecture Preparation/ Office Hours
11:00-12:00	Classical Mechanics PHY711	Physics Research	Classical Mechanics PHY711	Physics Research	Classical Mechanics PHY711
12:00-1:15			Physics Research		Condensed Matter Theory Journal Club
1:15-3:30	Physics Research		Physics Colloquium		Physics Research
3:30-5:00					

Schedule additional office hours by email:
natalie@wfu.edu

9/02/2016 PHY 711 Fall 2016 – Lecture 2 4


Comment on quiz questions

- $$g(t) = \int_0^t (x^2 + t) dx \quad \frac{dg}{dt} = \int_0^t \frac{d}{dt} (x^2 + t) dt + (x^2 + t) \Big|_{x=t}$$

$$= \int_0^t dt + (t^2 + t) = t^2 + 2t$$
- Evaluate the integral $\oint \frac{dz}{z}$ for a closed contour about the origin.

Suppose that $z = e^{i\theta} \quad dz = e^{i\theta} i d\theta \quad \oint \frac{dz}{z} = \int_0^{2\pi} \frac{e^{i\theta} i d\theta}{e^{i\theta}} = 2\pi i$
- $\frac{df}{dx} = f \quad \Rightarrow f(x) = Ae^x \quad f(x) = 1 \quad \Rightarrow A = 1$
- $\sum_{n=1}^N a^n = \frac{a - a^{N+1}}{1 - a} \quad \text{Let } S \equiv \sum_{n=1}^N a^n \quad \text{Note that } aS - S = a^{N+1} - a$


9/02/2016 PHY 711 Fall 2016 – Lecture 2 5



Differential cross section

$$\left(\frac{d\sigma}{d\Omega}\right) = \frac{\text{Number of detected particles at } \theta \text{ per target particle}}{\text{Number of incident particles per unit area}}$$

= Area of incident beam that is scattered into detector at angle θ



$$\left(\frac{d\sigma}{d\Omega}\right) = \frac{d\phi b db}{d\phi \sin\theta d\theta} = \frac{b}{\sin\theta} \left|\frac{db}{d\theta}\right|$$

Figure from Marion & Thorton, Classical Dynamics

9/02/2016

PHY 711 Fall 2016 – Lecture 2

7

Note: Notion of cross section is common to many areas of physics including classical mechanics, quantum mechanics, optics, etc. Only in the classical mechanics can we calculate it using geometric considerations




Figure from Marion & Thorton, Classical Dynamics

$$\left(\frac{d\sigma}{d\Omega}\right) = \frac{d\phi b db}{d\phi \sin\theta d\theta} = \frac{b}{\sin\theta} \left|\frac{db}{d\theta}\right|$$


Note: We are assuming that the process is isotropic in ϕ

9/02/2016

PHY 711 Fall 2016 – Lecture 2


8

Simple example – collision of hard spheres



$$\left(\frac{d\sigma}{d\Omega}\right) = \frac{b}{\sin\theta} \left|\frac{db}{d\theta}\right|$$

Microscopic view:



$$b(\theta) = ?$$

$$b(\theta) = D \sin\left(\frac{\pi}{2} - \frac{\theta}{2}\right)$$


$$\left(\frac{d\sigma}{d\Omega}\right) = \frac{D^2}{4}$$

9/02/2016

PHY 711 Fall 2016 – Lecture 2

9


Simple example – collision of hard spheres -- continued



Total scattering cross section:

$$\sigma = \int \left(\frac{d\sigma}{d\Omega} \right) d\Omega$$

Hard sphere:




$$\left(\frac{d\sigma}{d\Omega} \right) = \frac{D^2}{4}$$

$$\sigma = \pi D^2$$

9/02/2016 PHY 711 Fall 2016 -- Lecture 2 10

Relationship of scattering cross-section to particle interactions --
Classical mechanics of a conservative 2-particle system.



$$\frac{d\mathbf{p}_1}{dt} = \mathbf{F}_{12} \quad \frac{d\mathbf{p}_2}{dt} = \mathbf{F}_{21}$$

$$\mathbf{F}_{12} = -\nabla_1 V(\mathbf{r}_1 - \mathbf{r}_2) \Rightarrow E = \frac{1}{2} m_1 v_1^2 + \frac{1}{2} m_2 v_2^2 + V(\mathbf{r}_1 - \mathbf{r}_2)$$

9/02/2016 PHY 711 Fall 2016 -- Lecture 2 11

Typical two-particle interactions –

Central potential: $V(\mathbf{r}_1 - \mathbf{r}_2) = V(|\mathbf{r}_1 - \mathbf{r}_2|) \equiv V(r)$

Hard sphere: $V(r) = \begin{cases} \infty & r \leq a \\ 0 & r > a \end{cases}$

Coulomb or gravitational: $V(r) = \frac{K}{r}$

Lennard-Jones: $V(r) = \frac{A}{r^{12}} - \frac{B}{r^6}$

9/02/2016 PHY 711 Fall 2016 -- Lecture 2 12

Relationship between center of mass and laboratory frames of reference

Definition of center of mass $\mathbf{R}_{CM}$

$$m_1 \mathbf{r}_1 + m_2 \mathbf{r}_2 = (m_1 + m_2) \mathbf{R}_{CM}$$

$$m_1 \dot{\mathbf{r}}_1 + m_2 \dot{\mathbf{r}}_2 = (m_1 + m_2) \dot{\mathbf{R}}_{CM} = (m_1 + m_2) \mathbf{V}_{CM}$$

$$E = \frac{1}{2} m_1 v_1^2 + \frac{1}{2} m_2 v_2^2 + V(\mathbf{r}_1 - \mathbf{r}_2)$$

$$= \frac{1}{2} (m_1 + m_2) V_{CM}^2 + \frac{1}{2} \mu |\mathbf{v}_1 - \mathbf{v}_2|^2 + V(\mathbf{r}_1 - \mathbf{r}_2)$$

where: $\mu \equiv \frac{m_1 m_2}{m_1 + m_2}$

9/02/2016

PHY 711 Fall 2016 -- Lecture 2

13

Classical mechanics of a conservative 2-particle system -- continued

$$E = \frac{1}{2} (m_1 + m_2) V_{CM}^2 + \frac{1}{2} \mu |\mathbf{v}_1 - \mathbf{v}_2|^2 + V(\mathbf{r}_1 - \mathbf{r}_2)$$

For central potentials: $V(\mathbf{r}_1 - \mathbf{r}_2) = V(|\mathbf{r}_1 - \mathbf{r}_2|) \equiv V(r_{12})$

Relative angular momentum is also conserved:

$$\mathbf{L}_{12} \equiv \mathbf{r}_{12} \times \mu \mathbf{v}_{12}$$

$$E = \frac{1}{2} (m_1 + m_2) V_{CM}^2 + \frac{1}{2} \mu v_{12}^2 + \frac{L_{12}^2}{2\mu r_{12}^2} + V(r_{12})$$

Simpler notation:

$$E = \frac{1}{2} (m_1 + m_2) V_{CM}^2 + \frac{1}{2} \mu \dot{r}^2 + \frac{\ell^2}{2\mu r^2} + V(r)$$


9/02/2016

PHY 711 Fall 2016 -- Lecture 2

14

Note: The following analysis will be carried out in the center of mass frame of reference.


In laboratory frame:



$$\mu = \frac{m_1 m_{\text{target}}}{m_1 + m_{\text{target}}}$$

$$\ell = |\mathbf{r} \times \mu \mathbf{v}_1|$$

In center-of-mass frame:




Also note: We are assuming that the interaction between particle and target $V(r)$ conserves energy and angular momentum.

9/02/2016

PHY 711 Fall 2016 -- Lecture 2

15

For a continuous potential interaction in center of mass reference frame:



9/02/2016


PHY 711 Fall 2016 – Lecture 2

16


The results above were derived in the center of mass reference frame; relationship between normal laboratory reference and center of mass:

Laboratory reference frame:

Before




After




Center of mass reference frame:

Before



After



9/02/2016

PHY 711 Fall 2016 – Lecture 2

17

Relationship between center of mass and laboratory frames of reference -- continued

Since m_2 is initially at rest :

$$\mathbf{V}_{CM} = \frac{m_1}{m_1 + m_2} \mathbf{u}_1 \quad \mathbf{u}_1 = \mathbf{U}_1 + \mathbf{V}_{CM} \Rightarrow \mathbf{U}_1 = \frac{m_2}{m_1 + m_2} \mathbf{u}_1 = \frac{m_2}{m_1} \mathbf{V}_{CM}$$

$$\mathbf{u}_2 = \mathbf{U}_2 + \mathbf{V}_{CM} \Rightarrow \mathbf{U}_2 = -\frac{m_1}{m_1 + m_2} \mathbf{u}_1 = -\mathbf{V}_{CM}$$

$$\mathbf{v}_1 = \mathbf{V}_1 + \mathbf{V}_{CM}$$


$$\mathbf{v}_2 = \mathbf{V}_2 + \mathbf{V}_{CM}$$

9/02/2016

PHY 711 Fall 2016 – Lecture 2

18

Relationship between center of mass and laboratory frames of reference



$$\mathbf{v}_1 = \mathbf{V}_1 + \mathbf{V}_{CM}$$

$$v_1 \sin \psi = V_1 \sin \theta$$

$$v_1 \cos \psi = V_1 \cos \theta + V_{CM}$$

$$\tan \psi = \frac{\sin \theta}{\cos \theta + V_{CM} / V_1} = \frac{\sin \theta}{\cos \theta + m_1 / m_2} \quad \text{For elastic scattering}$$

9/02/2016

PHY 711 Fall 2016 – Lecture 2

19

Digression – elastic scattering

$$\begin{aligned} \frac{1}{2} m_1 U_1^2 + \frac{1}{2} m_2 U_2^2 + \frac{1}{2} (m_1 + m_2) V_{CM}^2 \\ = \frac{1}{2} m_1 V_1^2 + \frac{1}{2} m_2 V_2^2 + \frac{1}{2} (m_1 + m_2) V_{CM}^2 \end{aligned}$$

Also note:

$$m_1 \mathbf{U}_1 + m_2 \mathbf{U}_2 = 0 \quad m_1 \mathbf{V}_1 + m_2 \mathbf{V}_2 = 0$$

$$\mathbf{U}_1 = \frac{m_2}{m_1} \mathbf{V}_{CM} \quad \mathbf{U}_2 = -\mathbf{V}_{CM}$$

$$\Rightarrow |\mathbf{U}_1| = |\mathbf{V}_1| \quad \text{and} \quad |\mathbf{U}_2| = |\mathbf{V}_2| = |\mathbf{V}_{CM}|$$

$$\text{Also note that: } m_1 |\mathbf{U}_1| = m_2 |\mathbf{U}_2|$$


$$\text{So that: } V_{CM} / V_1 = V_{CM} / U_1 = m_1 / m_2$$

9/02/2016

PHY 711 Fall 2016 – Lecture 2

20

Relationship between center of mass and laboratory frames of reference – continued (elastic scattering)



$$\mathbf{v}_1 = \mathbf{V}_1 + \mathbf{V}_{CM}$$

$$v_1 \sin \psi = V_1 \sin \theta$$

$$v_1 \cos \psi = V_1 \cos \theta + V_{CM}$$

$$\tan \psi = \frac{\sin \theta}{\cos \theta + V_{CM} / V_1} = \frac{\sin \theta}{\cos \theta + m_1 / m_2}$$

$$\text{Also: } \cos \psi = \frac{\cos \theta + m_1 / m_2}{\sqrt{1 + 2m_1 / m_2 \cos \theta + (m_1 / m_2)^2}}$$

9/02/2016

PHY 711 Fall 2016 – Lecture 2

21

Differential cross sections in different reference frames

$$\left(\frac{d\sigma_{LAB}(\psi)}{d\Omega_{LAB}} \right) = \left(\frac{d\sigma_{CM}(\theta)}{d\Omega_{CM}} \right) \frac{d\Omega_{CM}}{d\Omega_{LAB}}$$

$$\frac{d\Omega_{CM}}{d\Omega_{LAB}} = \left| \frac{\sin \theta}{\sin \psi} \frac{d\theta}{d\psi} \right| = \left| \frac{d \cos \theta}{d \cos \psi} \right|$$

Using :

$$\cos \psi = \frac{\cos \theta + m_1 / m_2}{\sqrt{1 + 2(m_1 / m_2) \cos \theta + (m_1 / m_2)^2}}$$

$$\left| \frac{d \cos \psi}{d \cos \theta} \right| = \frac{(m_1 / m_2) \cos \theta + 1}{(1 + 2(m_1 / m_2) \cos \theta + (m_1 / m_2)^2)^{3/2}}$$

9/02/2016

PHY 711 Fall 2016 – Lecture 2

22

Differential cross sections in different reference frames – continued:

$$\left(\frac{d\sigma_{LAB}(\psi)}{d\Omega_{LAB}} \right) = \left(\frac{d\sigma_{CM}(\theta)}{d\Omega_{CM}} \right) \left| \frac{d \cos \theta}{d \cos \psi} \right|$$

$$\left(\frac{d\sigma_{LAB}(\psi)}{d\Omega_{LAB}} \right) = \left(\frac{d\sigma_{CM}(\theta)}{d\Omega_{CM}} \right) \frac{(1 + 2m_1 / m_2 \cos \theta + (m_1 / m_2)^2)^{3/2}}{(m_1 / m_2) \cos \theta + 1}$$

where : $\tan \psi = \frac{\sin \theta}{\cos \theta + m_1 / m_2}$

9/02/2016

PHY 711 Fall 2016 – Lecture 2

23

$$\left(\frac{d\sigma_{LAB}(\psi)}{d\Omega_{LAB}} \right) = \left(\frac{d\sigma_{CM}(\theta)}{d\Omega_{CM}} \right) \frac{(1 + 2m_1 / m_2 \cos \theta + (m_1 / m_2)^2)^{3/2}}{(m_1 / m_2) \cos \theta + 1}$$

where : $\tan \psi = \frac{\sin \theta}{\cos \theta + m_1 / m_2}$

Example: suppose $m_1 = m_2$

In this case : $\tan \psi = \frac{\sin \theta}{\cos \theta + 1} \Rightarrow \psi = \frac{\theta}{2}$

note that $0 \leq \psi \leq \frac{\pi}{2}$

$$\left(\frac{d\sigma_{LAB}(\psi)}{d\Omega_{LAB}} \right) = \left(\frac{d\sigma_{CM}(2\psi)}{d\Omega_{CM}} \right) \cdot 4 \cos \psi$$

9/02/2016

PHY 711 Fall 2016 – Lecture 2

24