

PHY 711 Classical Mechanics and Mathematical Methods 10-10:50 AM MWF online

Discussion for Lecture 33: Chap. 9 of F&W

Wave equation for sound beyond the linear approximation

- 1. Non-linear effects in sound waves**
- 2. Shock wave analysis**

27	Mon, 10/26/2020	Chap. 9	Mechanics of 3 dimensional fluids	#18	10/30/2020
28	Wed, 10/28/2020	Chap. 9	Mechanics of 3 dimensional fluids		
29	Fri, 10/30/2020	Chap. 9	Linearized hydrodynamics equations	#19	11/02/2020
30	Mon, 11/02/2020	Chap. 9	Linear sound waves	#20	11/04/2020
31	Wed, 11/04/2020	Chap. 9	Linear sound waves	Project topic	11/06/2020
32	Fri, 11/06/2020	Chap. 9	Sound sources and scattering; Non linear effects		
33	Mon, 11/09/2020	Chap. 9	Non linear effects in sound waves and shocks	#21	11/11/2020
34	Wed, 11/11/2020	Chap. 10	Surface waves in fluids		
35	Fri, 11/13/2020	Chap. 10	Surface waves in fluids; soliton solutions		
36	Mon, 11/16/2020	Chap. 11	Heat conduction		
37	Wed, 11/18/2020	Chap. 12	Viscous effects		
38	Fri, 11/20/2020	Chap. 13	Elasticity		
39	Mon, 11/23/2020		Review		
	Wed, 11/25/2020		Thanksgiving Holidaya		
	Fri, 11/27/2020		Thanksgiving Holidaya		
40	Mon, 11/30/2020		Review		
	Wed, 12/02/2020		Presentations I		
	Fri, 12/04/2020		Presentations II		

PHY 711 -- Assignment #21

Nov. 09, 2020

Finish reading Chapter 9 in **Fetter & Walecka**.

1. In class, we discussed how to visualize the non-linear behavior of an adiabatic ideal gas with parameter γ . Using Maple or Mathematica or other software and using a parametric plot formalism, create an animated gif file to show the traveling waveform $s(w)$, where s is a shape of your choice and $w = x - u(s(w))t$. You will also need to choose the value of γ as well.

Schedule for weekly one-on-one meetings (EST)

Nick – 11 AM Monday

Tim – 9 AM Tuesday

Gao – 9 PM Tuesday

Jeanette – 11 AM Friday

Derek – 12 PM Friday

Your questions –

From Tim –

1. Is γ = degrees of freedom?

From Gao –

1. Why can we assume ρ is a function of $(x-u(\rho)t)$?

Effects of nonlinearities in fluid equations

-- one dimensional case

Newton - Euler equation of motion :

$$\frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{v} = \mathbf{f}_{\text{applied}} - \frac{\nabla p}{\rho}$$

$$\text{Continuity equation : } \frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{v}) = 0$$

Assume spatial variation confined to x direction ;

assume that $\mathbf{v} = v \hat{\mathbf{x}}$ and $\mathbf{f}_{\text{applied}} = 0$.

$$\frac{\partial v}{\partial t} + v \frac{\partial v}{\partial x} = - \frac{1}{\rho} \frac{\partial p}{\partial x}$$

$$\frac{\partial \rho}{\partial t} + v \frac{\partial \rho}{\partial x} + \rho \frac{\partial v}{\partial x} = 0$$

$$\frac{\partial v}{\partial t} + v \frac{\partial v}{\partial x} + \frac{1}{\rho} \frac{\partial p}{\partial x} = 0$$

$$\frac{\partial \rho}{\partial t} + v \frac{\partial \rho}{\partial x} + \rho \frac{\partial v}{\partial x} = 0$$

Expressing p in terms of ρ : $p = p(\rho)$

$$\frac{\partial p}{\partial x} = \frac{\partial p}{\partial \rho} \frac{\partial \rho}{\partial x} \equiv c^2(\rho) \frac{\partial \rho}{\partial x} \quad \text{where} \quad \frac{\partial p}{\partial \rho} \equiv c^2(\rho)$$

For adiabatic ideal gas:

$$\frac{dp}{p} = \gamma \frac{d\rho}{\rho} \quad p = p_0 \left(\frac{\rho}{\rho_0} \right)^\gamma$$

$$c^2(\rho) = \frac{\gamma p}{\rho} = c_0^2 \left(\frac{\rho}{\rho_0} \right)^{\gamma-1} \quad \text{where} \quad c_0^2 \equiv \frac{\gamma p_0}{\rho_0}$$

Digression – What is gamma?

Internal energy for ideal gas: $pV = Nk_B T$

$$E_{\text{int}} = \frac{f}{2} Nk_B T \quad f \equiv \text{degrees of freedom; } 3 \text{ for atom, } 5 \text{ for diatomic molecule}$$

In terms of specific heat ratio: $\gamma \equiv \frac{C_p}{C_V}$

$$dE_{\text{int}} = dQ - dW$$

$$C_V = \left(\frac{dQ}{dT} \right)_V = \left(\frac{\partial E}{\partial T} \right)_V = \frac{f}{2} Nk_B$$

$$C_p = \left(\frac{dQ}{dT} \right)_p = \left(\frac{\partial E}{\partial T} \right)_p + p \left(\frac{\partial V}{\partial T} \right)_p = \frac{f}{2} Nk_B + Nk_B$$

$$\gamma = \frac{C_p}{C_V} = \frac{\frac{f}{2} + 1}{\frac{f}{2}} = 1 + \frac{2}{f} \quad \Rightarrow \quad \frac{f}{2} = \frac{1}{\gamma - 1} \quad E_{\text{int}} = \frac{1}{\gamma - 1} Nk_B$$

$$\frac{\partial v}{\partial t} + v \frac{\partial v}{\partial x} + \frac{c^2(\rho)}{\rho} \frac{\partial \rho}{\partial x} = 0$$

$$\frac{\partial \rho}{\partial t} + v \frac{\partial \rho}{\partial x} + \rho \frac{\partial v}{\partial x} = 0$$

Expressing variation of v in terms of $v(\rho)$:

$$\frac{\partial v}{\partial \rho} \frac{\partial \rho}{\partial t} + v \frac{\partial v}{\partial \rho} \frac{\partial \rho}{\partial x} + \frac{c^2(\rho)}{\rho} \frac{\partial \rho}{\partial x} = 0$$

$$\frac{\partial \rho}{\partial t} + v \frac{\partial \rho}{\partial x} + \rho \frac{\partial v}{\partial \rho} \frac{\partial \rho}{\partial x} = 0$$

Some more algebra :

$$\text{From Euler equation : } \frac{\partial v}{\partial \rho} \left(\frac{\partial \rho}{\partial t} + v \frac{\partial \rho}{\partial x} \right) + \frac{c^2(\rho)}{\rho} \frac{\partial \rho}{\partial x} = 0$$

$$\text{From continuity equation : } \frac{\partial \rho}{\partial t} + v \frac{\partial \rho}{\partial x} = -\rho \frac{\partial v}{\partial \rho} \frac{\partial \rho}{\partial x}$$

$$\text{Combined equation : } \frac{\partial v}{\partial \rho} \left(-\rho \frac{\partial v}{\partial \rho} \frac{\partial \rho}{\partial x} \right) + \frac{c^2(\rho)}{\rho} \frac{\partial \rho}{\partial x} = 0$$

$$\Rightarrow \left(\frac{\partial v}{\partial \rho} \right)^2 = \frac{c^2(\rho)}{\rho^2} \quad \frac{\partial v}{\partial \rho} = \pm \frac{c}{\rho}$$

$$\Rightarrow \frac{\partial \rho}{\partial t} + (v \pm c) \frac{\partial \rho}{\partial x} = 0$$

Assuming adiabatic process : $c^2 = c_0^2 \left(\frac{\rho}{\rho_0} \right)^{\gamma-1}$ $c_0^2 = \frac{\gamma p_0}{\rho_0}$

$$\frac{\partial v}{\partial \rho} = \frac{dv}{d\rho} = \pm \frac{c}{\rho} \quad \Rightarrow v = \pm c_0 \int_{\rho_0}^{\rho} \left(\frac{\rho'}{\rho_0} \right)^{(\gamma-1)/2} \frac{d\rho'}{\rho'}$$

$$\Rightarrow v = \pm \frac{2c_0}{\gamma-1} \left(\left(\frac{\rho}{\rho_0} \right)^{(\gamma-1)/2} - 1 \right)$$

$$\Rightarrow c = c_0 \left(\frac{\rho}{\rho_0} \right)^{(\gamma-1)/2}$$

Summary :

$$\frac{dv}{d\rho} = \pm \frac{c}{\rho}$$

$$\frac{\partial \rho}{\partial t} + (v \pm c) \frac{\partial \rho}{\partial x} = 0$$

Assuming adiabatic process : $c^2 = c_0^2 \left(\frac{\rho}{\rho_0} \right)^{\gamma-1}$ $c_0^2 = \frac{\gamma p_0}{\rho_0}$

$$c = c_0 \left(\frac{\rho}{\rho_0} \right)^{(\gamma-1)/2} \quad v = \pm \frac{2c_0}{\gamma-1} \left(\left(\frac{\rho}{\rho_0} \right)^{(\gamma-1)/2} - 1 \right)$$

Traveling wave solution:

Assume: $\rho = \rho_0 + f(x - u(\rho)t)$

Need to find self-consistent equations for propagation velocity $u(\rho)$ using equations

From previous derivations: $\frac{\partial \rho}{\partial t} + (v \pm c) \frac{\partial \rho}{\partial x} = 0$

Apparently: $u(\rho) \Leftrightarrow v \pm c$

For adiabatic ideal gas and + signs:

$$u = v + c = c_0 \left(\frac{\gamma + 1}{\gamma - 1} \left(\frac{\rho}{\rho_0} \right)^{(\gamma-1)/2} - \frac{2}{\gamma - 1} \right)$$

Some details

Assume: $\rho = \rho_0 + f(x - u(\rho)t)$

Need to find self-consistent equations for
propagation velocity $u(\rho)$ using equations

From previous derivations: $\frac{\partial \rho}{\partial t} + (v \pm c) \frac{\partial \rho}{\partial x} = 0$

Apparently: $u(\rho) \Leftrightarrow v \pm c$

Note that for $u = v + c$ (choice of + solution)

$\frac{\partial \rho}{\partial t} + u \frac{\partial \rho}{\partial x} = 0$ is satisfied by a function of the form

$$\rho(x, t) = \rho_0 + f(x - u(\rho(x, t))t)$$

Let $w \equiv x - u(\rho(x, t))t$

$$\frac{df}{dw} \frac{\partial w}{\partial t} + u \frac{df}{dw} \frac{\partial w}{\partial x} = \frac{df}{dw} (-u + u) = 0$$

Traveling wave solution -- continued:

$$\frac{\partial \rho}{\partial t} + (v \pm c) \frac{\partial \rho}{\partial x} = 0$$

Assume: $\rho = \rho_0 + f(x - u(\rho)t) = \rho_0 + f(x - (v \pm c)t)$

For adiabatic ideal gas and + signs :

$$u = v + c = c_0 \left(\frac{\gamma + 1}{\gamma - 1} \left(\frac{\rho}{\rho_0} \right)^{(\gamma-1)/2} - \frac{2}{\gamma - 1} \right)$$

Solution in linear approximation:

$$u = v + c \approx v_0 + c_0 = c_0 \left(\frac{\gamma + 1}{\gamma - 1} - \frac{2}{\gamma - 1} \right) = c_0$$

$$\Rightarrow \rho = \rho_0 + f(x - c_0 t)$$

Traveling wave solution -- full non-linear case:

Visualization for particular waveform: $\rho = \rho_0 + f(\underbrace{x - u(\rho)t}_w)$

Assume: $f(w) \equiv \rho_0 s(w)$

$$\frac{\rho}{\rho_0} = 1 + s(x - ut)$$

For adiabatic ideal gas:

$$u = c_0 \left(\frac{\gamma + 1}{\gamma - 1} \left(\frac{\rho}{\rho_0} \right)^{(\gamma-1)/2} - \frac{2}{\gamma - 1} \right)$$

$$u = c_0 \left(\frac{\gamma + 1}{\gamma - 1} \left(1 + s(x - ut) \right)^{(\gamma-1)/2} - \frac{2}{\gamma - 1} \right)$$

Visualization continued:

$$u = c_0 \left(\frac{\gamma + 1}{\gamma - 1} (1 + s(x - ut))^{(\gamma-1)/2} - \frac{2}{\gamma - 1} \right)$$

Plot $s(x - ut)$ for fixed t , as a function of x :

Let $w = x - ut$

$$x = w + ut = w + u(w)t \equiv x(w, t)$$

$$u(w) = c_0 \left(\frac{\gamma + 1}{\gamma - 1} (1 + s(w))^{(\gamma-1)/2} - \frac{2}{\gamma - 1} \right)$$

Parametric equations:

plot $s(w)$ vs $x(w, t)$ for range of w at each t

Summary

$$\frac{\partial \rho}{\partial t} + u(\rho) \frac{\partial \rho}{\partial x} = 0$$

Solution: $\rho = \rho_0 + f(x - u(\rho)t) = \rho_0 (1 + s(x - u(\rho)t))$

For linear case: $u(\rho) = c_0$

For non-linear case: $u(\rho) = c_0 \left(\frac{\gamma + 1}{\gamma - 1} (1 + s(x - ut))^{(\gamma-1)/2} - \frac{2}{\gamma - 1} \right)$


Plot $s(x - ut)$ for fixed t , as a function of x :

Let $w = x - ut \Rightarrow x = w + ut = w + u(w)t \equiv x(w, t)$

$$u(w) = c_0 \left(\frac{\gamma + 1}{\gamma - 1} (1 + s(w))^{(\gamma-1)/2} - \frac{2}{\gamma - 1} \right)$$

Parametric equations: plot $s(w)$ vs $x(w, t)$ for range of w


Linear wave:



Non-linear wave:



Linear wave




Non-linear wave




Analysis of shock wave

Plots of $\delta\rho$

Solution becomes
unphysical



Analysis of shock wave -- continued



Note that in this case u is assumed to be a given parameter of the system.

Analysis of shock wave – continued

While analysis in the shock region is complicated, we can use conservation laws to analyze regions 1 and 2

Assume $\rho(x,t) = \rho(x - ut)$

$$p(x,t) = p(x - ut)$$

$$v(x,t) = v(x - ut)$$


Continuity equation:

$$\frac{\partial \rho}{\partial t} + \frac{\partial(\rho v)}{\partial x} = 0 = \frac{\partial(\rho v - \rho u)}{\partial x} \quad \Rightarrow (v_2 - u)\rho_2 = (v_1 - u)\rho_1$$

Conservation of energy and momentum:

$$\Rightarrow p_2 + \rho_2 (v_2 - u)^2 = p_1 + \rho_1 (v_1 - u)^2$$

$$\Rightarrow \epsilon_2 + \frac{1}{2}(v_2 - u)^2 + \frac{p_2}{\rho_2} = \epsilon_1 + \frac{1}{2}(v_1 - u)^2 + \frac{p_1}{\rho_1}$$



Analysis of shock wave – continued


While analysis in the shock region is complicated, we can use conservation laws to analyze regions 1 and 2

Summary of equations

$$\Rightarrow (v_2 - u)\rho_2 = (v_1 - u)\rho_1$$

$$\Rightarrow p_2 + \rho_2 (v_2 - u)^2 = p_1 + \rho_1 (v_1 - u)^2$$

$$\Rightarrow \epsilon_2 + \frac{1}{2}(v_2 - u)^2 + \frac{p_2}{\rho_2} = \epsilon_1 + \frac{1}{2}(v_1 - u)^2 + \frac{p_1}{\rho_1}$$



Assume that within each regions (1 & 2), the ideal gas equations apply

$$\epsilon_1 + \frac{p_1}{\rho_1} = \frac{\gamma}{\gamma - 1} \frac{p_1}{\rho_1} \quad \epsilon_2 + \frac{p_2}{\rho_2} = \frac{\gamma}{\gamma - 1} \frac{p_2}{\rho_2}$$



It follows that

$$\frac{\gamma}{\gamma - 1} \frac{p_2}{\rho_2} + \frac{1}{2}(v_2 - u)^2 = \frac{\gamma}{\gamma - 1} \frac{p_1}{\rho_1} + \frac{1}{2}(v_1 - u)^2$$

Analysis of shock wave – continued

For adiabatic ideal gas, also considering energy and momentum conservation:

$$\frac{\rho_2}{\rho_1} = \frac{\frac{\gamma+1}{\gamma-1} \frac{p_2}{p_1} + 1}{\frac{\gamma+1}{\gamma-1} + \frac{p_2}{p_1}} \leq \frac{\gamma+1}{\gamma-1}$$



Analysis of shock wave – continued

For adiabatic ideal gas, entropy considerations::

Internal energy density: $\varepsilon = \frac{p}{(\gamma - 1)\rho} = C_V T$

First law of thermo: $d\varepsilon = Tds - pd\left(\frac{1}{\rho}\right)$

$$ds = \frac{1}{T} \left(d \left(\frac{p}{(\gamma - 1)\rho} \right) - pd \left(\frac{1}{\rho} \right) \right) = C_V d \ln \left(\frac{p}{\rho^\gamma} \right)$$

$$s = C_V \ln \left(\frac{p}{\rho^\gamma} \right) + (\text{constant})$$

$$s_2 - s_1 = C_V \ln \left(\frac{p_2}{p_1} \left(\frac{\rho_1}{\rho_2} \right)^\gamma \right) \quad 0 < s_2 - s_1 < C_V \left(\ln \left(\frac{p_2}{p_1} \right) - \gamma \ln \left(\frac{\gamma + 1}{\gamma - 1} \right) \right)$$