




# **PHY 711 Classical Mechanics and Mathematical Methods**

## **10-10:50 AM MWF in Olin 103**

### **Notes for Lecture 24: Chap. 7 & App. A-D (F&W)**

**Generalization of the one dimensional wave equation →  
various mathematical problems and techniques including:**

- 
- 1. Fourier transforms**
  - 2. Laplace transforms (very briefly)**
  - 3. Complex variables**
  - 4. Contour integrals**



19	Mon, 10/03/2022	Chap. 7	Sturm-Liouville equations	#15	10/05/2022
20	Wed, 10/05/2022	Chap. 7	Sturm-Liouville equations		
21	Fri, 10/07/2022	Chap. 1-4,6-7	Review		
	Mon, 10/10/2022	No class	Take home exam		
	Wed, 10/12/2022	No class	Take home exam		
	Fri, 10/14/2022	No class	Fall break		
22	Mon, 10/17/2022	Chap. 7	Green's function methods for one-dimensional Sturm-Liouville equations	#16	10/19/2022
23	Wed, 10/19/2022	Chap. 7	Fourier and other transform methods	#17	10/21/2022
24	Fri, 10/21/2022	Chap. 7	Complex variables and contour integration	#18	10/24/2022
25	Mon, 10/24/2022	Chap. 5	Rigid body motion		

## PHY 711 – Homework # 18

Read Appendix A of **Fetter and Walecka**.

1. Assume that  $a > 0$  and  $b > 0$ ; use contour integration methods to evaluate the integral:

$$\int_{-\infty}^{\infty} \frac{e^{iax}}{x^2 + b^2} dx.$$

Note that you may use Maple, Mathematica, or other software to evaluate this integral, but full credit will be earned by using the contour integration methods.

## Your questions

From Athul -- In slide 15, is it an example of analytic or non analytic functions? I might be missing something but I don't understand how different functions produce almost similar solutions in later slides.

## A brief introduction to Laplace transforms --

Laplace transforms are particularly useful in solving initial value problems.  
The Laplace transform of the function  $\phi(x)$  is defined:

$$L_{\phi(x)}(p) \equiv \int_0^{\infty} e^{-px} \phi(x) dx$$

Assuming that  $\phi(x)$  is well-behaved in the interval  $0 \leq x \leq \infty$ , the following identities can be shown:

$$L_{d\phi(x)/dx}(p) \equiv \int_0^{\infty} e^{-px} \frac{d\phi(x)}{dx} dx = -\phi(0) + pL_{\phi(x)}(p)$$

and

$$L_{d^2\phi(x)/dx^2}(p) \equiv \int_0^{\infty} e^{-px} \frac{d^2\phi(x)}{dx^2} dx = -\frac{d\phi(0)}{dx} - p\phi(0) + p^2 L_{\phi(x)}(p)$$

These identities allow us to turn a differential equation for  $\phi(x)$  into an algebraic equation for  $\mathcal{L}_\phi(p)$ . We then need to perform an inverse Laplace transform to find  $\phi(x)$ . For illustration, we will consider a simple example with  $\tau(x) = 1$ ,  $\sigma(x) = 1$ ,  $\lambda = 0$ . The differential equation then becomes

$$-\frac{d^2\phi(x)}{dx^2} = F(x), \quad (27)$$

where we will take the initial conditions to be  $\phi(0) = 0$  and  $d\phi(0)/dx = 0$ . For our example, we will also take  $F(x) = F_0 e^{-\gamma x}$ . Multiplying, both sides of the equation by  $e^{-px}$  and integrating  $0 \leq x \leq \infty$ , we find

$$\mathcal{L}_\phi(p) = -\frac{F_0}{p^2(\gamma + p)}. \quad (28)$$

In general the inverse Laplace transform involves performing a contour integral, but we can use the following simple relations

$$\mathcal{L}_1 = \int_0^\infty e^{-px} dx = \frac{1}{p}. \quad (29)$$

$$\mathcal{L}_x = \int_0^\infty x e^{-px} dx = \frac{1}{p^2}. \quad (30)$$

$$\mathcal{L}_{e^{-\gamma x}} = \int_0^\infty e^{-\gamma x} e^{-px} dx = \frac{1}{p + \gamma}. \quad (31)$$

Noting that

$$-\frac{F_0}{p^2(\gamma + p)} = -\frac{F_0}{\gamma^2} \left( \frac{1}{\gamma + p} - \frac{1}{p} + \frac{\gamma}{p^2} \right), \quad (32)$$

we see that the inverse Laplace transform gives us

$$\phi(x) = \frac{F_0}{\gamma^2} \left( 1 - e^{-\gamma x} - \gamma x \right). \quad (33)$$

We can check that this a solution to the differential equation

$$-\frac{d^2\phi}{dx^2} = F_0 e^{-\gamma x} \quad \text{for } \phi(0) = 0 \quad \text{and} \quad \frac{d\phi}{dx}(0) = 0$$

Using Laplace transforms to solve equation :

$$\left( -\frac{d^2}{dx^2} - 1 \right) \phi(x) = F_0 \sin\left(\frac{\pi x}{L}\right) \quad \text{with } \phi(0) = 0, \frac{d\phi(0)}{dx} = 0$$

$$\begin{aligned} \mathcal{L}_\phi(p) &= -\left(\frac{\pi}{L}\right) \frac{F_0}{\left(p^2 + 1\right)\left(p^2 + \left(\frac{\pi}{L}\right)^2\right)} \\ &= -F_0 \left(\frac{\pi/L}{(\pi/L)^2 - 1}\right) \left( \frac{1}{p^2 + 1} - \frac{1}{p^2 + \left(\frac{\pi}{L}\right)^2} \right) \end{aligned}$$

Note that :  $\int_0^\infty \sin(at)e^{-pt} dt = \frac{a}{a^2 + p^2}$

$$\Rightarrow \phi(x) = \frac{F_0}{(\pi/L)^2 - 1} \left( \sin\left(\frac{\pi x}{L}\right) - \frac{\pi}{L} \sin(x) \right)$$

# Table of Laplace transforms

## Laplace Transform Table

Largely modeled on a table in D'Azzo and Houpis, *Linear Control Systems Analysis and Design*, 1988

$F(s)$	$f(t) \quad 0 \leq t$
1. $1$	$\delta(t)$ unit impulse at $t = 0$
2. $\frac{1}{s}$	$1$ or $u(t)$ unit step starting at $t = 0$
3. $\frac{1}{s^2}$	$t \cdot u(t)$ or $t$ ramp function
4. $\frac{1}{s^n}$	$\frac{1}{(n-1)!} t^{n-1}$ $n = \text{positive integer}$
5. $\frac{1}{s} e^{-as}$	$u(t-a)$ unit step starting at $t = a$
6. $\frac{1}{s} (1 - e^{-as})$	$u(t) - u(t-a)$ rectangular pulse
7. $\frac{1}{s+a}$	$e^{-at}$ exponential decay
8. $\frac{1}{(s+a)^n}$	$\frac{1}{(n-1)!} t^{n-1} e^{-at}$ $n = \text{positive integer}$
9. $\frac{1}{s(s+a)}$	$\frac{1}{a} (1 - e^{-at})$
10. $\frac{1}{s(s+a)(s+b)}$	$\frac{1}{ab} (1 - \frac{b}{b-a} e^{-at} + \frac{a}{b-a} e^{-bt})$
11. $\frac{s+\alpha}{s(s+a)(s+b)}$	$\frac{1}{ab} [\alpha - \frac{b(\alpha-a)}{b-a} e^{-at} + \frac{a(\alpha-b)}{b-a} e^{-bt}]$
12. $\frac{1}{(s+a)(s+b)}$	$\frac{1}{b-a} (e^{-at} - e^{-bt})$
13. $\frac{s}{(s+a)(s+b)}$	$\frac{1}{a-b} (ae^{-at} - be^{-bt})$

<https://www.dartmouth.edu/~sullivan/22files/New%20Laplace%20Transform%20Table.pdf>

Inverse Laplace transform :

$$\mathcal{L}_\phi(p) = \int_0^\infty e^{-pt} \phi(t) dt$$

$$\phi(t) = \frac{1}{2\pi i} \int_{\lambda-i\infty}^{\lambda+i\infty} e^{pt} \mathcal{L}_\phi(p) dp$$

Check:

$$\begin{aligned} \frac{1}{2\pi i} \int_{\lambda-i\infty}^{\lambda+i\infty} e^{pt} \mathcal{L}_\phi(p) dp &= \frac{1}{2\pi i} \int_{\lambda-i\infty}^{\lambda+i\infty} e^{pt} dp \int_0^\infty e^{-pu} \phi(u) du \\ \frac{1}{2\pi i} \int_0^\infty \phi(u) du \int_{\lambda-i\infty}^{\lambda+i\infty} e^{p(t-u)} dp &= \frac{1}{2\pi i} \int_0^\infty \phi(u) du \int_{-\infty}^\infty e^{\lambda(t-u)} e^{is(t-u)} i ds \\ &= \frac{1}{2\pi i} \int_0^\infty \phi(u) du \left( e^{\lambda(t-u)} 2\pi i \delta(t-u) \right) \\ &= \begin{cases} \phi(t) & \text{if } t \geq 0 \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

In order to evaluate these integrals, we need to use complex analysis.



In general – to calculate inverse Laplace transforms, we need to introduce concepts of complex numbers and contour integration

## Introduction to complex variables

1. Basic properties
2. Notion of an analytic complex function
3. Cauchy integral theory
4. Analytic functions and functions with poles
5. Evaluating integrals of functions in the complex plane

## Complex numbers

$$i \equiv \sqrt{-1} \quad i^2 = -1$$

Define  $z = x + iy$

$$|z|^2 = zz^* = (x + iy)(x - iy) = x^2 + y^2$$

Polar representation

$$z = \rho(\cos\phi + i\sin\phi) = \rho e^{i\phi}$$

## Functions of complex variables

$$f(z) = \Re(f(z)) + i\Im(f(z)) \equiv u(x, y) + iv(x, y)$$

## Derivatives: Cauchy-Riemann equations

$$\frac{\partial f(z)}{\partial x} = \frac{\partial u(z)}{\partial x} + i \frac{\partial v(z)}{\partial x} \quad \frac{\partial f(z)}{\partial \bar{y}} = \frac{\partial u(z)}{\partial \bar{y}} + i \frac{\partial v(z)}{\partial \bar{y}} = \frac{\partial v(z)}{\partial y} - i \frac{\partial u(z)}{\partial y}$$

Argue that  $\frac{df}{dz} = \frac{\partial f(z)}{\partial x} = \frac{\partial f(z)}{i\partial y} \Rightarrow \frac{\partial u(z)}{\partial x} = \frac{\partial v(z)}{\partial y}$  and  $\frac{\partial v(z)}{\partial x} = -\frac{\partial u(z)}{\partial y}$



# Analytic function

$f(z)$  is analytic if it is:

- continuous
- single valued
- its first derivative satisfies Cauchy-Rieman conditions

## Examples of analytic functions

$$e^z = e^{x+iy} = e^x \cos(y) + ie^x \sin(y)$$

$$\frac{\partial u}{\partial x} = e^x \cos(y) = \frac{\partial v}{\partial y} \quad \frac{\partial v}{\partial x} = e^x \sin(y) = -\frac{\partial u}{\partial y}$$

$$z^2 = (x + iy)^2 = (x^2 - y^2) + 2ixy \equiv u(x, y) + iv(x, y)$$

$$\frac{\partial u}{\partial x} = 2x = \frac{\partial v}{\partial y} \quad \frac{\partial v}{\partial x} = 2y = -\frac{\partial u}{\partial y}$$



## Examples of non-analytic functions

Note that  $z = \rho e^{i\phi} = \rho e^{i\phi+i2\pi n}$  for any integer  $n$

$$\Rightarrow \ln z = \ln \rho + i(\phi + 2\pi n)$$

$\ln z$  is not analytic because it is multivalued

$$\Rightarrow z^\alpha = \rho^\alpha e^{i\alpha\phi} e^{i2\pi n\alpha}$$

$z^\alpha$  is not analytic for non-integer  $\alpha$   
because it is multivalued

Behavior of  $f(z) = \frac{1}{z^n}$  about the point  $z = 0$ :

For an integer  $n$ , consider

$$\oint \frac{1}{z^n} dz = \int_0^{2\pi} \frac{\rho e^{i\phi} id\phi}{\rho^n e^{in\phi}} = \rho^{1-n} \int_0^{2\pi} e^{i(1-n)\phi} id\phi = \begin{cases} 0 & n \neq 1 \\ 2\pi i & n = 1 \end{cases}$$

Behavior of  $f(z) = \frac{1}{z^n}$  about the point  $z = 0$ :

For an integer  $n$ , consider


$$\oint \frac{1}{z^n} dz = \int_0^{2\pi} \frac{\rho e^{i\phi} i d\phi}{\rho^n e^{in\phi}} = \rho^{1-n} \int_0^{2\pi} e^{i(1-n)\phi} i d\phi = \begin{cases} 0 & n \neq 1 \\ 2\pi i & n = 1 \end{cases}$$

This observation helps us to focus on a special kind  
of singularity called a "pole"

For  $f(z)$  in the vicinity of  $z = z_p$ :  $f(z) \approx \frac{g(z_p)}{z - z_p}$

Therefore:  $\oint f(z) dz = 0$       or       $\oint f(z) dz = g(z_p) \oint \frac{dz}{z - z_p} = 2\pi i g(z_p)$

Integration does  
not include  $z_p$       Integration does  
include  $z_p$





General formula for determining residue:

Suppose that in the neighborhood of  $z_p$ ,  $f(z) \approx \frac{h(z)}{(z - z_p)^m} \underset{z \rightarrow z_p}{\equiv} \frac{\text{Res}(f(z_p))}{z - z_p}$

Since  $h(z) \equiv (z - z_p)^m f(z)$  is analytic near  $z_p$ , we can make a Taylor expansion


about  $z_p$ :  $h(z) \approx h(z_p) + (z - z_p) \frac{dh(z_p)}{dz} + \dots + \frac{(z - z_p)^{m-1}}{(m-1)!} \frac{d^{m-1}h(z_p)}{dz^{m-1}} + \dots$

$$\Rightarrow \text{Res}(f(z_p)) = \lim_{z \rightarrow z_p} \left\{ \frac{1}{(m-1)!} \frac{d^{m-1} \left( (z - z_p)^m f(z) \right)}{dz^{m-1}} \right\}$$

In the following examples  $m=1$

Example:

$$\int_{-\infty}^{\infty} \frac{x^2}{1+x^4} dx = \int_{-\infty}^{\infty} \frac{x^2}{1+x^4} dx + 0 = \oint \frac{z^2}{1+z^4} dz$$



$$1+z^4 = (z - e^{i\pi/4})(z - e^{3i\pi/4})(z - e^{-i\pi/4})(z - e^{-3i\pi/4})$$

$$\oint \frac{z^2}{1+z^4} dz = 2\pi i \left( \text{Res}(z_p = e^{i\pi/4}) + \text{Res}(z_p = e^{3i\pi/4}) \right)$$

Note:  
 $m=1$

$$\text{Res}(z_p = e^{i\pi/4}) = \frac{e^{i\pi/4}}{4i} \quad \text{Res}(z_p = e^{3i\pi/4}) = -\frac{e^{3i\pi/4}}{4i}$$

$$\oint \frac{z^2}{1+z^4} dz = 2\pi i \left( \frac{e^{i\pi/4}}{4i} - \frac{e^{3i\pi/4}}{4i} \right) = \frac{\pi}{2} \left( \left( \sqrt{\frac{1}{2}} + i\sqrt{\frac{1}{2}} \right) - \left( -\sqrt{\frac{1}{2}} + i\sqrt{\frac{1}{2}} \right) \right) = \frac{\pi}{\sqrt{2}}$$



Some details:

$$f(z) = \frac{z^2}{1+z^4}$$

Note that:  $e^{i\pi} = -1 = e^{-i\pi}$

$$\begin{aligned}\text{Res}\left(f(z = e^{i\pi/4})\right) &= \frac{\left(e^{i\pi/4}\right)^2}{\left(e^{i\pi/4} - e^{3i\pi/4}\right)\left(e^{i\pi/4} - e^{-i\pi/4}\right)\left(e^{i\pi/4} - e^{-3i\pi/4}\right)} \\ &= \frac{e^{i\pi/2}}{\left(e^{i\pi/4} + e^{-i\pi/4}\right)\left(e^{i\pi/4} - e^{-i\pi/4}\right)\left(e^{i\pi/4} + e^{i\pi/4}\right)} \\ &= \frac{e^{i\pi/4}}{2(i - (-i))} = \frac{e^{i\pi/4}}{4i}\end{aligned}$$

Question – Could we have chosen the contour in the lower half plane?


- a. Yes
- b. No

Another example:  $I = \int_0^\infty \frac{\cos(ax)}{4x^4 + 5x^2 + 1} dx$

$$\int_0^\infty \frac{\cos(ax)}{4x^4 + 5x^2 + 1} dx = \frac{1}{2} \int_{-\infty}^\infty \frac{e^{iax}}{4x^4 + 5x^2 + 1} dx = \frac{1}{2} \oint \frac{e^{iaz}}{4z^4 + 5z^2 + 1} dz$$

$$4z^4 + 5z^2 + 1 = 4(z - i)(z - \frac{i}{2})(z + i)(z + \frac{i}{2})$$

Note:  
 $m=1$



$$I = 2\pi i \left( \operatorname{Res}(z_p = i) + \operatorname{Res}(z_p = \frac{i}{2}) \right)$$

$$\begin{aligned}
\int_0^\infty \frac{\cos(ax)}{4x^4 + 5x^2 + 1} dx &= \frac{1}{2} \oint \frac{e^{iaz}}{4z^4 + 5z^2 + 1} dz \\
&= 2\pi i \left( \text{Res}(z_p = i) + \text{Res}(z_p = \frac{i}{2}) \right) \\
&= \frac{\pi}{6} \left( -e^{-a} + 2e^{-a/2} \right)
\end{aligned}$$

Question – Could we have chosen the contour in the lower half plane?

- a. Yes
- b. No


Note that for  $a > 0$  and  $z_I > 0$

in the lower half plane:  $e^{iaz} = e^{iaz_R} e^{az_I}$

Another example:  $I = \int_{-\infty}^{\infty} \frac{x \sin kx}{x^2 + a^2} dx$  for  $k > 0$  and  $a > 0$

$$\int_{-\infty}^{\infty} \frac{x \sin kx}{x^2 + a^2} dx = \frac{1}{i} \int_{-\infty}^{\infty} \frac{xe^{ikx}}{x^2 + a^2} dx = \frac{1}{i} \oint \frac{ze^{ikz}}{z^2 + a^2} dz$$

$$z^2 + a^2 = (z - ia)(z + ia)$$



$$I = 2\pi i \left( \text{Res} \left( z_p = ia \right) \right) = \pi e^{-ka}$$

# Some details --

$$\int_{-\infty}^{\infty} \frac{x \sin kx}{x^2 + a^2} dx = \frac{1}{i} \int_{-\infty}^{\infty} \frac{xe^{ikx}}{x^2 + a^2} dx = \frac{1}{i} \oint \frac{ze^{ikz}}{z^2 + a^2} dz$$

$$z^2 + a^2 = (z - ia)(z + ia)$$


$$\frac{1}{i} \oint \frac{ze^{ikz}}{z^2 + a^2} dz = 2\pi i \lim_{z \rightarrow ia} \left( (z - ia) \frac{ze^{ikz}}{z^2 + a^2} \right)$$

$$= 2\pi i \frac{1}{i} \frac{iae^{-ka}}{2ia} = \pi e^{-ka}$$

From the Drude model of dielectric response --

$$G(\tau) = \frac{\omega_p^2}{2\pi} \int_{-\infty}^{\infty} d\omega \frac{e^{-i\omega\tau}}{\omega_0^2 - \omega^2 - i\gamma\omega} \quad \text{where } \omega_p, \omega_0, \text{ and } \gamma \text{ are positive constants}$$

Upper hemisphere:



$$z = x + iy \quad e^{-i\omega\tau} = e^{-ix\tau + y\tau} \\ \Rightarrow \text{Converges for } \tau < 0$$

Lower hemisphere:

$$v_0 \equiv \sqrt{\omega_0^2 - \frac{\gamma^2}{4}}$$

$$z = x - iy \quad e^{-i\omega\tau} = e^{-ix\tau - y\tau} \\ \Rightarrow \text{Converges for } \tau > 0$$



From the Drude model of dielectric response -- continued --

$$G(\tau) = \frac{\omega_p^2}{2\pi} \int_{-\infty}^{\infty} d\omega \frac{e^{-i\omega\tau}}{\omega_0^2 - \omega^2 - i\gamma\omega} \quad \text{where } \omega_p, \omega_0, \text{ and } \gamma \text{ are positive constants}$$

$$G(\tau) = \omega_p^2 \begin{cases} 0 & \text{for } \tau < 0 \\ e^{-\gamma\tau/2} \frac{\sin \nu_0 \tau}{\nu_0} & \text{for } \tau > 0 \end{cases}$$

Cauchy integral theorem for analytic function  $f(z)$ :


$$f(z) = \frac{1}{2\pi i} \oint_C \frac{f(z')}{z' - z} dz'.$$



## Example

Suppose  $f(|z| \rightarrow \infty) = 0$  and for  $z = x$ :


$$f(x) = a(x) + ib(x)$$





## Example -- continued


$$f(z) = \frac{1}{2\pi i} \oint \frac{f(z')}{z' - z} dz' \quad \text{where} \quad f(x) = a(x) + ib(x)$$




$$a(x) + ib(x) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{a(x') + ib(x')}{x' - x} dx' + 0$$



## Example -- continued



$$\begin{aligned} \int_{-\infty}^{\infty} \frac{f(x')}{x' - x} dx' &= \int_{-\infty}^{x-\varepsilon} \frac{f(x')}{x' - x} dx' + \int_{x+\varepsilon}^{\infty} \frac{f(x')}{x' - x} dx' + \int_{x-\varepsilon}^{x+\varepsilon} \frac{f(x')}{x' - x} dx' \\ &= P \int_{-\infty}^{\infty} \frac{f(x')}{x' - x} dx' + i\pi f(x) \end{aligned}$$



let  $u = x' - x$

let  $x \rightarrow x + i\eta$


$$\int_{x-\varepsilon}^{x+\varepsilon} \frac{f(x')}{x'-x} dx' \approx f(x) \lim_{\eta \rightarrow 0} \int_{-\varepsilon}^{\varepsilon} \frac{1}{u - i\eta} du = f(x) \lim_{\eta \rightarrow 0} \int_{-\varepsilon}^{\varepsilon} \frac{u + i\eta}{u^2 + \eta^2} du$$

$$= i\pi f(x) \quad \text{since} \quad \lim_{\eta \rightarrow 0} \frac{i\eta}{u^2 + \eta^2} \approx i\pi \delta(u)$$



More details --

$$\lim_{\eta \rightarrow 0} \frac{\eta}{u^2 + \eta^2} \approx \pi \delta(u)$$





## Example -- continued

$$\int_{-\infty}^{\infty} \frac{f(x')}{x'-x} dx' = \int_{-\infty}^{x-\varepsilon} \frac{f(x')}{x'-x} dx' + \int_{x+\varepsilon}^{\infty} \frac{f(x')}{x'-x} dx' + \int_{x-\varepsilon}^{x+\varepsilon} \frac{f(x')}{x'-x} dx'$$
$$= P \int_{-\infty}^{\infty} \frac{f(x')}{x'-x} dx' + i\pi f(x)$$

$$a(x) + ib(x) = \frac{P}{2\pi i} \int_{-\infty}^{\infty} \frac{a(x') + ib(x')}{x'-x} dx' + \frac{\pi i}{2\pi i} (a(x) + ib(x))$$


$$\Rightarrow a(x) = \frac{P}{\pi} \int_{-\infty}^{\infty} \frac{b(x')}{x'-x} dx' \quad b(x) = -\frac{P}{\pi} \int_{-\infty}^{\infty} \frac{a(x')}{x'-x} dx'$$

## Kramers-Kronig relationships



## Comment on evaluating principal parts integrals


$$a(x) = \frac{P}{\pi} \int_{-\infty}^{\infty} \frac{b(x')}{x' - x} dx' = \lim_{\epsilon \rightarrow 0} \left( \frac{1}{\pi} \int_{-\infty}^{x-\epsilon} \frac{b(x')}{x' - x} dx' + \frac{1}{\pi} \int_{x+\epsilon}^{\infty} \frac{b(x')}{x' - x} dx' \right)$$





Example:

$$b(x') = \begin{cases} 0 & \text{for } x' < -2L, \quad -L < x' < L, \quad x' > 2L \\ B_0 & \text{for } L < x' < 2L \\ -B_0 & \text{for } -2L < x' < -L \end{cases}$$



Example:

$$b(x') = \begin{cases} 0 & \text{for } x' < -2L, \quad -L < x' < L, \quad x' > 2L \\ B_0 & \text{for } L < x' < 2L \\ -B_0 & \text{for } -2L < x' < -L \end{cases}$$

$$a(x) = \frac{P}{\pi} \int_{-\infty}^{\infty} \frac{b(x')}{x' - x} dx' = \lim_{\epsilon \rightarrow 0} \left( \frac{1}{\pi} \int_{-\infty}^{x-\epsilon} \frac{b(x')}{x' - x} dx' + \frac{1}{\pi} \int_{x+\epsilon}^{\infty} \frac{b(x')}{x' - x} dx' \right)$$

For  $x < -2L$  or  $x > 2L$   $-L < x < L$ :


$$a(x) = \frac{-B_0}{\pi} \int_{-2L}^{-L} \frac{dx'}{x' - x} + \frac{B_0}{\pi} \int_L^{2L} \frac{dx'}{x' - x}$$

$$= \frac{-B_0}{\pi} \ln \left( \left| \frac{x+L}{x+2L} \right| \right) + \frac{B_0}{\pi} \ln \left( \left| \frac{x-2L}{x-L} \right| \right) = \frac{B_0}{\pi} \ln \left( \left| \frac{x^2 - 4L^2}{x^2 - L^2} \right| \right)$$

$$a(x) = \frac{P}{\pi} \int_{-\infty}^{\infty} \frac{b(x')}{x' - x} dx' = \lim_{\epsilon \rightarrow 0} \left( \frac{1}{\pi} \int_{-\infty}^{x-\epsilon} \frac{b(x')}{x' - x} dx' + \frac{1}{\pi} \int_{x+\epsilon}^{\infty} \frac{b(x')}{x' - x} dx' \right)$$

For our example:

$$a(x) = \frac{B_0}{\pi} \ln \left( \left| \frac{4L^2 - x^2}{L^2 - x^2} \right| \right)$$





# Summary

For a function  $f(x)$ , analytic along the real line:

$$f(x) = \Re(f(x)) + i\Im(f(x)) = a(x) + ib(x)$$

$$\Rightarrow a(x) = \frac{P}{\pi} \int_{-\infty}^{\infty} \frac{b(x')}{x' - x} dx', \quad b(x) = -\frac{P}{\pi} \int_{-\infty}^{\infty} \frac{a(x')}{x' - x} dx'$$

Example:


$$f(x) = \frac{1}{x+i} \quad a(x) = \frac{x}{x^2+1} \quad b(x) = -\frac{1}{x^2+1}$$

Check:

$$\frac{P}{\pi} \int_{-\infty}^{\infty} \frac{b(x')}{x' - x} dx' = -\frac{P}{\pi} \int_{-\infty}^{\infty} \frac{1}{(x' - x)(x'^2 + 1)} dx' \stackrel{?}{=} \frac{x}{x^2 + 1} = a(x)$$

$$a(\omega) = \frac{\omega - 10}{(\omega - 10)^2 + 1}$$

$$b(\omega) = \frac{1}{(\omega - 10)^2 + 1}$$



# Continued:

$$\begin{aligned}
 \frac{P}{\pi} \int_{-\infty}^{\infty} \frac{b(x')}{x' - x} dx' &= -\frac{P}{\pi} \int_{-\infty}^{\infty} \frac{1}{(x' - x)(x'^2 + 1)} dx' \\
 &= -\frac{P}{\pi} \int_{-\infty}^{\infty} \left( \frac{1}{(x' - x)(x'^2 + 1)} - \frac{1}{(x' - x)(x^2 + 1)} \right) dx' - \frac{1}{(x^2 + 1)} \frac{P}{\pi} \int_{-\infty}^{\infty} \frac{1}{x' - x} dx' \\
 &= -\frac{P}{\pi} \int_{-\infty}^{\infty} \left( \frac{x^2 - x'^2}{(x' - x)(x'^2 + 1)(x^2 + 1)} \right) dx' - \frac{1}{(x^2 + 1)} \frac{P}{\pi} \int_{-\infty}^{\infty} \frac{1}{x' - x} dx' \\
 &= \frac{P}{\pi} \int_{-\infty}^{\infty} \left( \frac{x + x'}{(x'^2 + 1)(x^2 + 1)} \right) dx' - \frac{1}{(x^2 + 1)} \frac{P}{\pi} \int_{-\infty}^{\infty} \frac{1}{x' - x} dx'
 \end{aligned}$$

Note that:  $\int_{x+\epsilon}^X \frac{1}{x' - x} dx' = \ln(X - x) - \ln(\epsilon) = \ln\left(\frac{X - x}{\epsilon}\right)$

$$\int_{-X}^{x-\epsilon} \frac{1}{x' - x} dx' = -\ln(-X - x) + \ln(-\epsilon) = -\ln\left(\frac{X + x}{\epsilon}\right)$$

$$\frac{P}{\pi} \int_{-\infty}^{\infty} \frac{1}{x' - x} dx' = \lim_{X \rightarrow \infty} \ln\left(\frac{X - x}{X + x}\right) = 0 \quad \frac{P}{\pi} \int_{-\infty}^{\infty} \frac{1}{x'^2 + 1} dx' = 1$$

$$\frac{P}{\pi} \int_{-\infty}^{\infty} \frac{b(x')}{x' - x} dx' = \frac{x}{x^2 + 1} = a(x)$$