



PHY 711 Classical Mechanics and Mathematical Methods

10-10:50 AM MWF in Olin 103

Notes on Lecture 33:

Chapter 10 in F & W: Surface waves

1. Water waves in a channel

2. Wave-like solutions; wave speed



28	Mon, 10/31/2022	Chap. 9	Mechanics of 3 dimensional fluids	#21	11/02/2022
29	Wed, 11/02/2022	Chap. 9	Mechanics of 3 dimensional fluids	#22	11/04/2022
30	Fri, 11/04/2022	Chap. 9	Linearized hydrodynamics equations	#23	11/07/2022
31	Mon, 11/07/2022	Chap. 9	Linear sound waves	#24	11/09/2022
32	Wed, 11/09/2022	Chap. 9	Scattering of sound and non-linear effects	#25	11/11/2022
33	Fri, 11/11/2022	Chap. 10	Surface waves in fluids	#26	11/16/2022
34	Mon, 11/14/2022	Chap. 10	Surface waves in fluids; soliton solutions		
35	Wed, 11/16/2022	Chap. 11	Heat conduction		
36	Fri, 11/18/2022	Chap. 12	Viscous effects on hydrodynamics		
37	Mon, 11/21/2022	Chap 1-12	Review		
	Wed, 11/23/2022		Thanksgiving Holiday		
	Fri, 11/25/2022		Thanksgiving Holiday		
	Mon, 11/28/2022		Presentations I		
	Wed, 11/30/2022		Presentations II		
	Fri, 12/02/2022		Presentations III		



PHY 711 -- Assignment #26

Nov. 11, 2022

Start reading Chapter 10 in **Fetter & Walecka**.

1. Work Problem 10.3 at the end of Chapter 10 in **Fetter and Walecka**.

Note that some of the ideas are discussed in today's lecture.

Reference: Chapter 10 of Fetter and Walecka

Physics of incompressible fluids and their surfaces






Consider a container of water with average height h and surface $h + \zeta(x, y, t)$; ($h \leftrightarrow z_0$ on some of the slides)

Atmospheric pressure is in equilibrium with the surface of water

Pressure at a height z above the bottom where the surface is at a height $h + \zeta$:

$$p(z) = \begin{cases} p_0 + \rho g (h + \zeta - z) & \text{For } z \leq h + \zeta \\ p_0 & \text{For } z > h + \zeta \end{cases}$$

Here ρ represents density of water



Why do we not consider ρ_{air} in this analysis?

- a. Because it is a reasonable approximation
- b. Because it simplifies the analysis
- c. Both of the above

Euler's equation for incompressible fluid :

$$\frac{d\mathbf{v}}{dt} = f_{\text{applied}} - \frac{\nabla p}{\rho} = -g\hat{\mathbf{z}} - \frac{\nabla p}{\rho}$$

Assume that $v_z \ll v_x, v_y \Rightarrow -g - \frac{1}{\rho} \frac{\partial p}{\partial z} \approx 0$

$$\Rightarrow p(x, y, z, t) = p_0 + \rho g (\zeta(x, y, t) + h - z) \quad \text{within the water}$$

Horizontal fluid motions (keeping leading terms):

$$\frac{dv_x}{dt} \approx \frac{\partial v_x}{\partial t} = -\frac{1}{\rho} \frac{\partial p}{\partial x} = -g \frac{\partial \zeta}{\partial x}$$

$$\frac{dv_y}{dt} \approx \frac{\partial v_y}{\partial t} = -\frac{1}{\rho} \frac{\partial p}{\partial y} = -g \frac{\partial \zeta}{\partial y}$$




Consider a surface $\zeta(x,t)$ wave moving in the x -direction in a channel of width $b(x)$ and height $h(x)$:

Continuity condition in integral form:

$$\frac{d}{dt} \int_V \rho dV + \int_A \rho \mathbf{v} \cdot d\mathbf{A} = 0$$

\uparrow $\int_V \rho dV$ \uparrow $\int_A \rho \mathbf{v} \cdot d\mathbf{A}$
 $b(x)(h(x) + \zeta(x,t)) dx$ $b(x)(h(x) + \zeta(x,t)) \hat{\mathbf{x}}$




Evaluating continuity condition:

$$b(x) \frac{\partial \zeta}{\partial t} = - \frac{\partial}{\partial x} (h(x) b(x) v(x,t))$$

Some details

Continuity condition in integral form:



$$\frac{d}{dt} \int_V \rho dV + \int_A \rho \mathbf{v} \cdot d\mathbf{A} = 0$$

\uparrow $\int_V \rho dV$ \uparrow $\int_A \rho \mathbf{v} \cdot d\mathbf{A}$
 $b(x)(h(x) + \zeta(x, t)) dx$ $b(x)(h(x) + \zeta(x, t)) \hat{\mathbf{x}}$


Here, we are assuming that ρ is constant

$$\begin{aligned} \frac{d}{dt} \int_V \rho dV + \int_A \rho \mathbf{v} \cdot d\mathbf{A} &= \rho \int b(x) \frac{\partial \zeta}{\partial t} dx + \rho \int \frac{\partial}{\partial x} (b(x)(h(x) + \zeta(x, t))v(x, t)) dx = 0 \\ \Rightarrow b(x) \frac{\partial \zeta}{\partial t} &= - \frac{\partial}{\partial x} (h(x)b(x)v(x, t)) \end{aligned}$$



From continuity condition:

$$b(x) \frac{\partial \zeta}{\partial t} = - \frac{\partial}{\partial x} (h(x)b(x)v(x,t))$$



Example (Problem 10.3):

$$b(x) = b_0 \quad h(x) = \kappa x$$

(A special case sometimes found at a beach.)

$$b_0 \frac{\partial \zeta}{\partial t} = - \frac{\partial}{\partial x} ((\kappa x)b_0 v(x,t))$$

$$\frac{\partial \zeta}{\partial t} = -\kappa \left(v + x \frac{\partial v}{\partial x} \right)$$

From Newton-Euler equation:

$$\frac{dv}{dt} \approx \frac{\partial v}{\partial t} = -g \frac{\partial \zeta}{\partial x}$$

Example continued

$$\frac{\partial \zeta}{\partial t} = -\kappa \left(v + x \frac{\partial v}{\partial x} \right) \quad \Rightarrow \quad \frac{\partial^2 \zeta}{\partial t^2} = -\kappa \left(\frac{\partial v}{\partial t} + x \frac{\partial^2 v}{\partial x \partial t} \right)$$

$$\frac{\partial v}{\partial t} = -g \frac{\partial \zeta}{\partial x} \quad \Rightarrow \quad \frac{\partial^2 \zeta}{\partial t^2} = \kappa g \left(\frac{\partial \zeta}{\partial x} + x \frac{\partial^2 \zeta}{\partial x^2} \right)$$

It can be shown that a solution can take the form:

$$\zeta(x, t) = C J_0 \left(\frac{2\omega}{\sqrt{\kappa g}} \sqrt{x} \right) \cos(\omega t)$$

Note that $J_0(u)$ satisfies the equation: $\left(\frac{d^2}{du^2} + \frac{1}{u} \frac{d}{du} + 1 \right) J_0(u) = 0$

Therefore, for $u = \frac{2\omega}{\sqrt{\kappa g}} \sqrt{x}$

$$\left(x \frac{d^2}{dx^2} + \frac{d}{dx} \right) J_0(u) = \frac{\omega^2}{\kappa g} \left(\frac{d^2}{du^2} + \frac{1}{u} \frac{d}{du} \right) J_0(u) = -\frac{\omega^2}{\kappa g} J_0(u)$$

Therefore, for $u = \frac{2\omega}{\sqrt{\kappa g}} \sqrt{x} \Rightarrow \frac{1}{\sqrt{x}} = \frac{2\omega}{\sqrt{\kappa g}} \frac{1}{u}$

$$\left(x \frac{d^2}{dx^2} + \frac{d}{dx} \right) J_0(u) = \frac{\omega^2}{\kappa g} \left(\frac{d^2}{du^2} + \frac{1}{u} \frac{d}{du} \right) J_0(u) = -\frac{\omega^2}{\kappa g} J_0(u)$$

Detail: $\frac{dJ_0(u)}{dx} = \frac{dJ_0(u)}{du} \frac{\omega}{\sqrt{\kappa g}} \frac{1}{\sqrt{x}}$

$$\frac{d^2 J_0(u)}{dx^2} = \frac{d^2 J_0(u)}{du^2} \left(\frac{\omega}{\sqrt{\kappa g}} \frac{1}{\sqrt{x}} \right)^2 - \frac{dJ_0(u)}{du} \frac{\omega}{2\sqrt{\kappa g}} \frac{1}{x\sqrt{x}}$$

Therefore:
$$\left(x \frac{d^2}{dx^2} + \frac{d}{dx} \right) J_0(u) = \left(\frac{\omega^2}{\kappa g} \frac{d^2 J_0(u)}{du^2} + \frac{dJ_0(u)}{du} \frac{\omega}{2\sqrt{\kappa g}} \frac{1}{\sqrt{x}} \right)$$

$$= \frac{\omega^2}{\kappa g} \left(\frac{d^2 J_0(u)}{du^2} + \frac{dJ_0(u)}{du} \frac{1}{u} \right)$$

Example continued


$$\frac{\partial^2 \zeta}{\partial t^2} = \kappa g \left(\frac{\partial \zeta}{\partial x} + x \frac{\partial^2 \zeta}{\partial x^2} \right)$$

$$\Rightarrow \zeta(x, t) = C J_0 \left(\frac{2\omega\sqrt{x}}{\sqrt{\kappa g}} \right) \cos(\omega t)$$

Check:

$$-\omega^2 C J_0 \left(\frac{2\omega\sqrt{x}}{\sqrt{\kappa g}} \right) \cos(\omega t) = \kappa g \left(\frac{\partial}{\partial x} + x \frac{\partial^2}{\partial x^2} \right) C J_0 \left(\frac{2\omega\sqrt{x}}{\sqrt{\kappa g}} \right) \cos(\omega t)$$

$$\zeta(x,t) = CJ_0\left(\frac{2\omega}{\sqrt{\kappa g}}\sqrt{x}\right)\cos(\omega t)$$




Imagine watching the waves at a beach – can you visualize the configuration for the surface wave pattern to approximation this situation?

- a. Long flat beach
- b. Beach in which average water level increases
- c. Beach in which average water level decreases





A simpler example:




Continuity condition:

$$b(x) \frac{\partial \zeta}{\partial t} = - \frac{\partial}{\partial x} (h(x)b(x)v(x,t))$$

Special case, where b and h are constant --
For constant b and h :

$$\frac{\partial \zeta}{\partial t} = -h \frac{\partial}{\partial x} (v(x,t))$$

Example with b and h constant -- continued



Continuity condition for flow of incompressible fluid:

$$\frac{\partial \zeta}{\partial t} + h \nabla \cdot \mathbf{v} = 0$$

From horizontal flow relations: $\frac{\partial \mathbf{v}}{\partial t} = -g \nabla \zeta$

Equation for surface function: $\frac{\partial^2 \zeta}{\partial t^2} - gh \nabla^2 \zeta = 0$



For uniform channel:

Surface wave equation:


$$\frac{\partial^2 \zeta}{\partial t^2} - c^2 \nabla^2 \zeta = 0 \qquad c^2 = gh$$

More complete analysis finds:

$$c^2 = \frac{g}{k} \tanh(kh) \qquad \text{where } k = \frac{2\pi}{\lambda}$$

More details: -- recall setup --

Consider a container of water with average height h and surface $h + \zeta(x, y, t)$



Equations describing fluid itself (without boundaries)

Euler's equation for incompressible fluid:

$$\frac{d\mathbf{v}}{dt} = \frac{\partial \mathbf{v}}{\partial t} + \mathbf{v} \cdot \nabla \mathbf{v} = \frac{\partial \mathbf{v}}{\partial t} + \nabla \left(\frac{1}{2} v^2 \right) + \mathbf{v} \times (\nabla \times \mathbf{v}) = -\nabla U - \frac{\nabla p}{\rho}$$


Assume that $\nabla \times \mathbf{v} = 0$ (irrotational flow) $\Rightarrow \mathbf{v} = -\nabla \Phi$

$$\Rightarrow \nabla \left(-\frac{\partial \Phi}{\partial t} + \frac{1}{2} v^2 + U + \frac{p}{\rho} \right) = 0$$

$$\Rightarrow -\frac{\partial \Phi}{\partial t} + \frac{1}{2} v^2 + U + \frac{p}{\rho} = \text{constant (within the fluid)}$$

For the same system, the continuity condition becomes

$$\nabla \cdot \mathbf{v} = -\nabla^2 \Phi = 0$$



Within fluid: $0 \leq z \leq h + \zeta$

$$-\frac{\partial \Phi}{\partial t} + \frac{1}{2} v^2 + g(z - h) = \text{constant} \quad (\text{We have absorbed } p_0 \text{ in "constant"})$$

$$-\nabla^2 \Phi = 0$$

At surface: $z = h + \zeta$ with $\zeta = \zeta(x, y, t)$

$$\frac{d\zeta}{dt} = \frac{\partial \zeta}{\partial t} + v_x \frac{\partial \zeta}{\partial x} + v_y \frac{\partial \zeta}{\partial y} \quad \text{where } v_{x,y} = v_{x,y}(x, y, h + \zeta, t)$$

Full equations:

Within fluid: $0 \leq z \leq h + \zeta$

$$-\frac{\partial \Phi}{\partial t} + \frac{1}{2} v^2 + g(z - h) = \text{constant} \quad (\text{We have absorbed } p_0 \text{ in "constant"})$$

$$-\nabla^2 \Phi = 0$$

At surface: $z = h + \zeta$ with $\zeta = \zeta(x, y, t)$

$$\frac{d\zeta}{dt} = \frac{\partial \zeta}{\partial t} + v_x \frac{\partial \zeta}{\partial x} + v_y \frac{\partial \zeta}{\partial y} \quad \text{where } v_{x,y} = v_{x,y}(x, y, h + \zeta, t)$$

Linearized equations:

$$\text{For } 0 \leq z \leq h + \zeta : \quad -\frac{\partial \Phi}{\partial t} + g(z - h) = 0 \quad -\nabla^2 \Phi = 0$$

$$\text{At surface: } z = h + \zeta \quad \frac{d\zeta}{dt} = \frac{\partial \zeta}{\partial t} = v_z(x, y, h + \zeta, t)$$

$$-\frac{\partial \Phi(x, y, h + \zeta, t)}{\partial t} + g\zeta = 0$$

For simplicity, keep only linear terms and assume that horizontal variation is only along x :

For $0 \leq z \leq h + \zeta$:
$$\nabla^2 \Phi = \left(\frac{\partial^2}{\partial z^2} + \frac{\partial^2}{\partial x^2} \right) \Phi(x, z, t) = 0$$

Consider a periodic waveform: $\Phi(x, z, t) = Z(z) \cos(k(x - ct))$

$$\Rightarrow \left(\frac{d^2}{dz^2} - k^2 \right) Z(z) = 0$$

Boundary condition at bottom of tank: $v_z(x, 0, t) = 0$

$$\Rightarrow \frac{dZ}{dz}(0) = 0 \quad Z(z) = A \cosh(kz)$$

For simplicity, keep only linear terms and assume that horizontal variation is only along x – continued:

$$\text{At surface: } z = h + \zeta \quad \frac{\partial \zeta}{\partial t} = v_z(x, h + \zeta, t) = -\frac{\partial \Phi(x, h + \zeta, t)}{\partial z}$$

$$-\frac{\partial \Phi(x, h + \zeta, t)}{\partial t} + g\zeta = 0$$

$$-\frac{\partial^2 \Phi(x, h + \zeta, t)}{\partial t^2} + g \frac{\partial \zeta}{\partial t} = -\frac{\partial^2 \Phi(x, h + \zeta, t)}{\partial t^2} - g \frac{\partial \Phi(x, h + \zeta, t)}{\partial z} = 0$$

$$\text{For } \Phi(x, (h + \zeta), t) = A \cosh(k(h + \zeta)) \cos(k(x - ct))$$


$$A \cosh(k(h + \zeta)) \cos(k(x - ct)) \left(k^2 c^2 - gk \frac{\sinh(k(h + \zeta))}{\cosh(k(h + \zeta))} \right) = 0$$

$$\Rightarrow c^2 = \frac{g \sinh(k(h + \zeta))}{k \cosh(k(h + \zeta))}$$

For simplicity, keep only linear terms and assume that horizontal variation is only along x – continued:

$$c^2 = \frac{g \sinh(k(h + \zeta))}{k \cosh(k(h + \zeta))} = \frac{g}{k} \tanh(k(h + \zeta))$$

Assuming $\zeta \ll h$: $c^2 = \frac{g}{k} \tanh(kh)$ $\lambda = \frac{2\pi}{k}$



For simplicity, keep only linear terms and assume that horizontal variation is only along x – continued:

$$c^2 \approx \frac{g}{k} \tanh(kh) \quad \text{For } \lambda \gg h, \quad c^2 \approx gh$$

$$\Phi(x, z, t) = A \cosh(kz) \cos(k(x - ct))$$


$$\zeta(x, t) = \frac{1}{g} \frac{\partial \Phi(x, h + \zeta, t)}{\partial t} \approx \frac{kc}{g} A \cosh(kh) \sin(k(x - ct))$$

Note that for $\lambda \gg h$, $c^2 \approx gh$

(solutions are consistent with previous analysis)



General problem
including
non-linearities



Within fluid: $0 \leq z \leq h + \zeta$

$$-\frac{\partial \Phi}{\partial t} + \frac{1}{2} v^2 + g(z - h) = \text{constant} \quad (\text{We have absorbed}$$

$$-\nabla^2 \Phi = 0 \quad p_0 \text{ in our constant.})$$

At surface: $z = h + \zeta$ with $\zeta = \zeta(x, y, t)$

$$\frac{d\zeta}{dt} = \frac{\partial \zeta}{\partial t} + v_x \frac{\partial \zeta}{\partial x} + v_y \frac{\partial \zeta}{\partial y} \quad \text{where } v_{x,y} = v_{x,y}(x, y, h + \zeta, t)$$