

PHY 341/641


Thermodynamics and Statistical Physics

Lecture 2

1. Continued discussion of microscopic models (Chapter 1)
 - a. Notion of equilibrium in statistical mechanics/thermodynamics
 - b. Macrostates/microstates
2. Introduction to thermodynamics (Chapter 2)
 - a. Definition of “the system”
 - b. Thermodynamic variables (T , P , V , N , ...)
 - c. First law of thermodynamics

Review from last time --

Chapter 1 – From Microscopic to Macroscopic Behavior




Assignment: Read Chapter 1 (quickly) during this week and checkout some of the corresponding simulations (HW 1 and HW 2) due Monday 1/23.

“The purpose of this introductory [material] is to whet your appetite... ” The chapter introduces a lot of the concepts that we will use (more carefully) throughout the course.

What will you learn?


Energy Analysis



Comment on simulation tools

Molecular dynamics

$$m_i \frac{d^2 \mathbf{r}_i}{dt^2} = \mathbf{F}_i$$



$$\mathbf{F}_i = -\nabla_i \sum_{j \neq i} u_{pair}(|\mathbf{r}_i - \mathbf{r}_j|)$$


Example model pair potential (Lennard-Jones):

$$u_{LJ}(r) = 4\epsilon \left[\left(\frac{\sigma}{r} \right)^{12} - \left(\frac{\sigma}{r} \right)^6 \right]$$

For an “ideal gas”--

$$m_i \frac{d^2 \mathbf{r}_i}{dt^2} = \mathbf{F}_i \equiv 0$$

Modeling of a dilute gas of non-interacting particles:



$$P_{L \rightarrow R} = \frac{n}{N}$$


$$P_{R \rightarrow L} = \frac{N - n}{N}$$

Enumeration of possibilities for N=4


Microstates				n	$P_{L \rightarrow R}$
L	L	L	L	4	1/16
L	L	L	R	3	4/16
L	L	R	L		
L	R	L	L		
R	L	L	L		
L	L	R	R	2	6/16
L	R	R	L		
R	R	L	L		
L	R	L	R		
R	L	R	L		
R	L	L	R		
R	R	R	L	1	4/16
R	R	L	R		
R	L	R	R		
L	R	R	R		
R	R	R	R	0	1/16

Simulation available from: http://www.compadre.org/STP/stp_ApproachToEquilibrium.jar


*Note: in order to easily control the simulation, you need to use:
Display → Switch GUI*



$N=64$, $t \approx 1000$ $\langle n \rangle = 32.08$, $\langle n^2 \rangle = 1,067.13$, $\sigma^2 = 38.12$, $\sigma/\langle n \rangle = 0.19$




$N=256$, $t \approx 1000$ $\langle n \rangle = 149.57$, $\langle n^2 \rangle = 23,086.17$, $\sigma^2 = 714.56$, $\sigma/\langle n \rangle = 0.18$



Macroscopic viewpoint – thermodynamics

(start reading Chapter 2)



Variables of thermodynamics

➤ Temperature

- ❖ T

- ❖ *Zeroth law of thermodynamics: Two systems in thermal equilibrium with a third system are in thermal equilibrium with each other.*

➤ Pressure

- ❖ $P=F/A$

➤ Volume

- ❖ V

➤ Number of particles

- ❖ N

The relationships between these variables depends on the “equation of state” of the system.

Examples of “Equations of State”

Ideal gas equation of state :


$$PV = Nk_B T \qquad k_B = 1.38 \times 10^{-23} \text{ J/K}$$

van der Waals equation of state :

$$\left(P + \frac{N^2}{V^2} a \right) (V - Nb) = Nk_B T$$

Comparison of

- Ideal gas equation
- van der Waals gas equation



In general the “state” of a system will depend on the thermodynamic variables. For example, the internal energy:

$$U = U(N, T, P, V)$$


Thermodynamic “processes” $\rightarrow$ change the state of the system

$$U_1(N_1, T_1, P_1, V_1) \Rightarrow U_2(N_2, T_2, P_2, V_2)$$

Thermodynamic process -- work (performed ON the system)

$$dW = -Fdx = -PA dx = -PdV$$


$$W_{1 \rightarrow 2} = -\int_{V_1}^{V_2} P(T, V) dV$$



For an ideal gas at constant T :

$$W_{1 \rightarrow 2} = -\int_{V_1}^{V_2} P(T, V) dV = -\int_{V_1}^{V_2} \frac{NkT}{V} dV = -NkT \ln \frac{V_2}{V_1}$$

Work performed during a cyclic process:



$$W_{net} = -(P_{high} - P_{low})(V_{high} - V_{low})$$

Thermodynamic process -- heat (added TO the system)

Q

For heat added TO the system: $Q > 0$

For heat withdrawn FROM the system: $Q < 0$

First law of thermodynamics

$$U_2 - U_1 \equiv \Delta U = W + Q$$

Example: Adiabatic expansion $V_1 \rightarrow V_2$ for an ideal gas system

$$U_2 - U_1 \equiv \Delta U = W + Q = W$$

$$PV = NkT$$

To be continued.....