

PHY 341/641 Thermodynamics and Statistical Physics

Lecture 20

Methodologies of statistical mechanics. (Chapter 5 in STP)

- Phase transitions in spin systems
- Mean field approximation

3/9/2012

PHY 341/641 Spring 2012 -- Lecture 20

1

9	2/06/2012	Probability distributions	3.4-3.5	HW 9	2/8/2012
10	2/08/2012	Continuous distributions/Central limit theorem	3.6-3.10	HW 10	2/10/2012
11	2/10/2012	Introduction to statistical mechanics	4.1-4.2	HW 11	2/13/2012
12	2/13/2012	Enumeration of microstates	4.3	HW 12	2/15/2012
13	2/15/2012	Many particle systems	4.4-4.5	HW 13	2/17/2012
14	2/17/2012	Microcanonical ensemble	4.6	HW 14	2/20/2012
15	2/20/2012	Canonical ensemble	4.7-4.8	HW 15	2/22/2012
16	2/22/2012	Grand canonical ensemble	4.9-4.12	HW 16	2/24/2012
17	2/24/2012	Introduction to magnetic systems	5.1-5.5		
	2/27/2012	APS -- no class; take-home exam			
	2/29/2012	APS -- no class; take-home exam			
	3/02/2012	APS -- no class; take-home exam			
18	3/05/2012	Exam due -- Ising model	5.5	HW 17	03/07/2012
19	3/07/2012	Ising model	5.6-5.7	HW 18	03/09/2012
20	3/09/2012	Phase transformation	5.8-5.10		
	3/12/2012	Spring Break			
	3/14/2012	Spring Break			
	3/16/2012	Spring Break			
21	3/19/2012	Many particle systems	6.1-6.2		

3/9/2012

PHY 341/641 Spring 2012 -- Lecture 20

2


Ising model in various lattices -- mean field solutions

Microstate s :


$$E_s = -J \sum_{i,j(nn)} s_i s_j - H \sum_{i=1}^N s_i$$

Mean fields approximat ion :

$$E_s^{MF} = -J \sum_i s_i q \langle s_j \rangle - H \sum_{i=1}^N s_i$$



$q=4/2$



$q=6/2$

$$\langle H_{eff} \rangle = Jq \langle s_j \rangle + H \equiv Jqm + H$$

Note: textbooks differ with counting neighbors q without double counting. Here q implies "unique" neighbors.

3/9/2012

PHY 341/641 Spring 2012 -- Lecture 20

3

Self-consistency condition for mean field treatment

$$\langle H_{\text{eff}} \rangle = Jq \langle s_j \rangle + H \equiv Jqm + H$$

$$Z_1 = \sum_{s_1 = \pm 1} e^{-\beta \langle H_{\text{eff}} \rangle s_1} = 2 \cosh(\beta(Jqm + H))$$

$$\frac{F}{N} = -kT \ln Z_1$$

$$m = -\frac{1}{N} \frac{\partial F}{\partial H} = \tanh(\beta(Jqm + H))$$


3/9/2012

PHY 341/641 Spring 2012 -- Lecture 20

4

Mean field self-consistency condition:

$$m = \tanh(\beta(Jqm + H))$$



Condition for non-trivial solution for m at $H = 0$:
 $\beta J q \geq 1$

→ Mean field solutions exhibit "critical behavior" (phase transition) at $\beta_c J q = 1$.

3/9/2012


PHY 341/641 Spring 2012 -- Lecture 20

5

Mean field self-consistency condition for $H=0$: $m = \tanh(\beta(Jqm))$

Define: $\beta_c J q = 1$

$$m = \tanh(\beta(Jqm)) = \tanh\left(\frac{\beta}{\beta_c} m\right) = \tanh\left(\frac{T_c}{T} m\right)$$



3/9/2012

PHY 341/641 Spring 2012 -- Lecture 20

6

Summary of results for mean field treatment of Ising model

$$Z_N = [2 \cosh(\beta(Jqm + H))]^N$$

$$F = -kTN \ln[2 \cosh(\beta(Jqm + H))]$$

$$\text{where: } m = \tanh(\beta(Jqm + H))$$

Internal energy :

$$E = -\frac{\partial \ln Z_N}{\partial \beta}$$

$$= -N(Jqm + H) [\tanh(\beta(Jqm + H))]$$

$$= -N(Jqm + H)m$$

3/9/2012

PHY 341/641 Spring 2012 -- Lecture 20

7

Summary of results for mean field treatment of Ising model (continued)

Internal energy for $H = 0$:

$$E = -NJqm^2$$

$$\text{where } m = \tanh(\beta(Jqm))$$

Heat capacity :

$$C = \left(\frac{\partial E}{\partial T} \right)_N = 2Nk\beta^2 Jqm \left(\frac{\partial m}{\partial \beta} \right)_N$$

$$\left(\frac{\partial m}{\partial \beta} \right)_N = \left(Jqm + \beta Jq \left(\frac{\partial m}{\partial \beta} \right)_N \right) \text{sech}^2(\beta(Jqm))$$

$$\left(\frac{\partial m}{\partial \beta} \right)_N = \frac{Jqm}{\cosh^2(\beta(Jqm)) - \beta Jq}$$

$$C = \frac{2Nk\beta^2 J^2 q^2 m^2}{\cosh^2(\beta(Jqm)) - \beta Jq}$$

3/9/2012

PHY 341/641 Spring 2012 -- Lecture 20

8

Summary of results for mean field treatment of Ising model (continued)

Behavior of heat capacity for $H = 0$

in terms of $T_c = \frac{Jq}{k}$:

$$C = \frac{2Nk\beta^2 J^2 q^2 m^2}{\cosh^2(\beta(Jqm)) - \beta Jq}$$

$$= \frac{2Nkm^2}{\cosh^2\left(\frac{T_c}{T}m\right) - \left(\frac{T_c}{T}\right)} \left(\frac{T_c}{T}\right)^2$$

$$\text{Note that: } m = \begin{cases} \tanh\left(\frac{T_c}{T}m\right) & \text{for } T < T_c \\ 0 & \text{for } T > T_c \end{cases}$$

3/9/2012

PHY 341/641 Spring 2012 -- Lecture 20

9

Summary of results for mean field treatment of Ising model
(continued)

Behavior for T near T_c where m is small

$$m = \begin{cases} \tanh\left(\frac{T_c}{T} m\right) & \text{for } T < T_c \\ 0 & \text{for } T > T_c \end{cases}$$

for $T < T_c$ and $T \approx T_c$:

$$m \approx \frac{T_c}{T} m - \frac{1}{3} \left(\frac{T_c}{T} m\right)^3 + \dots$$

$$m \approx \sqrt{3} \left(\frac{T}{T_c}\right) \left(1 - \frac{T}{T_c}\right)^{1/2}$$

3/9/2012

PHY 341/641 Spring 2012 -- Lecture 20

10

Summary of results for mean field treatment of Ising model
(continued) -- behavior for T near T_c where m is small

$$m \approx \sqrt{3} \left(\frac{T}{T_c}\right) \left(1 - \frac{T}{T_c}\right)^{1/2}$$

$$C = \frac{2Nkm^2}{\cosh^2\left(\frac{T_c}{T} m\right) - \left(\frac{T_c}{T}\right)} \left(\frac{T_c}{T}\right)^2$$

$$C \equiv 0 \quad \text{for } T > T_c$$

3/9/2012

PHY 341/641 Spring 2012 -- Lecture 20

11

Summary of results for mean field treatment of Ising model
(continued) -- behavior for T near T_c where m is small

$$m \approx \sqrt{3} \left(\frac{T}{T_c}\right) \left(1 - \frac{T}{T_c}\right)^{1/2}$$

$$C = \frac{2Nkm^2}{\cosh^2\left(\frac{T_c}{T} m\right) - \left(\frac{T_c}{T}\right)} \left(\frac{T_c}{T}\right)^2$$

for $T < T_c$ and $T \approx T_c$:

$$C \approx \frac{2Nk}{1 - \left(\frac{T_c}{T}\right) + 3\left(1 - \frac{T}{T_c}\right)} \left(\frac{T_c}{T}\right)^2 \cdot 3 \left(\frac{T}{T_c}\right)^2 \left(1 - \frac{T}{T_c}\right) \approx \frac{6Nk}{3 - \frac{T_c}{T}} \approx 3Nk$$

3/9/2012

PHY 341/641 Spring 2012 -- Lecture 20

12

Behavior of magnetic susceptibility (in scaled units):

$$m = \tanh(\beta(Jqm + H))$$

$$\chi(T, H) = \left(\frac{\partial m}{\partial H} \right)_T$$

$$\left(\frac{\partial m}{\partial H} \right)_T = \left(\beta + \beta Jq \left(\frac{\partial m}{\partial H} \right)_T \right) \operatorname{sech}^2(\beta(Jqm + H))$$

$$\chi(T, H) = \left(\frac{\partial m}{\partial H} \right)_T = \frac{\beta}{\cosh^2(\beta(Jqm + H)) - \beta Jq}$$

For $H = 0$:

$$\chi(T, 0) = \frac{1}{Jq} \frac{\frac{T_c}{T}}{\cosh^2\left(\frac{T_c}{T} m\right) - \frac{T_c}{T}}$$

3/9/2012

PHY 341/641 Spring 2012 -- Lecture 20

13

Behavior of magnetic susceptibility (in scaled units) -- continued:

$$\chi(T, 0) = \frac{1}{Jq} \frac{\frac{T_c}{T}}{\cosh^2\left(\frac{T_c}{T} m\right) - \frac{T_c}{T}}$$

For $T > T_c$; $m = 0$

$$\chi(T, 0) = \frac{1}{Jq} \frac{\frac{T_c}{T}}{1 - \frac{T_c}{T}} = \frac{T_c}{Jq} \frac{1}{T - T_c} \quad \text{Curie - Weiss relation}$$

For $T < T_c$ and $T \approx T_c$; $m \approx \sqrt{3} \left(\frac{T}{T_c} \right) \left(1 - \frac{T}{T_c} \right)^{1/2}$

$$\cosh^2\left(\frac{T_c}{T} m\right) \approx 1 + 3 \left(1 - \frac{T}{T_c} \right)$$

$$\chi(T, 0) \approx \frac{1}{Jq} \frac{\frac{T_c}{T}}{\left(1 - \frac{T}{T_c} \right) \left(3 - \frac{T_c}{T} \right)} \approx \frac{T_c}{2Jq} \frac{1}{T_c - T}$$

3/9/2012

PHY 341/641 Spring 2012 -- Lecture 20

14

Behavior of magnetization at T_c :

$$m = \tanh(\beta(Jqm + H)) = \tanh\left(\frac{T_c}{T} m + \frac{H}{kT}\right)$$

At $T = T_c$:

$$m(T_c) = \tanh\left(m(T_c) + \frac{H}{kT_c}\right) \approx m(T_c) + \frac{H}{kT_c} - \frac{1}{3} \left(m(T_c) + \frac{H}{kT_c} \right)^3$$

$$\Rightarrow m(T_c) \approx \left(\frac{3H}{kT_c} \right)^{1/3}$$

3/9/2012

PHY 341/641 Spring 2012 -- Lecture 20

15

Critical exponents:

$$\varepsilon \equiv \left(1 - \frac{T}{T_c}\right)$$

Quantity	Critical behavior	values of the exponents		
		$d = 2$ (exact)	$d = 3$	mean-field theory
specific heat	$C \sim \varepsilon^{-\alpha}$	0 (logarithmic)	0.113	0 (jump)
order parameter	$m \sim \varepsilon^\beta$	1/8	0.324	1/2
susceptibility	$\chi \sim \varepsilon^{-\gamma}$	7/4	1.238	1
equation of state ($\epsilon = 0$)	$m \sim H^{1/\delta}$	15	4.82	3
correlation length	$\xi \sim \varepsilon^{-\nu}$	1	0.629	1/2
correlation function $\epsilon = 0$	$G(r) \sim 1/r^{d-2+\eta}$	1/4	0.031	0

Table 5.1: Values of the critical exponents for the Ising model in two and three dimensions. The values of the critical exponents of the Ising model are known exactly in two dimensions and are ratios of integers. The results in three dimensions are not ratios of integers and are approximate. The exponents predicted by mean-field theory are discussed in Sections 5.7 and 9.1, pages 236 and 334.
