

PHY 341/641
Thermodynamics and Statistical Physics

Lecture 8

1. Complete discussion of example 2.21
2. Begin introduction to probability concepts (Chapter 3 in STP)

2/3/2012

PHY 341/641 Spring 2012 -- Lecture 8

1

Example from your text on functional dependences in black body radiation following analysis by Boltzmann in 1884.

Assume that we have a system characterized by :

Internal energy: $U(T, V) \equiv u(T)V$ Entropy: $S(T, V)$

Pressure: $P(T) \equiv \frac{1}{3}u(T)$

$$dS = \frac{V}{T} \frac{du}{dT} dT + \left(\frac{\partial S}{\partial V} \right)_T dV$$

$$\left(\frac{\partial S}{\partial T} \right)_V = \frac{V}{T} \frac{du}{dT} \quad \left(\frac{\partial S}{\partial V} \right)_T = \left(\frac{\partial}{\partial T} \right)_V \left(\left(\frac{\partial S}{\partial V} \right)_T \right) \Rightarrow u(T) = KT^4, \quad S(T, V) = \frac{4}{3}KT^3V$$

2/3/2012

PHY 341/641 Spring 2012 -- Lecture 8

2

Example from your text on functional dependences in black body radiation following analysis by Boltzmann in 1884 – continued --

Some details :

$$dS = \left(\frac{\partial S}{\partial T} \right)_V dT + \left(\frac{\partial S}{\partial V} \right)_T dV = \frac{d(uV)}{T} + \frac{P}{T} dV$$

$$\left(\frac{\partial S}{\partial T} \right)_V = \frac{V}{T} \frac{du}{dT} \quad \left(\frac{\partial S}{\partial V} \right)_T = \frac{u}{T} + \frac{u}{3T} = \frac{4u}{3T}$$

$$\left(\frac{\partial}{\partial V} \right)_T \left(\frac{V}{T} \frac{du}{dT} \right) = \left(\frac{\partial}{\partial T} \right)_V \left(\frac{4u}{3T} \right) \Rightarrow \frac{du}{dT} = 4 \frac{u}{T}$$

$$\Rightarrow u(T) = KT^4 \quad S(T, V) = \frac{4}{3}KT^3V$$

2/3/2012

PHY 341/641 Spring 2012 -- Lecture 8

3

Concepts of Probability – Chapter 3 of STP

Probability analysis is helpful for correlating microscopic and macroscopic physics.

Example – 2-sided coin toss

$P(\text{heads}) = \frac{1}{2}$
 $P(\text{tails}) = \frac{1}{2}$

Suppose we have 4 coins tossed at once. On any given toss, what is the probability of having n occurrences of heads?

2/3/2012 PHY 341/641 Spring 2012 -- Lecture 8 4

Outcomes for tossing 4 coins


Configurations	n	$P(n)$
TTTT	0	1/16
HTTT,THTT,TTHT,TTTH	1	4/16
TTHH,THTH,HTHT,HTTH	2	6/16
TTHH,HTHH,HHTH,HHHT	3	4/16
HHHH	4	1/16

Simulation of coin tosses:
[stp_CoinToss.jar](#)


2/3/2012 PHY 341/641 Spring 2012 -- Lecture 8 5

Distribution of outcomes for tossing N coins

$N=4$ -- 316 trials



$N=256$ -- 443 trials



2/3/2012 PHY 341/641 Spring 2012 -- Lecture 8 6

Mathematics of probability theory

- For each event i , we assign a probability $P(i)$ with the conditions: $P(i) \geq 0$ and $\sum_i P(i) = 1$
- For two independent events i and j , the probability of i or j is

$$P(i \text{ or } j) = P(i) + P(j)$$

- For two independent events i and j , the probability of i and j is

$$P(i \text{ and } j) = P(i)P(j)$$

2/3/2012

PHY 341/641 Spring 2012 -- Lecture 8

7

Example (similar to 3.8 of your textbook)

Suppose that in a certain class of 100 students the grade distribution averaged over many years is given as shown in the second column. What is the probability of getting each grade assuming that the distribution remains the same:

Grade	100 student Distribution	Probability
A	16	0.16
B	32	0.32
C	48	0.48
D	4	0.04
F	0	0.00

2/3/2012

PHY 341/641 Spring 2012 -- Lecture 8

8

Calculation of averages

Define: $P(i) \equiv$ probability of x_i

$$\Rightarrow \text{Average } \langle x \rangle = \sum_{i=1}^n x_i P(i)$$

$$\Rightarrow \text{Moment } \langle x^m \rangle = \sum_{i=1}^n x_i^m P(i)$$

$$\Rightarrow \text{Standard deviation } \sigma = \sqrt{\langle \Delta x^2 \rangle} \equiv \sqrt{\langle (x - \langle x \rangle)^2 \rangle}$$

note that : $\langle \Delta x^2 \rangle = \langle x^2 - \langle x \rangle^2 \rangle$

2/3/2012

PHY 341/641 Spring 2012 -- Lecture 8

9

Example:

Grade	100 student Distribution	Probability
A (4)	16	0.16
B (3)	32	0.32
C (2)	48	0.48
D (1)	4	0.04
F (0)	0	0.00

$$\Rightarrow \text{Average } \langle G \rangle = \sum_{i=1}^n G_i P(i) = 2.6$$

$$\Rightarrow \text{Standard deviation } \sigma = \sqrt{\langle \Delta x^2 \rangle} \equiv \sqrt{\langle (x - \langle x \rangle)^2 \rangle} = 0.8$$

2/3/2012

PHY 341/641 Spring 2012 -- Lecture 8

10

Example: Dice throws



Figure 3.1: The six possible outcomes of the toss of a die.

Example 3.10. On average, how many times must a die be thrown until "4" appears?

Let p = probability of getting 4 on one throw ($p = 1/6$)

Let q = probability of not getting 4 on one throw

Probability of first getting 4 on n th throw: $P_n = pq^{n-1}$

$$\begin{aligned} \text{Mean \# of throws: } m &= \sum_{n=1}^{\infty} npq^{n-1} = p \frac{d}{dq} \sum_{n=0}^{\infty} q^n \\ &= p \frac{d}{dq} \frac{1}{1-q} = p \frac{1}{(1-q)^2} = \frac{1}{p} \end{aligned}$$

2/3/2012

PHY 341/641 Spring 2012 -- Lecture 8

11