

PHY 341/641
Thermodynamics and Statistical Physics

Lecture 9

1. Probability concepts (Chapter 3 in STP)
 - A. Bayes' theorem
 - B. Binomial distribution

2/6/2012

PHY 341/641 Spring 2012 -- Lecture 8

1

Construction of an "uncertainty" function

Define: $\Omega \equiv$ total number of outcomes of an "experiment"

Define: $S(\Omega) \equiv$ uncertainty function

Necessary properties:

$$S(\Omega = 1) = 0$$

$$\text{If } \Omega_1 > \Omega_2 \text{ then } S(\Omega_1) > S(\Omega_2)$$

$$\text{If } \Omega = \Omega_1 \Omega_2 \text{ then } S(\Omega) = S(\Omega_1) + S(\Omega_2)$$

Suppose the for each outcome i there is an associated probability

$$P_i \text{ for } 1 \leq i \leq \Omega; \text{ For a uniform system, we expect } P_i = \frac{1}{\Omega}.$$

$$\text{Argue that } S(\Omega; [P_i]) = -\sum_{i=1}^{\Omega} P_i \ln P_i.$$

2/6/2012

PHY 341/641 Spring 2012 -- Lecture 8

2

Bayes' theorem

Conditional probability --

Define $P(A|B) \equiv$ probability of A given B

Define $P(A|\bar{B}) \equiv$ probability of A given not B

$$\Rightarrow P(A) = P(A|B) + P(A|\bar{B})$$

$$\Rightarrow P(A \text{ and } B) = P(A|B)P(B) = P(B|A)P(A)$$

Bayes' theorem:

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

2/6/2012

PHY 341/641 Spring 2012 -- Lecture 8

3

Bayes' theorem continued:

Example: Suppose B represents a data set and A represents a model which can be generalized to several mutually exclusive models A_i

Generalized Bayes' theorem :

$$P(A_i | B) = \frac{P(B | A_i)P(A_i)}{P(B)}$$

If we assume that the set of A_i span all possible outcomes, then

$$P(B) = \sum_i P(B | A_i)P(A_i)$$

$$\Rightarrow P(A_i | B) = \frac{P(B | A_i)P(A_i)}{\sum_j P(B | A_j)P(A_j)}$$

2/6/2012

PHY 341/641 Spring 2012 -- Lecture 8

4

Example 3.16. Bayes' theorem and the problem of false positives

Even though you have no symptoms, your doctor wishes to test you for a rare disease that only 1 in 10,000 people of your age contract. The test is 98% accurate, which means that, if you have the disease, 98% of the times the test will come out positive and 2% negative. Also, if you do not have the disease, the test will come out negative 98% of the time and positive 2% of the time. You take the test and it comes out positive. What is the probability that you have the disease? Answer the question before you read the solution using Bayes' theorem.

Let $P(D)$ \equiv probability of having the disease (0.0001 in this case)

$P(\bar{D}) \equiv P(N)$ \equiv probability of not having disease (0.9999)

$P(+ | D) = 0.98$ $P(- | D) = 0.02$

$P(+ | N) = 0.02$ $P(- | N) = 0.98$

Want to know $P(D | +)$ \equiv probability of having disease given positive test

$$\begin{aligned} P(D | +) &= \frac{P(+ | D)P(D)}{P(+ | D)P(D) + P(+ | N)P(N)} \\ &= \frac{0.98 \cdot 0.0001}{0.98 \cdot 0.0001 + 0.02 \cdot 0.9999} = 0.0049 = 0.49\% \end{aligned}$$

Compare with $P(D) = 0.0001 = 0.01\%$

2/6/2012

PHY 341/641 Spring 2012 -- Lecture 8

5

Binomial distribution

Appropriate for describing systems with exactly two elemental outcomes such as

1. Coin Toss – HT -- $P_H = p$, $P_T = 1 - p = q$

2. Spin $\uparrow \downarrow$ -- $P_\uparrow = p$, $P_\downarrow = 1 - p = q$

Typically, we are interested in the number n of elemental outcomes of type 1 (H, \uparrow , etc.) in a total of N processes.

Example: $N=4$, $n=\#H$'s, $p=1/2=q$

Configurations	n	P(n)
TTTT	0	1/16
HTTT, THTT, TTHT, TTTH	1	4/16
TTHH, THTH, HHTT, THTH, HTHT, HTTH	2	6/16
THHH, HTHH, HHTH, HHHT	3	4/16
HHHH	4	1/16

2/6/2012

PHY 341/641 Spring 2012 -- Lecture 8

6

Binomial distribution

Example: $N=4$, $n=\#H's$, $p=1/2=q$

Configurations	n	P(n)
TTTT	0	1/16
HTTT, THTT, TTHT, TTTH	1	4/16
TTTH, THTT, HHTT, THTH, HTHT, HTTH	2	6/16
TTHH, HTTH, HHTH, HHHT	3	4/16
HHHH	4	1/16

Example: $N=3$, $n=\#H's$, $p=1/2=q$

Configurations	n	P(n)
TTT	0	1/8
HTT, THT, TTH	1	3/8
TTH, HTH, HHT	2	3/8
HHH	3	1/8

2/6/2012

PHY 341/641 Spring 2012 -- Lecture 8

7

Binomial distribution (continued)

In each of these cases,

$$P_N(n) = \frac{N!}{n!(N-n)!} p^n q^{N-n}$$

Note that

$$\sum_{n=0}^N P_N(n) = \sum_{n=0}^N \frac{N!}{n!(N-n)!} p^n q^{N-n} = 1$$

Because

$$1^N = (p+q)^N = \sum_{n=0}^N \frac{N!}{n!(N-n)!} p^n q^{N-n}$$

2/6/2012

PHY 341/641 Spring 2012 -- Lecture 8

8

Average value of n in binomial distribution

$$\langle n \rangle = \sum_{n=0}^N n P_N(n) = \sum_{n=0}^N n \frac{N!}{n!(N-n)!} p^n q^{N-n}$$

Note that : $n p^n q^{N-n} = p \left(\frac{\partial}{\partial p} \right)_q p^n q^{N-n}$

$$\langle n \rangle = p \left(\frac{\partial}{\partial p} \right)_q \sum_{n=0}^N \frac{N!}{n!(N-n)!} p^n q^{N-n} = p \left(\frac{\partial}{\partial p} \right)_q (p+q)^N = pN$$

2/6/2012

PHY 341/641 Spring 2012 -- Lecture 8

9

Similarly, we can show that :

$$\langle n^2 \rangle = \sum_{n=0}^N n^2 P_N(n) = \sum_{n=0}^N n^2 \frac{N!}{n!(N-n)!} p^n q^{N-n}$$

Note that : $n^2 p^n q^{N-n} = \left(p \frac{\partial}{\partial p} \right)_q^2 p^n q^{N-n}$

$$\langle n^2 \rangle = \left(p \frac{\partial}{\partial p} \right)_q^2 (p+q)^N = pN + p^2 N(N-1)$$

Variance :

$$\langle n^2 \rangle - \langle n \rangle^2 = pN + p^2 N(N-1) - p^2 N^2 = Np(1-p) = Npq$$

$$\langle n^2 \rangle - \langle n \rangle^2 \equiv \sigma^2 = Npq$$

2/6/2012

PHY 341/641 Spring 2012 -- Lecture 8


10

Binomial distribution in the limit $N \rightarrow \infty$

$$P_N(n) = \frac{N!}{n!(N-n)!} p^n q^{N-n}$$

$$P_{N \rightarrow \infty}(n) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(n-\langle n \rangle)^2 / 2\sigma^2}$$

Result from [stp_Binomial.jar](#)



2/6/2012

PHY 341/641 Spring 2012 -- Lecture 8

11

Stirling's approximation to factorial

For large m , Stirling's approximation to $m!$ is :

$$\ln m! \approx m \ln m - m + \frac{1}{2} \ln(2\pi m)$$

Evaluate Binomial distribution as Taylor's expansion of $\ln(P_N(n))$ about $\langle n \rangle$:

$$\ln P_N(n) \approx \ln P_N(\langle n \rangle) + (n - \langle n \rangle) \left. \frac{d \ln P_N(n)}{dn} \right|_{\langle n \rangle} + \frac{1}{2} (n - \langle n \rangle)^2 \left. \frac{d^2 \ln P_N(n)}{dn^2} \right|_{\langle n \rangle} \dots$$

\downarrow \downarrow \downarrow
 $\ln A$ 0 $-B$

$$\Rightarrow P_N(n) \approx A e^{-\frac{B}{2}(n-\langle n \rangle)^2}$$

2/6/2012

PHY 341/641 Spring 2012 -- Lecture 8

12

$$A \approx \frac{1}{\sqrt{2\pi\sigma^2}} \quad B \approx \frac{1}{\sigma^2}$$

Summary :

$$P_N(n) = \frac{N!}{n!(N-n)!} p^n q^{N-n}$$

$$P_\infty(n) \approx \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(n-\langle n \rangle)^2}{2\sigma^2}}$$

2/6/2012

PHY 341/641 Spring 2012 -- Lecture 8

13

Poisson probability distribution –

Approximation to binomial distribution for small p

$$P_N(n) = \frac{N!}{n!(N-n)!} p^n q^{N-n}$$

Assume $N \gg n$ $p \ll 1$

$$\frac{N!}{n!(N-n)!} \approx \frac{N^n}{n!}$$

$$\ln q^{N-n} = (N-n) \ln(1-p) \approx -(N-n)p \approx -Np$$

$$P_N(n) \approx \frac{N^n}{n!} p^n e^{-Np} = \frac{\langle n \rangle^n}{n!} e^{-\langle n \rangle}$$

2/6/2012

PHY 341/641 Spring 2012 -- Lecture 8

14
