

PHY 712 Electrodynamics

11-11:50 AM MWF Olin 107

Plan for Lecture 28:

Start reading Chap. 14 – Radiation by moving charges

- A. Comments on Exam 2**
- B. Radiation due to moving charges**
 - 1. Motion in a line**
 - 2. Motion in a circle**

	03-22(Fri)	<i>APS Meeting</i>	(no class)	Exam
25	03-25(Mon)	Chap. 11	Lorentz transformations	#17
26	03-27(Wed)	Chap. 11	Transformations between electromagnetic fields	#18
27	03-28(Thur)	Chap. 11	Liénard-Wiechert potentials revisited	
	03-29(Fri)	<i>Good Friday</i>	(no class)	
28	04-01(Mon)	Chap. 14	Radiation by accelerated charges	#19
29	04-03(Wed)	Chap. 14	Radiation by accelerated charges	
30	04-05(Fri)	Chap. 14	Synchrotron radiation spectrum	
31	04-08(Mon)			
32	04-10(Wed)			
33	04-12(Fri)			
	04-15(Mon)		(no class -- presentation preparation)	
	04-17(Wed)		(no class -- presentation preparation)	
	04-19(Fri)		(no class -- presentation preparation)	
34	04-22(Mon)			
35	04-24(Wed)			
36	04-26(Fri)			
	04-29(Mon)		Student presentations I	
	05-01(Wed)		Student presentations II	
	05-02(Thurs)		Student presentations III	


Comments on Exam 2

Radiation from a moving charged particle

Variables (notation) :

$$\dot{\mathbf{R}}_q(t_r) \equiv \frac{d\mathbf{R}_q(t_r)}{dt_r} \equiv \mathbf{v}$$

$$\mathbf{R}(t_r) \equiv \mathbf{r} - \mathbf{R}_q(t_r) \equiv \mathbf{R}$$



Liénard-Wiechert fields (cgs Gaussian units):

$$\mathbf{E}(\mathbf{r}, t) = \frac{q}{\left(R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}\right)^3} \left[\left(\mathbf{R} - \frac{\mathbf{v}R}{c} \right) \left(1 - \frac{v^2}{c^2} \right) + \left(\mathbf{R} \times \left\{ \left(\mathbf{R} - \frac{\mathbf{v}R}{c} \right) \times \frac{\dot{\mathbf{v}}}{c^2} \right\} \right) \right]. \quad (19)$$

$$\mathbf{B}(\mathbf{r}, t) = \frac{q}{c} \left[\frac{-\mathbf{R} \times \mathbf{v}}{\left(R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}\right)^3} \left(1 - \frac{v^2}{c^2} + \frac{\dot{\mathbf{v}} \cdot \mathbf{R}}{c^2} \right) - \frac{\mathbf{R} \times \dot{\mathbf{v}}/c}{\left(R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}\right)^2} \right]. \quad (20)$$

In this case, the electric and magnetic fields are related according to

$$\mathbf{B}(\mathbf{r}, t) = \frac{\mathbf{R} \times \mathbf{E}(\mathbf{r}, t)}{R}. \quad (21)$$

$$\dot{\mathbf{R}}_q(t_r) \equiv \frac{d\mathbf{R}_q(t_r)}{dt_r} \equiv \mathbf{v} \quad \mathbf{R}(t_r) \equiv \mathbf{r} - \mathbf{R}_q(t_r) \equiv \mathbf{R} \quad \dot{\mathbf{v}} \equiv \frac{d^2\mathbf{R}_q(t_r)}{dt_r^2}$$

Electric field far from source:

$$\mathbf{E}(\mathbf{r}, t) = \frac{q}{\left(R - \frac{\mathbf{v} \cdot \mathbf{R}}{c} \right)^3} \left\{ \mathbf{R} \times \left[\left(\mathbf{R} - \frac{\mathbf{v}R}{c} \right) \times \frac{\dot{\mathbf{v}}}{c^2} \right] \right\}$$

$$\mathbf{B}(\mathbf{r}, t) = \frac{\mathbf{R} \times \mathbf{E}(\mathbf{r}, t)}{R}$$

Let $\hat{\mathbf{R}} \equiv \frac{\mathbf{R}}{R}$ $\boldsymbol{\beta} \equiv \frac{\mathbf{v}}{c}$ $\dot{\boldsymbol{\beta}} \equiv \frac{\dot{\mathbf{v}}}{c}$

$$\mathbf{E}(\mathbf{r}, t) = \frac{q}{cR \left(1 - \boldsymbol{\beta} \cdot \hat{\mathbf{R}} \right)^3} \left\{ \hat{\mathbf{R}} \times \left[\left(\hat{\mathbf{R}} - \boldsymbol{\beta} \right) \times \dot{\boldsymbol{\beta}} \right] \right\}$$

$$\mathbf{B}(\mathbf{r}, t) = \hat{\mathbf{R}} \times \mathbf{E}(\mathbf{r}, t)$$

Poynting vector:

$$\mathbf{S}(\mathbf{r}, t) = \frac{c}{4\pi} (\mathbf{E} \times \mathbf{B})$$

$$\mathbf{E}(\mathbf{r}, t) = \frac{q}{cR(1 - \boldsymbol{\beta} \cdot \hat{\mathbf{R}})^3} \left\{ \hat{\mathbf{R}} \times [(\hat{\mathbf{R}} - \boldsymbol{\beta}) \times \dot{\boldsymbol{\beta}}] \right\}$$

$$\mathbf{B}(\mathbf{r}, t) = \hat{\mathbf{R}} \times \mathbf{E}(\mathbf{r}, t)$$

$$\mathbf{S}(\mathbf{r}, t) = \frac{c}{4\pi} \hat{\mathbf{R}} |\mathbf{E}(\mathbf{r}, t)|^2 = \frac{q^2}{4\pi c R^2} \hat{\mathbf{R}} \frac{|\hat{\mathbf{R}} \times [(\hat{\mathbf{R}} - \boldsymbol{\beta}) \times \dot{\boldsymbol{\beta}}]|^2}{(1 - \boldsymbol{\beta} \cdot \hat{\mathbf{R}})^6}$$

Power radiated

$$\mathbf{S}(\mathbf{r}, t) = \frac{c}{4\pi} \hat{\mathbf{R}} |\mathbf{E}(\mathbf{r}, t)|^2 = \frac{q^2}{4\pi c R^2} \hat{\mathbf{R}} \frac{|\hat{\mathbf{R}} \times [(\hat{\mathbf{R}} - \beta) \times \dot{\beta}]|^2}{(1 - \beta \cdot \hat{\mathbf{R}})^6}$$

$$\frac{dP}{d\Omega} = \mathbf{S} \cdot \hat{\mathbf{R}} R^2 = \frac{q^2}{4\pi c} \frac{|\hat{\mathbf{R}} \times [(\hat{\mathbf{R}} - \beta) \times \dot{\beta}]|^2}{(1 - \beta \cdot \hat{\mathbf{R}})^6}$$

In the non - relativistic limit : $\beta \ll 1$


$$\frac{dP}{d\Omega} = \frac{q^2}{4\pi c} |\hat{\mathbf{R}} \times [\hat{\mathbf{R}} \times \dot{\beta}]|^2 = \frac{q^2}{4\pi c^3} |\dot{\mathbf{v}}|^2 \sin^2 \Theta$$

Radiation from a moving charged particle

Variables (notation) :

$$\dot{\mathbf{R}}_q(t_r) \equiv \frac{d\mathbf{R}_q(t_r)}{dt_r} \equiv \mathbf{v}$$

$$\mathbf{R}(t_r) \equiv \mathbf{r} - \mathbf{R}_q(t_r) \equiv \mathbf{R}$$



$$\frac{dP}{d\Omega} = \frac{q^2}{4\pi c^3} |\dot{\mathbf{v}}|^2 \sin^2 \Theta$$

Radiation power in non-relativistic case -- continued

$$\frac{dP}{d\Omega} = \frac{q^2}{4\pi c^3} |\dot{\mathbf{v}}|^2 \sin^2 \Theta$$

$$P = \int d\Omega \frac{dP}{d\Omega} = \frac{2}{3} \frac{q^2}{c^3} |\dot{\mathbf{v}}|^2$$

Radiation distribution in the relativistic case

$$\frac{dP}{d\Omega} = \mathbf{S} \cdot \hat{\mathbf{R}} R^2 = \frac{q^2}{4\pi c} \left| \frac{\hat{\mathbf{R}} \times [(\hat{\mathbf{R}} - \boldsymbol{\beta}) \times \dot{\boldsymbol{\beta}}]^2}{(1 - \boldsymbol{\beta} \cdot \hat{\mathbf{R}})^6} \right|_{t_r=t-R/c}$$

This expression gives us the energy per unit field time t . We are often interested in the power per unit retarded time $t_r = t - R/c$:

$$\frac{dP(t)}{d\Omega} = \frac{dP_r(t_r)}{d\Omega} \frac{dt}{dt_r} \quad \frac{dt}{dt_r} = 1 - \boldsymbol{\beta} \cdot \hat{\mathbf{R}}$$

$$\frac{dP_r(t_r)}{d\Omega} = \frac{q^2}{4\pi c} \left| \frac{\hat{\mathbf{R}} \times [(\hat{\mathbf{R}} - \boldsymbol{\beta}) \times \dot{\boldsymbol{\beta}}]^2}{(1 - \boldsymbol{\beta} \cdot \hat{\mathbf{R}})^5} \right|_{t_r=t-R/c}$$

Radiation distribution in the relativistic case -- continued

$$\frac{dP_r(t_r)}{d\Omega} = \frac{q^2}{4\pi c} \left| \hat{\mathbf{R}} \times \left[(\hat{\mathbf{R}} - \beta) \times \dot{\beta} \right]^2 \right| \Bigg|_{t_r=t-R/c}$$

For linear acceleration: $\beta \times \dot{\beta} = 0$

$$\frac{dP_r(t_r)}{d\Omega} = \frac{q^2}{4\pi c} \left| \hat{\mathbf{R}} \times (\hat{\mathbf{R}} \times \dot{\beta})^2 \right| \Bigg|_{t_r=t-R/c} = \frac{q^2}{4\pi c^3} |\dot{\mathbf{v}}|^2 \frac{\sin^2 \theta}{(1 - \beta \cos \theta)^5}$$

Power from linearly accelerating particle

$$\frac{dP_r(t_r)}{d\Omega} = \frac{q^2}{4\pi c} \frac{\left| \hat{\mathbf{R}} \times (\hat{\mathbf{R}} \times \dot{\beta}) \right|^2}{(1 - \beta \cdot \hat{\mathbf{R}})^5} \Bigg|_{t_r = t - R/c} = \frac{q^2}{4\pi c^3} |\dot{\mathbf{v}}|^2 \frac{\sin^2 \theta}{(1 - \beta \cos \theta)^5}$$

