

PHY 712 Electrodynamics
10-10:50 AM MWF Olin 107

Plan for Lecture 10:

Complete reading of Chapter 4

A. Microscopic \leftrightarrow macroscopic polarizability

B. Clausius-Mossotti equation

C. Electrostatic energy in dielectric media

02/07/2014 PHY 712 Spring 2014 -- Lecture 10 1

Course schedule for Spring 2014

(Preliminary schedule -- subject to frequent adjustment.) Please note that makeup lectures (indicated in red) are scheduled for Tuesdays or Thursdays at 11 AM - 12:15 PM in Olin 107.

Lecture date	JDJ Reading	Topic	Assign.	Due date
1 Wed 01/15/2014	Chap. 1	Introduction, units and Poisson equation	#1	01/31/2014
2 Thu 01/16/2014	Chap. 1	Electrostatic energy calculations	#2	01/31/2014
3 Fri 01/17/2014	Chap. 1	Poisson equation and Green's theorem	#3	01/31/2014
Mon 01/20/2014		MLK Holiday - no class		
4 Wed 01/22/2014	Chap. 1	Green's functions for cartesian coordinates	#4	01/31/2014
5 Thu 01/23/2014	Chap. 1	Brief introduction to numerical methods	#5	01/31/2014
6 Fri 01/24/2014	Chap. 2	Method of images	#6	01/31/2014
Mon 01/27/2014		NAWH out of town - no class		
Wed 01/29/2014		NAWH out of town - no class		
7 Fri 01/31/2014	Chap. 3	Cylindrical and spherical geometries	#7	02/05/2014
8 Mon 02/03/2014	Chap. 4	Multipole analysis of charge distributions	#8	02/05/2014
9 Wed 02/05/2014	Chap. 4	Dipoles and dielectrics	#9	02/07/2014
10 Fri 02/07/2014	Chap. 4	Dipoles and dielectrics	#10	02/10/2014
11 Mon 02/10/2014	Chap. 5	Magnetostatics	#11	02/12/2014

02/07/2014 PHY 712 Spring 2014 -- Lecture 10 2

Focus on dipolar fields:

Dipole moment \mathbf{p} :

$$\mathbf{p} \equiv \int d^3r' \mathbf{r}' \rho(\mathbf{r}')$$

For r outside the extent of $\rho(\mathbf{r})$:

Electrostatic potential from single dipole :

$$\Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \left(\frac{\mathbf{p} \cdot \mathbf{r}}{r^3} \right)$$

Electrostatic field from single dipole :


$$\mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \left(\frac{3\mathbf{r}(\mathbf{p} \cdot \mathbf{r}) - r^2\mathbf{p}}{r^5} - \frac{4\pi}{3} \mathbf{p} \delta^3(\mathbf{r}) \right)$$

02/07/2014 PHY 712 Spring 2014 -- Lecture 10 3

Microscopic origin of dipole moments

- Polarizable isotropic atoms/molecules
- Charge anisotropic molecules

Polarizable isotropic atoms/molecules




At equilibrium:

$$qE - m\omega_0^2 \delta x = 0$$

$$\delta x = \frac{qE}{m\omega_0^2}$$

02/07/2014 PHY 712 Spring 2014 – Lecture 10 4

Polarizable isotropic atoms/molecules – continued:



At equilibrium:

$$qE - m\omega_0^2 \delta x = 0$$

$$\delta x = \frac{qE}{m\omega_0^2}$$

Induced dipole moment:

$$p = q\delta x = \frac{q^2}{m\omega_0^2} E \equiv \epsilon_0 \gamma_{mol} E$$

02/07/2014 PHY 712 Spring 2014 – Lecture 10 5

Alignment of molecules with permanent dipoles p_0 :




For a freely rotating dipole its average moment in an electric field, estimated assuming a Boltzmann distribution:

$$\langle p_{mol} \rangle = \frac{E \int d\Omega p_0 \cos \theta e^{-p_0 E \cos \theta / kT}}{\int d\Omega e^{-p_0 E \cos \theta / kT}}$$

$$= \frac{1}{3} \frac{p_0^2}{kT} E \equiv \epsilon_0 \gamma_{mol} E$$

02/07/2014 PHY 712 Spring 2014 – Lecture 10 6

Field due to collection of induced dipoles



$$\mathbf{E}_{tot} = \sum_i \mathbf{E}_i^0 + \mathbf{E}_{ext}$$

$$\mathbf{E}_{site(i)} = \sum_{j \neq i} \mathbf{E}_j^0 + \mathbf{E}_{ext}$$


$$= \mathbf{E}_{tot} - \mathbf{E}_i^0$$

Electrostatic field from single dipole :

$$\mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \left(\frac{3\mathbf{r}(\mathbf{p} \cdot \mathbf{r}) - r^2\mathbf{p}}{r^5} - \frac{4\pi}{3} \mathbf{p} \delta^3(\mathbf{r}) \right)$$

02/07/2014 PHY 712 Spring 2014 -- Lecture 10 7

Field due to collection of induced dipoles -- continued



$$\mathbf{E}_{tot} = \sum_i \mathbf{E}_i^0 + \mathbf{E}_{ext}$$

$$\mathbf{E}_{site(i)} = \sum_{j \neq i} \mathbf{E}_j^0 + \mathbf{E}_{ext}$$

$$= \mathbf{E}_{tot} - \mathbf{E}_i^0$$


$$\mathbf{E}(\mathbf{r})_{tot} = \frac{1}{4\pi\epsilon_0} \sum_i \left(\frac{3\mathbf{r}(\mathbf{p}_i \cdot \mathbf{r}) - r^2\mathbf{p}_i}{r^5} - \frac{4\pi}{3} \mathbf{p}_i \delta^3(\mathbf{r}) \right) + \mathbf{E}_{ext}$$

$$\mathbf{E}(\mathbf{r})_{site(i)} = \frac{1}{4\pi\epsilon_0} \left(\sum_{j \neq i} \frac{3\mathbf{r}(\mathbf{p}_j \cdot \mathbf{r}) - r^2\mathbf{p}_j}{r^5} \right) + \mathbf{E}_{ext}$$

$$\langle \mathbf{E}_{site(i)} \rangle = \langle \mathbf{E}_{tot} \rangle + \frac{1}{V} \frac{1}{3\epsilon_0} \langle \mathbf{P} \rangle = \langle \mathbf{E}_{tot} \rangle + \frac{1}{3\epsilon_0} \langle \mathbf{P} \rangle$$

02/07/2014 PHY 712 Spring 2014 -- Lecture 10 8

Field due to collection of induced dipoles -- continued



$$\langle \mathbf{E}_{site(i)} \rangle = \langle \mathbf{E}_{tot} \rangle + \frac{1}{3\epsilon_0} \langle \mathbf{P} \rangle$$

$$\langle \mathbf{p} \rangle = \epsilon_0 \gamma_{mol} \langle \mathbf{E}_{site} \rangle$$

$$\langle \mathbf{P} \rangle = \frac{1}{V} \langle \mathbf{p} \rangle = \frac{\epsilon_0 \gamma_{mol}}{V} \left(\langle \mathbf{E}_{tot} \rangle + \frac{1}{3\epsilon_0} \langle \mathbf{P} \rangle \right)$$

$$\langle \mathbf{P} \rangle = \frac{\epsilon_0 \gamma_{mol}}{V} \frac{\langle \mathbf{E}_{tot} \rangle}{1 - \frac{\gamma_{mol}}{V}} = \epsilon_0 \chi_e \langle \mathbf{E}_{tot} \rangle$$

Claussius-Mossotti equation

$$\chi_e = \frac{\frac{\gamma_{mol}}{V}}{1 - \frac{\gamma_{mol}}{3V}} \quad \gamma_{mol} = 3V \left(\frac{\epsilon / \epsilon_0 - 1}{\epsilon / \epsilon_0 + 2} \right)$$

02/07/2014 PHY 712 Spring 2014 -- Lecture 10 9

Example of the Clausius-Mossotti equation –
 Pentane (C₅H₁₂) at various densities

Density (g/cm ³)	Mol/m ³	ϵ/ϵ_0	$3V^*(\epsilon/\epsilon_0 - 1)/(\epsilon/\epsilon_0 + 2)$
0.613	5.12536E+27	1.82	1.25646E-28
0.701	5.86114E+27	1.96	1.24084E-28
0.796	6.65544E+27	2.12	1.22536E-28
0.865	7.23236E+27	2.24	1.2131E-28
0.907	7.58353E+27	2.33	1.2151E-28

02/07/2014 PHY 712 Spring 2014 – Lecture 10 10

Re-examination of electrostatic energy in dielectric media

$$W = \frac{1}{2} \int d^3r \rho_{mono}(\mathbf{r}) \Phi(\mathbf{r})$$

In terms of displacement field:

$$\nabla \cdot \mathbf{D} = \rho_{mono}(\mathbf{r})$$

$$W = \frac{1}{2} \int d^3r \nabla \cdot \mathbf{D} \Phi(\mathbf{r}) = \frac{1}{2} \int d^3r \nabla \cdot (\mathbf{D}(\mathbf{r}) \Phi(\mathbf{r})) - \frac{1}{2} \int d^3r \mathbf{D}(\mathbf{r}) \cdot \nabla \Phi(\mathbf{r})$$

$$= 0 + \frac{1}{2} \int d^3r \mathbf{D}(\mathbf{r}) \cdot \mathbf{E}(\mathbf{r})$$

$$W = \frac{1}{2} \int d^3r \mathbf{D}(\mathbf{r}) \cdot \mathbf{E}(\mathbf{r})$$

02/07/2014 PHY 712 Spring 2014 – Lecture 10 11

Comment on the “Modern Theory of Polarization”
 Some references:

- R. D.King-Smith and D. Vanderbilt, Phys. Rev. B 47, 1651 (1993)
- R. Resta, Rev. Mod. Physics 66, 699 (1994)
- R, Resta, J. Phys. Condens. Matter 23, 123201 (2010)

Basic equations :

$$\epsilon_0 \nabla \cdot \mathbf{E} = \rho_{tot} = \rho_{bound} + \rho_{mono}$$

$$\nabla \cdot \mathbf{P} = \rho_{bound}$$

$$\nabla \cdot \mathbf{D} = \rho_{mono}$$

$$\epsilon_0 \mathbf{E} = \mathbf{D} + \mathbf{P}$$

Note: In general \mathbf{P} is highly dependent on the boundary values; often it is more convenient/meaningful to calculate $\Delta \mathbf{P}$.

02/07/2014 PHY 712 Spring 2014 – Lecture 10 12

Comment on the "Modern Theory of Polarization"
-- continued

$$\nabla \cdot \Delta \mathbf{P} = \Delta \rho_{\text{bound}} = \Delta \rho_{\text{bound}}^{\text{nuclear}} + \Delta \rho_{\text{bound}}^{\text{electronic}}$$
$$\Delta \mathbf{P}^{\text{electronic}} = -\frac{e}{V_{\text{crystal}}} \sum_n \langle w_{n0} | \mathbf{r} | w_{n0} \rangle$$

02/07/2014

PHY 712 Spring 2014 -- Lecture 10

13
