

PHY 712 Electrodynamics
10-10:50 AM MWF Olin 107
(Make up lecture 2/17/2014 at 9 AM)

Plan for Lecture 13:

Continue reading Chapter 5

1. Hyperfine interaction
 2. Macroscopic magnetization density M
 3. H field and its relation to B
 4. Magnetic boundary values

02/17/2014

PHY 712 Spring 2014 – Lecture 13

1

02/17/2014

PHY 712 Spring 2014 – Lecture 13

2

Hyperfine interaction energy:

Interactions between magnetic dipoles

Sources of magnetic dipoles and other sources of magnetism in an atom:

- Intrinsic magnetic moment of a nucleus μ_N
 - Intrinsic magnetic moment of an electron μ_e
 - Magnetic field due to electron current $\mathbf{J}(\mathbf{r})$

Interaction energy between a magnetic dipole \mathbf{m} and a magnetic field \mathbf{B} : $E_{int} = -\mathbf{m} \cdot \mathbf{B}$

$$\mu_N$$

In this case: $E_{int} = -\mu_a \cdot \mathbf{B}_a - \mu_N \cdot \mathbf{B}_N(0)$

$$\mathbf{B}_{\mu_N}(\mathbf{r}) = \frac{\mu_0}{4\pi} \left\{ \frac{3\hat{\mathbf{r}}(\boldsymbol{\mu}_N \cdot \hat{\mathbf{r}}) - \boldsymbol{\mu}_N}{r^3} + \frac{8\pi}{3} \boldsymbol{\mu}_N \delta^3(\mathbf{r}) \right\}$$

02/17/2014

PHY 712 Spring 2014 – Lecture 13

3

Hyperfine interaction energy: -- continued

$$E_{int} = -\mu_e \cdot \mathbf{B}_{\mu_N} - \mu_N \cdot \mathbf{B}_{J_e}(0)$$

Evaluation of the magnetic field at the nucleus due to the electron current density:
The vector potential associated with an electron in a bound state of an atom as described by a quantum mechanical wavefunction $\psi_{nlm_l}(\mathbf{r})$ can be written:

$$\mathbf{A}_{J_e}(\mathbf{r}) = -\frac{\mu_0 e \hbar}{4\pi m_e} m_l \int d^3 r' \frac{\hat{\mathbf{z}} \times \mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|} \frac{|\psi_{nlm_l}(\mathbf{r}')|^2}{r'^2 \sin^2 \theta'}$$

We want to evaluate the magnetic field $\mathbf{B} = \nabla \times \mathbf{A}$ in the vicinity of the nucleus ($\mathbf{r} \rightarrow 0$).

02/17/2014

PHY 712 Spring 2014 -- Lecture 13

4

Hyperfine interaction energy: -- continued

$$\begin{aligned} \mathbf{B}_{J_e}(0) &= \nabla \times \mathbf{A}_{J_e} \Big|_{\mathbf{r} \rightarrow 0} = -\frac{\mu_0 e \hbar}{4\pi m_e} m_l \int d^3 r' \nabla \times \frac{\hat{\mathbf{z}} \times \mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|} \frac{|\psi_{nlm_l}(\mathbf{r}')|^2}{r'^2 \sin^2 \theta'} \Big|_{\mathbf{r} \rightarrow 0} \\ \mathbf{B}_o(\mathbf{r}) &= \frac{\mu_0 e \hbar}{4\pi m_e} m_l \int d^3 r' \frac{(\mathbf{r} - \mathbf{r}') \times (\hat{\mathbf{z}} \times \mathbf{r}') |\psi_{nlm_l}(\mathbf{r}')|^2}{|\mathbf{r} - \mathbf{r}'|^3} \Big|_{\mathbf{r} \rightarrow 0} \\ \mathbf{B}_o(0) &= -\frac{\mu_0 e \hbar}{4\pi m_e} m_l \int d^3 r' \frac{\mathbf{r}' \times (\hat{\mathbf{z}} \times \mathbf{r}') |\psi_{nlm_l}(\mathbf{r}')|^2}{r'^3} \Big|_{\mathbf{r} \rightarrow 0} \\ \hat{\mathbf{r}}' \times (\hat{\mathbf{z}} \times \hat{\mathbf{r}}') &= \hat{\mathbf{z}}(1 - \cos^2 \theta') - \hat{\mathbf{x}} \cos \theta' \sin \theta' \cos \phi' - \hat{\mathbf{y}} \cos \theta' \sin \theta' \sin \phi'. \\ \mathbf{B}_o(0) &= -\frac{\mu_0 e \hbar}{4\pi m_e} m_l \int d^3 r' \frac{\hat{\mathbf{z}} r'^2 \sin^2 \theta' |\psi_{nlm_l}(\mathbf{r}')|^2}{r'^3} = -\frac{\mu_0 e \hbar}{4\pi m_e} m_l \hat{\mathbf{z}} \int d^3 r' \frac{|\psi_{nlm_l}(\mathbf{r}')|^2}{r'^3} \\ &= -\frac{\mu_0 e \hbar}{4\pi m_e} m_l \hat{\mathbf{z}} \left\langle \frac{1}{r'^3} \right\rangle \end{aligned}$$

02/17/2014

PHY 712 Spring 2014 -- Lecture 13

5

Hyperfine interaction energy: -- continued

$$E_{int} \equiv H_{HF} = -\mu_e \cdot \mathbf{B}_{\mu_N} - \mu_N \cdot \mathbf{B}_{J_e}(0)$$

Putting all of the terms together:

$$H_{HF} = -\frac{\mu_0}{4\pi} \left(\left\langle \frac{3(\mu_N \cdot \hat{\mathbf{r}})(\mu_e \cdot \hat{\mathbf{r}}) - \mu_N \cdot \mu_e}{r^3} + \frac{8\pi}{3} \mu_N \cdot \mu_e \delta^3(\mathbf{r}) \right\rangle + \frac{e}{m_e} \left\langle \frac{\mathbf{L} \cdot \mu_N}{r^3} \right\rangle \right).$$

In this expression the brackets $\langle \rangle$ indicate evaluating the expectation value relative to the electronic state.

02/17/2014

PHY 712 Spring 2014 -- Lecture 13

6

Macroscopic dipolar effects --
Magnetic dipole moment

$$\mathbf{m} = \frac{1}{2} \int d^3r \mathbf{r} \times \mathbf{J}(\mathbf{r})$$

Note that the intrinsic spin of elementary particles is associated with a magnetic dipole moment, but we often do not have a detailed knowledge of $\mathbf{J}(\mathbf{r})$.

Vector potential for magnetic dipole moment

$$\mathbf{A}(\mathbf{r}) = \frac{\mu_0}{4\pi} \frac{\mathbf{m} \times \mathbf{r}}{|\mathbf{r}|^3}$$

02/17/2014

PHY 712 Spring 2014 -- Lecture 13

7

Macroscopic magnetization

$$\mathbf{M}(\mathbf{r}) = \sum_i \mathbf{m}_i \delta^3(\mathbf{r} - \mathbf{r}_i)$$

Vector potential due to "free" current $\mathbf{J}_{\text{free}}(\mathbf{r})$ and macroscopic magnetization $\mathbf{M}(\mathbf{r})$. Note: the designation $\mathbf{J}_{\text{free}}(\mathbf{r})$ implies that this current does not also contribute to the magnetization density.

$$\mathbf{A}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int d^3r' \left(\frac{\mathbf{J}_{\text{free}}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} + \frac{\mathbf{M}(\mathbf{r}') \times (\mathbf{r} - \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|^3} \right)$$

02/17/2014

PHY 712 Spring 2014 -- Lecture 13

8

Vector potential contributions from macroscopic magnetization -- continued

$$\mathbf{A}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int d^3r' \left(\frac{\mathbf{J}_{\text{free}}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} + \frac{\mathbf{M}(\mathbf{r}') \times (\mathbf{r} - \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|^3} \right)$$

Note that :

$$\begin{aligned} \frac{\mathbf{M}(\mathbf{r}') \times (\mathbf{r} - \mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|^3} &= \mathbf{M}(\mathbf{r}') \times \nabla' \frac{1}{|\mathbf{r} - \mathbf{r}'|} \\ &= -\nabla' \times \left(\frac{\mathbf{M}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} \right) + \frac{\nabla' \times \mathbf{M}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} \\ \Rightarrow \mathbf{A}(\mathbf{r}) &= \frac{\mu_0}{4\pi} \int d^3r' \frac{\mathbf{J}_{\text{free}}(\mathbf{r}') + \nabla' \times \mathbf{M}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} \end{aligned}$$

02/17/2014

PHY 712 Spring 2014 -- Lecture 13

9

Vector potential contributions from macroscopic magnetization -- continued

$$\mathbf{A}(\mathbf{r}) = \frac{\mu_0}{4\pi} \int d^3 r' \frac{\mathbf{J}_{free}(\mathbf{r}') + \nabla' \times \mathbf{M}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}$$

Note that for the case that $\nabla \cdot \mathbf{A} = 0$:

$$\begin{aligned} \nabla \times \mathbf{B}(\mathbf{r}) &= \nabla \times (\nabla \times \mathbf{A}(\mathbf{r})) = -\nabla^2 \mathbf{A}(\mathbf{r}) \\ &= \frac{\mu_0}{4\pi} \int d^3 r' (4\pi \delta^3(\mathbf{r} - \mathbf{r}')) (\mathbf{J}_{free}(\mathbf{r}') + \nabla' \times \mathbf{M}(\mathbf{r}')) \\ &= \mu_0 (\mathbf{J}_{free}(\mathbf{r}) + \nabla \times \mathbf{M}(\mathbf{r})) \\ \Rightarrow \nabla \times (\mathbf{B}(\mathbf{r}) - \mu_0 \mathbf{M}(\mathbf{r})) &= \mu_0 \mathbf{J}_{free}(\mathbf{r}) \end{aligned}$$

02/17/2014

PHY 712 Spring 2014 -- Lecture 13

10

Magnetic field contributions

$$\nabla \times (\mathbf{B}(\mathbf{r}) - \mu_0 \mathbf{M}(\mathbf{r})) = \mu_0 \mathbf{J}_{free}(\mathbf{r})$$

Define the magnetic flux density :

$$\begin{aligned} \mu_0 \mathbf{H}(\mathbf{r}) &\equiv \mathbf{B}(\mathbf{r}) - \mu_0 \mathbf{M}(\mathbf{r}) \\ \Rightarrow \nabla \times \mathbf{H}(\mathbf{r}) &= \mathbf{J}_{free}(\mathbf{r}) \end{aligned}$$

$$\nabla \times (\mathbf{B}(\mathbf{r}) - \mu_0 \mathbf{M}(\mathbf{r})) = \mu_0 \mathbf{J}_{free}(\mathbf{r})$$

Note that $\mathbf{B}(\mathbf{r}) \equiv$ the magnetic flux density

Define $\mathbf{H}(\mathbf{r}) \equiv$ the magnetic field

$$\begin{aligned} \mu_0 \mathbf{H}(\mathbf{r}) &\equiv \mathbf{B}(\mathbf{r}) - \mu_0 \mathbf{M}(\mathbf{r}) \\ \Rightarrow \nabla \times \mathbf{H}(\mathbf{r}) &= \mathbf{J}_{free}(\mathbf{r}) \end{aligned}$$

02/17/2014

PHY 712 Spring 2014 -- Lecture 13

11

Summary of equations of magnetostatics :

$$\nabla \times \mathbf{B}(\mathbf{r}) = \mu_0 \mathbf{J}_{total}(\mathbf{r})$$

$$\nabla \times \mathbf{H}(\mathbf{r}) = \mathbf{J}_{free}(\mathbf{r})$$

$$\mathbf{B}(\mathbf{r}) = \mu_0 (\mathbf{H}(\mathbf{r}) + \mathbf{M}(\mathbf{r}))$$

$$\nabla \cdot \mathbf{B}(\mathbf{r}) = 0$$

For the case that $\mathbf{J}_{free}(\mathbf{r}) = 0$:



$$\nabla \times \mathbf{H}(\mathbf{r}) = 0$$

$$\nabla \cdot \mathbf{B}(\mathbf{r}) = 0$$

02/17/2014

PHY 712 Spring 2014 -- Lecture 13

12

For the case that $\mathbf{J}_{free}(\mathbf{r})$:

$$\nabla \times \mathbf{H}(\mathbf{r}) = 0$$

$$\nabla \cdot \mathbf{B}(\mathbf{r}) = 0$$

At boundary:

$$\mathbf{H}_1 \times \hat{\mathbf{n}} = \mathbf{H}_2 \times \hat{\mathbf{n}}$$

$$\mathbf{B}_1 \cdot \hat{\mathbf{n}} = \mathbf{B}_2 \cdot \hat{\mathbf{n}}$$



02/17/2014

PHY 712 Spring 2014 – Lecture 13

13

Example magnetostatic boundary value problem

M₀

$$\mathbf{M}(\mathbf{r}) = \begin{cases} M_0 \hat{\mathbf{z}} & r \leq a \\ 0 & r > a \end{cases}$$

$$\nabla \times \mathbf{H}(\mathbf{r}) = 0 \quad \Rightarrow \mathbf{H}(\mathbf{r}) = -\nabla \Phi_H(\mathbf{r})$$

$$\mathbf{B}(\mathbf{r}) = \mu_0 (\mathbf{H}(\mathbf{r}) + \mathbf{M}(\mathbf{r}))$$

$$\nabla \cdot \mathbf{B}(\mathbf{r}) = 0 = \mu_0 \nabla \cdot (\mathbf{H}(\mathbf{r}) + \mathbf{M}(\mathbf{r}))$$

$$\Rightarrow \nabla^2 \Phi_H(\mathbf{r}) = \nabla \cdot \mathbf{M}(\mathbf{r})$$

02/17/2014

PHY 712 Spring 2014 – Lecture 13

14

Example magnetostatic boundary value problem -- continued

M₀

$$\mathbf{M}(\mathbf{r}) = \begin{cases} M_0 \hat{\mathbf{z}} & r \leq a \\ 0 & r > a \end{cases}$$

$$\nabla^2 \Phi_H(\mathbf{r}) = \nabla \cdot \mathbf{M}(\mathbf{r})$$

$$\Rightarrow \Phi_H(\mathbf{r}) = -\frac{1}{4\pi} \int d^3 r' \frac{\nabla' \cdot \mathbf{M}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}$$

$$= -\frac{1}{4\pi} \int d^3r' \left[\nabla' \cdot \left(\frac{\mathbf{M}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} \right) - \mathbf{M}(\mathbf{r}') \cdot \nabla' \frac{1}{|\mathbf{r} - \mathbf{r}'|} \right]$$

$$= -\frac{1}{4\pi} \nabla \cdot \int d^3r' \frac{\mathbf{M}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}$$

02/17/2014

PHY 712 Spring 2014 – Lecture 13

15

Example magnetostatic boundary value problem -- continued

$$\mathbf{M}(\mathbf{r}) = \begin{cases} M_0 \hat{\mathbf{z}} & r \leq a \\ 0 & r > a \end{cases} \quad \Phi_H(\mathbf{r}) = -\frac{1}{4\pi} \nabla \cdot \int d^3r' \frac{\mathbf{M}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}$$

For this example:

$$\Phi_H(\mathbf{r}) = -\frac{M_0}{4\pi} \frac{\partial}{\partial z} \left(4\pi \int_0^a r'^2 dr' \frac{1}{r'} \right)$$

$$\text{For } r \leq a: \quad \Phi_H(\mathbf{r}) = -M_0 \frac{\partial}{\partial z} \left(\frac{a^2}{2} - \frac{r^2}{6} \right) = \frac{M_0 z}{3}$$

$$\text{For } r > a: \quad \Phi_H(\mathbf{r}) = -M_0 \frac{\partial}{\partial z} \left(\frac{a^3}{3r} \right) = \frac{M_0 a^3 z}{3r^3}$$

02/17/2014

PHY 712 Spring 2014 – Lecture 13

16

Example magnetostatic boundary value problem -- continued

$$\mathbf{M}(\mathbf{r}) = \begin{cases} M_0 \hat{\mathbf{z}} & r \leq a \\ 0 & r > a \end{cases}$$

$$\text{For } r \leq a: \quad \Phi_H(\mathbf{r}) = \frac{M_0 z}{3} \quad \mathbf{H}(\mathbf{r}) = -\nabla \Phi_H(\mathbf{r}) = -\frac{M_0}{3} \hat{\mathbf{z}}$$

$$\text{For } r > a : \quad \Phi_H(\mathbf{r}) = \frac{M_0 a^3 z}{3r^3} \quad \mathbf{H}(\mathbf{r}) = -\nabla \Phi_H(\mathbf{r}) = -\frac{M_0 a^3}{3} \left(\frac{\hat{\mathbf{z}}}{r^3} - \frac{3z\mathbf{r}}{r^5} \right)$$

$$\mathbf{B}(\mathbf{r}) = \mu_0 (\mathbf{H}(\mathbf{r}) + \mathbf{M}(\mathbf{r}))$$

$$\text{For } r \leq a : \quad \mathbf{H}(\mathbf{r}) = -\frac{M_0 \hat{\mathbf{z}}}{3} \quad \mathbf{B}(\mathbf{r}) = \mu_0 \frac{2M_0 \hat{\mathbf{z}}}{3}$$

$$\text{For } r > a : \quad \mathbf{H}(\mathbf{r}) = -\frac{M_0 a^3}{3} \left(\frac{\hat{\mathbf{z}}}{r^3} - \frac{3\mathbf{z}\mathbf{r}}{r^5} \right)$$

$$\mathbf{B}(\mathbf{r}) = -\mu_0 \frac{M_0 a^3}{3} \left(\frac{\hat{\mathbf{z}}}{r^3} - \frac{3z\mathbf{r}}{r^5} \right)$$

02/17/2014

PHY 712 Spring 2014 – Lecture 13

17

Check boundary values:

$$\text{For } r \leq a : \quad \mathbf{H}(\mathbf{r}) = -\frac{M_0 \hat{\mathbf{z}}}{3} \quad \mathbf{H}(a\hat{\mathbf{r}}) \times \hat{\mathbf{r}} = -\frac{M_0}{3} \hat{\mathbf{z}} \times \hat{\mathbf{r}}$$

$$\text{For } r > a : \quad \mathbf{H}(\mathbf{r}) = -\frac{M_0 a^3}{3} \left(\frac{\hat{\mathbf{z}}}{r^3} - \frac{3z\mathbf{r}}{r^5} \right)$$

$$\mathbf{H}(a\hat{\mathbf{r}}) \times \hat{\mathbf{r}} = -\frac{M_0 a^3}{3} \frac{\hat{\mathbf{z}} \times \hat{\mathbf{r}}}{a^3}$$

$$\text{For } r \leq a : \quad \mathbf{B}(\mathbf{r}) = \mu_0 \frac{2M_0 \hat{\mathbf{z}}}{3} \quad \mathbf{B}(a\hat{\mathbf{r}}) \cdot \hat{\mathbf{r}} = \mu_0 \frac{2M_0}{3} \hat{\mathbf{z}} \cdot \hat{\mathbf{r}}$$

$$\text{For } r > a : \quad \mathbf{B}(\mathbf{r}) = -\mu_0 \frac{M_0 a^3}{3} \left(\frac{\hat{\mathbf{z}}}{r^3} - \frac{3z\mathbf{r}}{r^5} \right)$$

$$\mathbf{B}(a\hat{\mathbf{r}}) \cdot \hat{\mathbf{r}} = -\mu_0 \frac{M_0 a^3}{3} \hat{\mathbf{z}} \cdot \hat{\mathbf{r}} \left(\frac{1}{a^3} - \frac{3a^2}{a^5} \right)$$

03/13/2014

PHY 712, Spring 2014 – Lecture 13

18

Example: permalloy, mumetal $\mu/\mu_0 \sim 10^4$ -- continued

Applying boundary conditions

(only $l = 1$ terms contribute):

$$\text{At } r = a \quad \delta_1 = \frac{\mu}{\mu_0} \left(\beta_1 - 2 \frac{\gamma_1}{a^3} \right)$$

$$a\delta_1 = a\beta_1 + \frac{\gamma_1}{a^2}$$


$$\text{At } r = b \quad \frac{\mu}{\mu_0} \left(\beta_1 - 2 \frac{\gamma_1}{b^3} \right) = - \frac{B_0}{\mu_0} - 2 \frac{\alpha_1}{b^3}$$

$$b\beta_1 + \frac{\gamma_1}{b^2} = -b \frac{B_0}{\mu} + \frac{\alpha_1}{b^2}$$


02/17/2014

PHY 712 Spring 2014 – Lecture 13

25



Example: permalloy, mumetal $\mu/\mu_0 \approx 10^4$ -- continued



When the dust clears:

$$\delta_1 = \left(\frac{-9\mu/\mu_0}{(2\mu/\mu_0 + 1)(\mu/\mu_0 + 2) - 2(a/b)^3(\mu/\mu_0 - 1)^2} \right) \frac{B_0}{\mu_0}$$

$$\approx \frac{1}{\mu/\mu_0} \left(\frac{-9/2}{(1 - (a/b)^3)} \frac{B_0}{\mu_0} \right)$$

02/17/2014

PHY 712 Spring 2014 – Lecture 13

26

Energy associated with magnetic fields

Note : We previously used without proof --

the force on a magnetic dipole \mathbf{m} in an external \mathbf{B} field is:

$$\mathbf{F} = \nabla(\mathbf{m} \cdot \mathbf{B})$$

This implies that energy associated with aligning a magnetic dipole \mathbf{m} in an external \mathbf{B} field is given by :

$$U = -\mathbf{m} \cdot \mathbf{B}$$

Macroscopic energies --

It can be shown that : $W_B = \frac{1}{2} \int d^3r \mathbf{B}(\mathbf{r}) \cdot \mathbf{H}(\mathbf{r})$

$$\text{In analogy to:} \quad W_E = \frac{1}{2} \int d^3r \mathbf{E(r)} \cdot \mathbf{D(r)}$$

02/17/2014

PHY 712 Spring 2014 – Lecture 13

27