

PHY 712 Electrodynamics
9-9:50 AM MWF Olin 105

Plan for Lecture 1:

Reading: Appendix 1 and Chapters I&1

- 1. Course structure and expectations**
- 2. Units – SI vs Gaussian**
- 3. Electrostatics – Poisson equation**

1/17/2018

PHY 712 Spring 2018 – Lecture 1

1

The screenshot shows the WFU Physics website. The 'Events' section contains two entries. The first entry, dated Jan. 17, 2018 at 4 PM, is circled in red and has the word 'Postponed' written over it in a large, semi-transparent font. The second entry, dated Jan. 24, 2018 at 4 PM, is for a colloquium by Professor Yasuke Kanai.

1/17/2018

PHY 712 Spring 2018 – Lecture 1

2

<http://users.wfu.edu/natalie/s18phy712/>

PHY 712 Electrodynamics

MWF 9-9:50 AM | OPL 105 | <http://www.wfu.edu/~natalie/s18phy712/>

Instructor: [Natalie Holzwarth](mailto:natalie@wfu.edu) | Phone: 758-5510 | Office: 300 OPL | e-mail: natalie@wfu.edu

- [General information](#)
- [Syllabus and homework assignments](#)
- [Lecture notes](#)
- [Computer codes](#)
- [Some presentation ideas](#)

Last modified: Monday, 15-Jan-2018 10:11:09 EST

1/17/2018

PHY 712 Spring 2018 – Lecture 1


3

Spring 2018 Schedule
for [N. A. W. Holzwarth](#)

	Monday	Tuesday	Wednesday	Thursday	Friday
9:00-10:00	Electrodynamics PHY712		Electrodynamics PHY712		Electrodynamics PHY712
10:00-12:00	Office Hours		Office Hours		Office Hours
12:00-1:00	Condensed Matter Theory Journal Club	Physics Research	Physics Research	Physics Research	
1:00-3:30	Physics Research		Physics Research		Physics Research
3:30-5:00			Physics Colloquium		


1/17/2018 PHY 712 Spring 2018 – Lecture 1 4

Textbook



1/17/2018

Optional supplement



PHY 712 Spring 2018 – Lecture 1

5

General Information

This course is a one semester survey of Electrodynamics at the graduate level, using the textbook: **Classical Electrodynamics**, 3rd edition, by John David Jackson (John Wiley & Sons, Inc., 1999) – “JDJ”. [link to errata](#) The more recent textbook: **Modern Electrodynamics**, by Andrew Zangwill (Cambridge University Press, 2013) will be used as a supplement.

It is likely that your grade for the course will depend upon the following factors:

Problem sets*	45%
Presentation	10%
Exams	45%

*The schedule notes the “due” date for each assignment. Homeworks may be turned in 1 lecture past their due date without grade penalty. After that, the homework grade will be reduced by 10% for each succeeding late date. According to the honor system, all work submitted for grading purposes should represent the student’s own best efforts. This means that students who work together on homework assignments should all contribute roughly equally and independently verify all derivations and results.

1/17/2018 PHY 712 Spring 2018 – Lecture 1 6

PHY 712 Electrodynamics

MWF 9-9:50 AM | OPL 105 | <http://www.wfu.edu/~natalie/s18phy712/>

Instructor: Natalie Holzwarth | Phone: 758-5510 | Office: 300 OPL | e-mail: natalie@wfu.edu

Course schedule for Spring 2018

(Preliminary schedule -- subject to frequent adjustment.)

Lecture date	JDJ Reading	Topic	HW	Due date
Wed: 01/17/2018	No class	Snow		
1 Fri: 01/19/2018	Chap. 1 & I	Introduction, units and Poisson equation	#1	01/26/2018
2 Mon: 01/22/2018				
3 Wed: 01/24/2018				
4 Thu: 01/25/2018				
5 Fri: 01/26/2018				
6 Mon: 01/29/2018				

1/17/2018 PHY 712 Spring 2018 -- Lecture 1 7

Comment about HW #1: (Jackson problem 1.5)

The time-averaged potential of a neutral hydrogen atom is given by:

$$\Phi(r) = -\frac{q}{4\pi\epsilon_0} \frac{e^{-2r/a_0}}{r} \left(1 + \frac{r}{a_0} \right)$$

where q denotes the magnitude of the elementary charge of an electron or proton and where a_0 denotes the Bohr radius. Find the distribution of charge (both continuous and discrete) that will give this potential and interpret your results physically.

1/17/2018 PHY 712 Spring 2018 -- Lecture 1 8

Some Ideas for Computational Project

The purpose of the "Computational Project" is to provide an opportunity for you to study a topic of your choice in greater depth. The general guideline for your choice of project is that it should have something to do with electrodynamics, and there should be some degree of computation or analysis with the project. The completed project will include a short write-up and a ~20min presentation to the class. You may design your own project or use one of the following list (which will be updated throughout the term).

- Evaluate the ewald sum of various ionic crystals using Maple or a programing language. (Template available in Fortan code.)
- Work out the details of the finite difference or finite element methods.
- Work out the details of the hyperfine Hamiltonian as discussed in Chapter 5 of Jackson.
- Work out the details of Jackson problem 7.2 and related problems.
- Work out the details of reflection and refraction from birefringent materials.
- Analyze the Kramers-Kronig transform of some optical data or calculations.
- Determine the classical electrodynamics associated with an infrared or optical laser.
- Analyze the radiation intensity and spectrum from an interesting source such as an atomic or molecular transition, a free electron laser, etc.
- Work out the details of Jackson problem 14.15.

1/17/2018 PHY 712 Spring 2018 -- Lecture 1 9

Material discussed in Appendix of textbook --

Units - SI vs Gaussian

Coulomb's Law

$$F = K_C \frac{q_1 q_2}{r_{12}^2} \tag{1}$$

Ampere's Law

$$F = K_A \frac{i_1 i_2}{r_{12}^2} d\mathbf{s}_1 \times d\mathbf{s}_2 \times \hat{\mathbf{r}}_{12} \tag{2}$$

In the equations above, the current and charge are related by $i_1 = dq_1/dt$ for all unit systems. The two constants K_C and K_A are related so that their ratio K_C/K_A has the units of $(m/s)^2$ and it is *experimentally* known that the ratio has the value $K_C/K_A = c^2$, where c is the speed of light.

1/17/2018

PHY 712 Spring 2018 – Lecture 1

10

Units - SI vs Gaussian – continued

The choices for these constants in the SI and Gaussian units are given below:

	CGS (Gaussian)	SI
K_C	1	$\frac{1}{4\pi\epsilon_0}$
K_A	$\frac{1}{c^2}$	$\frac{\mu_0}{4\pi}$

Here, $\frac{\mu_0}{4\pi} \equiv 10^{-7} N/A^2$ and $\frac{1}{4\pi\epsilon_0} = c^2 \cdot 10^{-7} N/A^2 = 8.98755 \times 10^9 N \cdot m^2/C^2$.

1/17/2018

PHY 712 Spring 2018 – Lecture 1

11

Units - SI vs Gaussian – continued

Below is a table comparing SI and Gaussian unit systems. The fundamental units for each system are so labeled and are used to define the derived units.

Variable	SI		Gaussian		SI/Gaussian
	Unit	Relation	Unit	Relation	
length	<i>m</i>	fundamental	<i>cm</i>	fundamental	100
mass	<i>kg</i>	fundamental	<i>gm</i>	fundamental	1000
time	<i>s</i>	fundamental	<i>s</i>	fundamental	1
force	<i>N</i>	$kg \cdot m^2/s$	<i>dyne</i>	$gm \cdot cm^2/s$	10^5
current	<i>A</i>	fundamental	<i>statampere</i>	<i>statcoulomb/s</i>	$\frac{1}{10c}$
charge	<i>C</i>	$A \cdot s$	<i>statcoulomb</i>	$\sqrt{dyne \cdot cm^2}$	$\frac{1}{10c}$

1/17/2018

PHY 712 Spring 2018 – Lecture 1

12

Units - SI vs Gaussian – continued

One advantage of the Gaussian system is that the field vectors: \mathbf{E} , \mathbf{D} , \mathbf{B} , \mathbf{H} , \mathbf{P} , \mathbf{M} all have the same physical dimensions. In vacuum, the following equalities hold: $\mathbf{B} = \mathbf{H}$ and $\mathbf{E} = \mathbf{D}$. Also, in the Gaussian system, the dielectric and permittivity constants ϵ and μ are dimensionless.

1/17/2018 PHY 712 Spring 2018 – Lecture 1 13


Basic equations of electrodynamics

CGS (Gaussian)	SI
$\nabla \cdot \mathbf{D} = 4\pi\rho$	$\nabla \cdot \mathbf{D} = \rho$
$\nabla \cdot \mathbf{B} = 0$	$\nabla \cdot \mathbf{B} = 0$
$\nabla \times \mathbf{E} = -\frac{1}{c} \frac{\partial \mathbf{B}}{\partial t}$	$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t}$
$\nabla \times \mathbf{H} = \frac{4\pi}{c} \mathbf{J} + \frac{1}{c} \frac{\partial \mathbf{D}}{\partial t}$	$\nabla \times \mathbf{H} = \mathbf{J} + \frac{\partial \mathbf{D}}{\partial t}$
$\mathbf{F} = q(\mathbf{E} + \frac{\mathbf{v}}{c} \times \mathbf{B})$	$\mathbf{F} = q(\mathbf{E} + \mathbf{v} \times \mathbf{B})$
$u = \frac{1}{8\pi} (\mathbf{E} \cdot \mathbf{D} + \mathbf{B} \cdot \mathbf{H})$	$u = \frac{1}{2} (\mathbf{E} \cdot \mathbf{D} + \mathbf{B} \cdot \mathbf{H})$
$\mathbf{S} = \frac{c}{4\pi} (\mathbf{E} \times \mathbf{H})$	$\mathbf{S} = (\mathbf{E} \times \mathbf{H})$

1/17/2018 PHY 712 Spring 2018 – Lecture 1 14

Units choice for this course:
 SI units for Jackson in Chapters 1-10
 Gaussian units for Jackson in Chapters 11-16

Electrostatics



$$\mathbf{E}(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \sum_i q_i \frac{\mathbf{r} - \mathbf{r}_i}{|\mathbf{r} - \mathbf{r}_i|^3}$$

$$= \frac{1}{4\pi\epsilon_0} \int d^3r' \rho(\mathbf{r}') \frac{\mathbf{r} - \mathbf{r}'}{|\mathbf{r} - \mathbf{r}'|^3}$$

1/17/2018 PHY 712 Spring 2018 – Lecture 1 15

Electrostatics

Discrete versus continuous charge distributions

In terms of Dirac delta function:

$$\rho(\mathbf{r}) = \sum_i q_i \delta(\mathbf{r} - \mathbf{r}_i)$$

Digression: Note that in cartesian coordinates --

$$\delta(\mathbf{r} - \mathbf{r}_i) = \delta(x - x_i)\delta(y - y_i)\delta(z - z_i)$$

1/17/2018

PHY 712 Spring 2018 -- Lecture 1

16

Differential equations --

Electrostatics

$$\nabla \cdot \mathbf{E} = \rho / \epsilon_0$$

$$\nabla \times \mathbf{E} = 0$$

Electrostatic potential

$$\mathbf{E} = -\nabla\Phi(r).$$

$$\nabla^2\Phi(r) = -\rho(r)/\epsilon_0.$$

1/17/2018

PHY 712 Spring 2018 -- Lecture 1

17

Relationship between integral and differential forms of electrostatics --

Differential form

$$\nabla^2\Phi(\mathbf{r}) = -\rho(\mathbf{r}) / \epsilon_0$$

Integral form

$$\Phi(\mathbf{r}) =$$

$$\frac{1}{4\pi\epsilon_0} \int d^3r' \frac{\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}$$

1/17/2018

PHY 712 Spring 2018 -- Lecture 1

18

Relationship between integral and differential forms of electrostatics --

Need to show: $\nabla^2 \left(\frac{1}{|\mathbf{r} - \mathbf{r}'|} \right) = -4\pi \delta^3(\mathbf{r} - \mathbf{r}')$.

Noting that

$$\int_{\text{small sphere about } \mathbf{r}'} d^3r \delta^3(\mathbf{r} - \mathbf{r}') f(\mathbf{r}) = f(\mathbf{r}'),$$

we see that we must show that

$$\int_{\text{small sphere about } \mathbf{r}'} d^3r \nabla^2 \left(\frac{1}{|\mathbf{r} - \mathbf{r}'|} \right) f(\mathbf{r}) = -4\pi f(\mathbf{r}').$$

1/17/2018 PHY 712 Spring 2018 - Lecture 1 19

We introduce a small radius a such that:

$$\frac{1}{|\mathbf{r} - \mathbf{r}'|} = \lim_{a \rightarrow 0} \frac{1}{\sqrt{|\mathbf{r} - \mathbf{r}'|^2 + a^2}}.$$


For a fixed value of a ,

$$\nabla^2 \frac{1}{\sqrt{|\mathbf{r} - \mathbf{r}'|^2 + a^2}} = \frac{-3a^2}{(|\mathbf{r} - \mathbf{r}'|^2 + a^2)^{5/2}}.$$

1/17/2018 PHY 712 Spring 2018 - Lecture 1 20

If the function $f(\mathbf{r})$ is continuous, we can make a Taylor expansion of it about the point $\mathbf{r} = \mathbf{r}'$, keeping only the first term. The integral over the small sphere about \mathbf{r}' can be carried out analytically, by changing to a coordinate system centered at \mathbf{r}' ;

so that

$$\int_{\text{small sphere about } \mathbf{r}'} d^3r \nabla^2 \left(\frac{1}{|\mathbf{r} - \mathbf{r}'|} \right) f(\mathbf{r}) \approx f(\mathbf{r}') \lim_{a \rightarrow 0} \int_{u < R} d^3u \frac{-3a^2}{(u^2 + a^2)^{5/2}}.$$


$\mathbf{u} = \mathbf{r} - \mathbf{r}'$

$$\int_{u < R} d^3u \frac{-3a^2}{(u^2 + a^2)^{5/2}} = 4\pi \int_0^R du \frac{-3a^2 u^2}{(u^2 + a^2)^{5/2}} = 4\pi \frac{-R^3}{(R^2 + a^2)^{3/2}}.$$

1/17/2018 PHY 712 Spring 2018 - Lecture 1 21

$$\int_{u < R} d^3u \frac{-3a^2}{(u^2 + a^2)^{5/2}} = 4\pi \int_0^R du \frac{-3a^2 u^2}{(u^2 + a^2)^{5/2}} = 4\pi \frac{-R^3}{(R^2 + a^2)^{3/2}}.$$

$$\text{For } a \ll R, \quad 4\pi \frac{-R^3}{(R^2 + a^2)^{3/2}} \approx -4\pi$$

$$\rightarrow \int_{\text{small sphere about } \mathbf{r}'} d^3r \nabla^2 \left(\frac{1}{|\mathbf{r} - \mathbf{r}'|} \right) f(\mathbf{r}) \approx f(\mathbf{r}')(-4\pi),$$

$$\rightarrow \nabla^2 \left(\frac{1}{|\mathbf{r} - \mathbf{r}'|} \right) = -4\pi \delta^3(\mathbf{r} - \mathbf{r}')$$

1/17/2018

PHY 712 Spring 2018 – Lecture 1

22

Example in HW1

The electrostatic potential of a neutral H atom is given by:

$$\Phi(r) = \frac{q}{4\pi\epsilon_0} \frac{e^{-\alpha r}}{r} \left(1 + \frac{\alpha r}{2} \right).$$

Find the charge density (both continuous and discrete) for this potential.

Hint #1: For continuous contribution you can use

$$\text{the identity: } \nabla^2 \Phi(r) = \frac{1}{r} \frac{\partial^2 (r\Phi(r))}{\partial r^2}$$

Hint #2: Don't forget to consider possible discrete contributions.

1/17/2018

PHY 712 Spring 2018 – Lecture 1

23
