

PHY 712 Electrodynamics
9-9:50 AM MWF Olin 105
Make-up lecture
Plan for Lecture 4:

Reading: Chapter 1 & 2 in JDJ

Electrostatic potentials

1. One, two, and three dimensions (Cartesian coordinates)
2. Mean value theorem for the electrostatic potential

1/25/2018 PHY 712 Spring 2018 – Lecture 4 1

PHY 712 Electrodynamics

MWF 9-9:50 AM | OPL 105 | <http://www.wfu.edu/~natalie/s18phy712/>

Instructor: Natalie Holzwarth | Phone: 758-5510 | Office: 300 OPL | e-mail: natalie@wfu.edu

Course schedule for Spring 2018
(Preliminary schedule -- subject to frequent adjustment.)

Lecture date	JDJ Reading	Topic	HW	Due date
Wed: 01/17/2018	No class	Snow		
1 Fri: 01/19/2018	Chap. 1 & Appen.	Introduction, units and Poisson equation	#1	01/26/2018
2 Mon: 01/22/2018	Chap. 1	Electrostatic energy calculations	#2	01/26/2018
3 Wed: 01/24/2018	Chap. 1	Poisson's equation and Green's theorem	#3	01/26/2018
4 Thu: 01/25/2018	Chap. 1 & 2	Poisson's equation in 2 and 3 dimensions		
5 Fri: 01/26/2018				
6 Mon: 01/29/2018				
7 Wed: 01/31/2018				

1/25/2018 PHY 712 Spring 2018 – Lecture 4 2

Poisson Equation

$$\nabla^2 \Phi_p(\mathbf{r}) = -\frac{\rho(\mathbf{r})}{\epsilon_0}$$

Solution to Poisson equation using Green's function $G(\mathbf{r}, \mathbf{r}')$:

$$\Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V d^3r' \rho(\mathbf{r}') G(\mathbf{r}, \mathbf{r}') +$$

$$\frac{1}{4\pi} \int_S d^2r' [G(\mathbf{r}, \mathbf{r}') \nabla' \Phi(\mathbf{r}') - \Phi(\mathbf{r}') \nabla' G(\mathbf{r}, \mathbf{r}')] \cdot \hat{\mathbf{r}}'$$

1/25/2018 PHY 712 Spring 2018 – Lecture 4 3

Orthogonal function expansion -- continued

Suppose the orthogonal functions satisfy an eigenvalue equation:

$$\frac{d^2}{dx^2} u_n(x) = -\alpha_n u_n(x)$$

where the functions $u_n(x)$ also satisfy the appropriate boundary conditions, then we can construct the Green's function:

$$G(x, x') = 4\pi \sum_n \frac{u_n(x)u_n(x')}{\alpha_n}$$

Check:

$$\begin{aligned} \frac{d^2}{dx^2} G(x, x') &= 4\pi \sum_n \frac{(-\alpha_n u_n(x))u_n(x')}{\alpha_n} = -4\pi \sum_n u_n(x)u_n(x') \\ &= -4\pi\delta(x - x') \end{aligned}$$

1/25/2018

PHY 712 Spring 2018 -- Lecture 4

7

Example

For example, consider the previous example in the interval

$-a \leq x \leq a$:

$$\rho(x) = \begin{cases} 0 & \text{for } x < -a \\ -\rho_0 & \text{for } -a < x < 0 \\ +\rho_0 & \text{for } 0 < x < a \\ 0 & \text{for } x > a \end{cases}$$

We want to solve the Poisson equation with boundary condition $d\Phi(-a)/dx = 0$ and $d\Phi(a)/dx = 0$. We may choose

$$u_n(x) = \sqrt{\frac{1}{a}} \sin\left(\frac{[2n+1]\pi x}{2a}\right)$$
 and the corresponding Green's function

$$G(x, x') = \frac{4\pi}{a} \sum_{n=0}^{\infty} \frac{\sin\left(\frac{[2n+1]\pi x}{2a}\right) \sin\left(\frac{[2n+1]\pi x'}{2a}\right)}{\left(\frac{[2n+1]\pi}{2a}\right)^2}$$

1/25/2018

PHY 712 Spring 2018 -- Lecture 4

8

Example -- continued

This form of the one-dimensional Green's function only allows us to find a solution to the Poisson equation within the interval $-a \leq x \leq a$ from

$$\Phi(x) = -\frac{1}{4\pi\epsilon_0} \int_{-a}^a dx' G(x, x') \rho(x') + C_1$$

$$\Rightarrow \Phi(x) = \frac{\rho_0 a^2}{\epsilon_0} \left(16 \sum_{n=0}^{\infty} \frac{\sin\left(\frac{[2n+1]\pi x}{2a}\right)}{([2n+1]\pi)^3} + \frac{1}{2} \right),$$

choosing C_1 so that $\Phi(-a) = 0$.

$$\text{Exact result: } \Phi(x) = \begin{cases} 0 & \text{for } x < -a \\ \frac{\rho_0}{2\epsilon_0} (x+a)^2 & \text{for } -a < x < 0 \\ -\frac{\rho_0}{2\epsilon_0} (x-a)^2 + \frac{\rho_0 a^2}{\epsilon_0} & \text{for } 0 < x < a \\ \frac{\rho_0}{\epsilon_0} a^2 & \text{for } x > a \end{cases}$$

1/25/2018

PHY 712 Spring 2018 -- Lecture 4

9

Example -- continued

$$\Phi(x) = \frac{\rho_0 a^2}{\epsilon_0} \left(16 \sum_{n=0}^{\infty} \frac{\sin\left(\frac{[2n+1]\pi x}{2a}\right)}{([2n+1]\pi)^3} + \frac{1}{2} \right)$$

1/25/2018 PHY 712 Spring 2018 - Lecture 4 10

Orthogonal function expansions in 2 and 3 dimensions

$$\nabla^2 \Phi(\mathbf{r}) \equiv \frac{\partial^2 \Phi(\mathbf{r})}{\partial x^2} + \frac{\partial^2 \Phi(\mathbf{r})}{\partial y^2} + \frac{\partial^2 \Phi(\mathbf{r})}{\partial z^2} = -\rho(\mathbf{r}) / \epsilon_0.$$

Let $\{u_n(x)\}$, $\{v_m(y)\}$, $\{w_n(z)\}$ denote complete orthogonal function sets in the x , y , and z dimensions, respectively. The Green's function construction becomes:

$$G(x, x', y, y', z, z') = 4\pi \sum_{lmn} \frac{u_l(x)u_l(x')v_m(y)v_m(y')w_n(z)w_n(z')}{\alpha_l + \beta_m + \gamma_n},$$

where

$$\frac{d^2}{dx^2} u_l(x) = -\alpha_l u_l(x), \quad \frac{d^2}{dy^2} v_m(y) = -\beta_m v_m(y), \quad \text{and} \quad \frac{d^2}{dz^2} w_n(z) = -\gamma_n w_n(z).$$

(See Eq. 3.167 in Jackson for example.)

1/25/2018 PHY 712 Spring 2018 - Lecture 4 11

Combined orthogonal function expansion and homogeneous solution construction of Green's function in 2 and 3 dimensions.

An alternative method of finding Green's functions for a second order ordinary differential equations (in 1 dimension) is based on a product of two independent solutions of the homogeneous equation, $\phi_1(x)$ and $\phi_2(x)$:

$$G(x, x') = K \phi_1(x_<) \phi_2(x_>), \quad \text{where} \quad K \equiv \frac{4\pi}{\frac{d\phi_1}{dx} \phi_2 - \phi_1 \frac{d\phi_2}{dx}},$$

where $x_<$ denotes the smaller of x and x' .

For the two and three dimensional cases, we can use this technique in one of the dimensions in order to reduce the number of summation terms. These ideas are discussed in Section 3.11 of Jackson.

1/25/2018 PHY 712 Spring 2018 - Lecture 4 12

Green's function construction -- continued
 For the two dimensional case, for example, we can assume that the Green's function can be written in the form:

$$G(x, x', y, y') = \sum_n u_n(x) u_n(x') g_n(y, y')$$

The y dependence of this equation will have the required behavior, if we choose: $\left[-\alpha_n + \frac{\partial^2}{\partial y^2}\right] g_n(y, y') = -4\pi\delta(y - y')$,

which in turn can be expressed in terms of the two independent solutions $v_{n_1}(y)$ and $v_{n_2}(y)$ of the homogeneous equation:

$$\left[\frac{d^2}{dy^2} - \alpha_n\right] v_{n_i}(y) = 0,$$

and the Wronskian constant: $K_n \equiv \frac{dv_{n_1}}{dy} v_{n_2} - v_{n_1} \frac{dv_{n_2}}{dy}$

1/25/2018 PHY 712 Spring 2018 - Lecture 4 13

$$\left[-\alpha_n + \frac{\partial^2}{\partial y^2}\right] g_n(y, y') = -4\pi\delta(y - y')$$

$$g_n(y, y') = \frac{4\pi}{K_n} v_{n_1}(y_{<}) v_{n_2}(y_{>})$$


where: $\left[\frac{d^2}{dy^2} - \alpha_n\right] v_{n_i}(y) = 0,$

and $K_n \equiv \frac{dv_{n_1}}{dy} v_{n_2} - v_{n_1} \frac{dv_{n_2}}{dy}$

For example, choose $v_{n_1}(y) = \sinh(\sqrt{\alpha_n} y)$ and $v_{n_2}(y) = \sinh(\sqrt{\alpha_n} (b - y))$
 where $K_n = \sqrt{\alpha_n} \sinh(\sqrt{\alpha_n} b)$
 using the identity: $\cosh(r)\sinh(s) + \sinh(r)\cosh(s) = \sinh(r + s)$

1/25/2018 PHY 712 Spring 2018 - Lecture 4 14

Example:



Two dimensional box with sides a and b with boundary conditions: $\phi(0, y) = \phi(a, y) = \phi(x, 0) = \phi(x, b) = 0$


$$\Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V d^3r' \rho(\mathbf{r}') G(\mathbf{r}, \mathbf{r}') + \frac{1}{4\pi} \int_S d^2r' [G(\mathbf{r}, \mathbf{r}') \nabla' \Phi(\mathbf{r}') - \Phi(\mathbf{r}') \nabla' G(\mathbf{r}, \mathbf{r}')] \cdot \hat{\mathbf{r}}'$$

Don't know this term $\left(\frac{1}{4\pi} \int_S d^2r' [G(\mathbf{r}, \mathbf{r}') \nabla' \Phi(\mathbf{r}') - \Phi(\mathbf{r}') \nabla' G(\mathbf{r}, \mathbf{r}')] \cdot \hat{\mathbf{r}}'\right)$ Know this term $\left(\frac{1}{4\pi\epsilon_0} \int_V d^3r' \rho(\mathbf{r}') G(\mathbf{r}, \mathbf{r}')\right)$

$$G(x, x', y, y') = \sum_n u_n(x) u_n(x') \frac{4\pi}{K_n} v_{n_1}(y_{<}) v_{n_2}(y_{>}).$$

1/25/2018 PHY 712 Spring 2018 - Lecture 4 15

Example:



Two dimensional box with sides a and b with boundary conditions: $\phi(0,y)=\phi(a,y)=\phi(x,0)=\phi(x,b)=0$

For this type of problem, it is prudent to construct $G(x,x',y,y')$ so that it vanishes on the boundary:
 $G(x,x',y,0) = G(x,x',y,b) = G(x,0,y,y') = G(x,a,y,y') = 0$

1/25/2018 PHY 712 Spring 2018 – Lecture 4 16

$$G(x,x',y,y') = \sum_n u_n(x)u_n(x') \frac{4\pi}{K_n} v_{n_1}(y_<)v_{n_2}(y_>).$$

$$\frac{d^2}{dx^2} u_n(x) = -\alpha_n u_n(x) \quad \text{where } u_n(0) = u_n(a) = 0$$

$$\Rightarrow u_n(x) = \sqrt{\frac{2}{a}} \sin\left(\frac{n\pi x}{a}\right) \quad \alpha_n = \left(\frac{n\pi}{a}\right)^2$$

$$\left[\frac{d^2}{dy^2} - \left(\frac{n\pi}{a}\right)^2 \right] v_n(y) = 0$$

$$v_{n_1}(y) = \sinh\left(\frac{n\pi}{a} y\right) \quad v_{n_2}(y) = \sinh\left(\frac{n\pi}{a} (b-y)\right)$$


$$K_n = \frac{n\pi}{a} \sinh\left(\frac{n\pi b}{a}\right)$$

1/25/2018 PHY 712 Spring 2018 – Lecture 4 17


Green's function construction -- continued

$$G(x,x',y,y') = \sum_n u_n(x)u_n(x') K_n v_{n_1}(y_<)v_{n_2}(y_>).$$

For example, a Green's function for a two-dimensional rectangular system with $0 \leq x \leq a$ and $0 \leq y \leq b$, which vanishes on the rectangular boundaries:

$$G(x,x',y,y') = 8 \sum_{n=1}^{\infty} \frac{\sin\left(\frac{n\pi x}{a}\right) \sin\left(\frac{n\pi x'}{a}\right) \sinh\left(\frac{n\pi y_<}{a}\right) \sinh\left(\frac{n\pi}{a} (b-y_>)\right)}{n \sinh\left(\frac{n\pi b}{a}\right)}$$


1/25/2018 PHY 712 Spring 2018 – Lecture 4 18



$$\Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V d^3r' \rho(\mathbf{r}') G(\mathbf{r}, \mathbf{r}') + \frac{1}{4\pi} \int_S d^2r' [G(\mathbf{r}, \mathbf{r}') \nabla' \Phi(\mathbf{r}') - \Phi(\mathbf{r}') \nabla' G(\mathbf{r}, \mathbf{r}')] \cdot \hat{\mathbf{r}}'$$

$$G(x, x', y, y') = 8 \sum_{n=1}^{\infty} \frac{\sin\left(\frac{n\pi x}{a}\right) \sin\left(\frac{n\pi x'}{a}\right) \sinh\left(\frac{n\pi y_{<}}{a}\right) \sinh\left(\frac{n\pi}{a}(b - y_{>})\right)}{n \sinh\left(\frac{n\pi b}{a}\right)}$$

1/25/2018 PHY 712 Spring 2018 – Lecture 4 19

$$G(x, x', y, y') = 8 \sum_{n=1}^{\infty} \frac{\sin\left(\frac{n\pi x}{a}\right) \sin\left(\frac{n\pi x'}{a}\right) \sinh\left(\frac{n\pi y_{<}}{a}\right) \sinh\left(\frac{n\pi}{a}(b - y_{>})\right)}{n \sinh\left(\frac{n\pi b}{a}\right)}$$

Example: $\rho(x, y) = \rho_0 \sin\left(\frac{\pi x}{a}\right) \sin\left(\frac{\pi y}{b}\right)$

$$\Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V d^3r' \rho(\mathbf{r}') G(\mathbf{r}, \mathbf{r}')$$

In this example, only n=1 contributes because

$$\int_0^a dx' \sin\left(\frac{\pi x'}{a}\right) \sin\left(\frac{n\pi x'}{a}\right) = \frac{a}{2} \delta_{n1}$$

$$\Phi(x, y) = \frac{8\rho_0}{4\pi\epsilon_0} \frac{a}{2 \sinh(\pi a/b)} \sin\left(\frac{\pi x}{a}\right) \times \left[\sinh\left(\frac{\pi(b-y)}{a}\right) \int_0^b dy' \sin\left(\frac{\pi y'}{b}\right) \sinh\left(\frac{\pi y'}{a}\right) + \sinh\left(\frac{\pi y}{a}\right) \int_y^b dy' \sin\left(\frac{\pi y'}{b}\right) \sinh\left(\frac{\pi(b-y')}{a}\right) \right]$$

When the dust clears: $\Phi(x, y) = \frac{\rho_0}{\epsilon_0} \frac{a^2 b^2}{\pi^2 (a^2 + b^2)} \sin\left(\frac{\pi x}{a}\right) \sin\left(\frac{\pi y}{b}\right)$

1/25/2018 PHY 712 Spring 2018 – Lecture 4 20

A useful theorem for electrostatics

The mean value theorem (Problem 1.10 in Jackson)

The “mean value theorem” value theorem (problem 1.10 of your textbook) states that the value of $\Phi(\mathbf{r})$ at the arbitrary (charge-free) point \mathbf{r} is equal to the average of $\Phi(\mathbf{r}')$ over the surface of any sphere centered on the point \mathbf{r} (see Jackson problem #1.10). One way to prove this theorem is the following. Consider a point $\mathbf{r}' = \mathbf{r} + \mathbf{u}$, where \mathbf{u} will describe a sphere of radius R about the fixed point \mathbf{r} . We can make a Taylor series expansion of the electrostatic potential $\Phi(\mathbf{r}')$ about the fixed point \mathbf{r} :

$$\Phi(\mathbf{r} + \mathbf{u}) = \Phi(\mathbf{r}) + \mathbf{u} \cdot \nabla \Phi(\mathbf{r}) + \frac{1}{2!} (\mathbf{u} \cdot \nabla)^2 \Phi(\mathbf{r}) + \frac{1}{3!} (\mathbf{u} \cdot \nabla)^3 \Phi(\mathbf{r}) + \frac{1}{4!} (\mathbf{u} \cdot \nabla)^4 \Phi(\mathbf{r}) + \dots \quad (1)$$

According to the premise of the theorem, we want to integrate both sides of the equation 1 over a sphere of radius R in the variable \mathbf{u} :

$$\int_{\text{sphere}} dS_{\mathbf{u}} = R^2 \int_0^{2\pi} d\phi_u \int_{-1}^{+1} d\cos(\theta_u). \quad (2)$$

1/25/2018 PHY 712 Spring 2018 – Lecture 4 21

Mean value theorem - continued

We note that

$$R^2 \int_0^{2\pi} d\phi_u \int_{-1}^{+1} d\cos(\theta_u) 1 = 4\pi R^2,$$

$$R^2 \int_0^{2\pi} d\phi_u \int_{-1}^{+1} d\cos(\theta_u) \mathbf{u} \cdot \nabla = 0,$$

$$R^2 \int_0^{2\pi} d\phi_u \int_{-1}^{+1} d\cos(\theta_u) (\mathbf{u} \cdot \nabla)^2 = \frac{4\pi R^4}{3} \nabla^2,$$

$$R^2 \int_0^{2\pi} d\phi_u \int_{-1}^{+1} d\cos(\theta_u) (\mathbf{u} \cdot \nabla)^3 = 0,$$

and

$$R^2 \int_0^{2\pi} d\phi_u \int_{-1}^{+1} d\cos(\theta_u) (\mathbf{u} \cdot \nabla)^4 = \frac{4\pi R^6}{5} \nabla^4.$$

Since $\nabla^2 \Phi(\mathbf{r}) = 0$, the only non-zero term of the average is thus the first term:

$$R^2 \int_0^{2\pi} d\phi_u \int_{-1}^{+1} d\cos(\theta_u) \Phi(\mathbf{r} + \mathbf{u}) = 4\pi R^2 \Phi(\mathbf{r}),$$

or

$$\Phi(\mathbf{r}) = \frac{1}{4\pi R^2} R^2 \int_0^{2\pi} d\phi_u \int_{-1}^{+1} d\cos(\theta_u) \Phi(\mathbf{r} + \mathbf{u}) \equiv \frac{1}{4\pi R^2} \int_{\text{sphere}} dS_u \Phi(\mathbf{r} + \mathbf{u}).$$

Since this result is independent of the radius R , we see that we have the theorem.

1/25/2018

PHY 712 Spring 2018 - Lecture 4

22
