

PHY 712 Electrodynamics

11-11:50 AM in Olin 103

Class notes for Lecture 4:

Reading: Chapter 1 in JDJ

- 1. Review of electrostatics with one-dimensional examples**
- 2. Poisson and Laplace Equations**
- 3. Green's Theorem and its use in electrostatics**

1/19/2022

PHY 712 Spring 2022 -- Lecture 4

1

In this lecture, we will return to the materials presented in our textbook. Some of the ideas were presented in PHY 711.

PHYSICS COLLOQUIUM

4 PM Olin 101

THURSDAY

JANUARY 20, 2022

"Epigenetic Inheritance from Yeast to Man: Molecular Memory Systems in Health and Disease"

In addition to the well-understood role for DNA as the central carrier for heritable information in biology on this planet, over the past few decades it has become clear that additional information can be passed from one generation to the next via so-called "epigenetic inheritance." The Rando lab focuses on the How, What, and Why of epigenetic inheritance. I will discuss two areas of interest for our lab. In the first, I will describe our efforts to understand the polymer folding puzzle of how the genome is organized in 3-dimensional space, and how this folding affects the function of the genome. In the second, I will describe our studies on effects of a father's diet and lifestyle on health and disease in offspring, and our attempts to address the question of how much information mammalian sperm carry.



Oliver Rando, Ph.D.

Professor
Department of Biochemistry and
Molecular Biotechnology
University of Massachusetts
Medical School
Worcester MA

4:00 pm - Olin 101*

*Link provided for those unable to attend in person.
Note: For additional information on the seminar
or to obtain the video conference link, contact
wfuophys@wfu.edu

1/19/2022

PHY 712 Spring 2022 -- Lecture 4

2

PHY 712 Electrodynamics

MWF 11-11:50 AM Olin 103 Webpage: <http://www.wfu.edu/~natalie/s22phy712/>

Instructor: [Natalie Holzwarth](#) Office: 300 OPL e-mail: natalie@wfu.edu

Course schedule for Spring 2022

(Preliminary schedule -- subject to frequent adjustment.)

	Lecture date	JDJ Reading	Topic	HW	Due date
1	Mon: 01/10/2022	Chap. 1 & Appen.	Introduction, units and Poisson equation	#1	01/14/2022
2	Wed: 01/12/2022	Chap. 1	Electrostatic energy calculations	#2	01/19/2022
3	Fri: 01/14/2022	Chap. 1	Electrostatic energy calculations	#3	01/21/2022
	Mon: 01/17/2022		MLK Holiday -- no class		
4	Wed: 01/19/2022	Chap. 1 & 2	Electrostatic potentials and fields	#4	01/24/2022
5	Fri: 01/21/2022	Chap. 1 - 3	Poisson's equation in 2 and 3 dimensions		

1/19/2022

PHY 712 Spring 2022 -- Lecture 4

3

Updated schedule. Note new homework assignment which follows from today's lecture.

January 19, 2022

PHY 712 – Problem Set #4

Continue reading Chapter 1 & 2 in **Jackson**

1. Consider a one-dimensional charge distribution of the form:

$$\rho(x) = \begin{cases} 0 & \text{for } x < -a \\ \rho_0 \sin(\pi x/a) & \text{for } -a \leq x \leq a \\ 0 & \text{for } x > a, \end{cases}$$

where ρ_0 and a are constants.

- (a) Solve the Poisson equation for the electrostatic potential $\Phi(x)$ with the boundary conditions $\Phi(-a) = 0$ and $\frac{d\Phi}{dx}(-a) = 0$.
- (b) Find the corresponding electrostatic field $E(x)$.
- (c) Plot $\Phi(x)$ and $E(x)$.
- (d) Discuss your results in terms of elementary application of Gauss's Law arguments.

1/19/2022

PHY 712 Spring 2022 -- Lecture 4

4

Content of IHW 3.

Poisson and Laplace Equations

We are concerned with finding solutions to the Poisson equation:

$$\nabla^2 \Phi_P(\mathbf{r}) = -\frac{\rho(\mathbf{r})}{\epsilon_0}$$

and the Laplace equation:

$$\nabla^2 \Phi_L(\mathbf{r}) = 0$$

The Laplace equation is the “homogeneous” version of the Poisson equation. The Green's theorem allows us to determine the electrostatic potential from volume and surface integrals:

$$\Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V d^3r' \rho(\mathbf{r}') G(\mathbf{r}, \mathbf{r}') + \frac{1}{4\pi} \int_S d^2r' [G(\mathbf{r}, \mathbf{r}') \nabla' \Phi(\mathbf{r}') - \Phi(\mathbf{r}') \nabla' G(\mathbf{r}, \mathbf{r}')] \cdot \hat{\mathbf{r}}'.$$

1/19/2022

PHY 712 Spring 2022 -- Lecture 4

5

Here we start our systematic derivations of solution of the electrostatic equation for a potential with a given charge source and the associated homogeneous equation.

Poisson equation -- continued

Note that we have previously shown that the differential and integral forms of Coulomb's law is given by:

$$\nabla^2 \Phi(\mathbf{r}) = -\frac{\rho(\mathbf{r})}{\epsilon_0} \quad \text{and} \quad \Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V d^3r' \frac{\rho(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}$$

Generalization of analysis for non-trivial boundary conditions:

$$\Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V d^3r' \rho(\mathbf{r}') G(\mathbf{r}, \mathbf{r}') + \frac{1}{4\pi} \int_S d^2r' [G(\mathbf{r}, \mathbf{r}') \nabla' \Phi(\mathbf{r}') - \Phi(\mathbf{r}') \nabla' G(\mathbf{r}, \mathbf{r}')] \cdot \hat{\mathbf{r}}'.$$

1/19/2022

PHY 712 Spring 2022 -- Lecture 4

6

What we discussed last week is still true for isolated charges. Now we consider the case where the charges are within a volume V whose surface may have some imposed restrictions (boundary conditions).

General comments on Green's theorem

$$\Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V d^3r' \rho(\mathbf{r}') G(\mathbf{r}, \mathbf{r}') + \frac{1}{4\pi} \int_S d^2r' \left[G(\mathbf{r}, \mathbf{r}') \nabla' \Phi(\mathbf{r}') - \Phi(\mathbf{r}') \nabla' G(\mathbf{r}, \mathbf{r}') \right] \cdot \hat{\mathbf{r}}'.$$

This general form can be used in 1, 2, or 3 dimensions. In general, the Green's function must be constructed to satisfy the appropriate (Dirichlet or Neumann) boundary conditions. Alternatively, or in addition, boundary conditions can be adjusted using the fact that for any solution to the Poisson equation, $\Phi_p(\mathbf{r})$ other solutions may be generated by use of solutions of the Laplace equation

$$\Phi(\mathbf{r}) = \Phi_p(\mathbf{r}) + C\Phi_L(\mathbf{r}), \text{ for any constant } C.$$

1/19/2022

PHY 712 Spring 2022 -- Lecture 4

7

Comment about how the boundary conditions may or may not work. Note that it is important to not over specify the boundary conditions..

The Green of Green Functions

In 1828, an English miller from Nottingham published a mathematical essay that generated little response. George Green's analysis, however, has since found applications in areas ranging from classical electrostatics to modern quantum field theory.

Lawrie Challis and Fred Sheard

Nottingham, an attractive and thriving town in the English Midlands, is famous for its association with Robin Hood, whose statue stands in the shadow of the castle wall. The Sheriff of Nottingham still has a special role in the city government although happily no longer strikes terror into the hearts of the good citizens.

Recently a new attraction, a windmill, has appeared on the Nottingham skyline (see figure 1). The sails turn on windy days and the adjoining mill shop sells packets of stone ground flour but also, more surprisingly, tracts on mathematical physics. The connection between the flour and the physics is part of the mill's unique character and is explained by a plaque once attached to the side of the mill tower that said,

HERE LIVED AND LABOURED
GEORGE GREEN
MATHEMATICIAN
B.1793–D.1841.

his family built a house next to the mill, Green spent most of his days and many of his nights working and indeed living in the mill. When he was 31, Jane Smith bore him a daughter. They had seven children in all but never married. It was said that Green's father felt that Jane was not a suitable wife for the son of a prosperous tradesman and landowner and threatened to disinherit him.

Little is known about Green's life from 1802 until 1823. In particular, it is not known whether he received any help in his mathematical development or if he was entirely self-taught. He may have received help from John Toplis, a fellow of Queens' College in the University of Cambridge and headmaster of the Nottingham Grammar School. Toplis's translation of Pierre-Simon Laplace's book *Mécanique Céleste*, published in Nottingham in 1814, seems a likely source of Green's interest in potential theory. The work was unusual in Britain at that time inasmuch as Toplis used Gottfried Leibniz's more convenient notation for differentials rather than Isaac Newton's. Because Green adapted the Leibniz notation, it seems plausible that Green was influenced by Toplis, but there is no evidence that Toplis acted in any way as his tutor.

In 1823, Green joined the Nottingham Subscription Library, the center of intellectual activity in the town. The library was situated in Bromley House (see figure 2). Library membership provided Green with encouragement

“Derivation” of Green's Theorem

$$\text{Poisson equation: } \nabla^2 \Phi(\mathbf{r}) = -\frac{\rho(\mathbf{r})}{\epsilon_0}$$

$$\text{Green's relation: } \nabla'^2 G(\mathbf{r}, \mathbf{r}') = -4\pi\delta^3(\mathbf{r} - \mathbf{r}').$$

$$\text{Divergence theorem: } \int_V d^3r \nabla \cdot \mathbf{A} = \oint_S d^2r \mathbf{A} \cdot \hat{\mathbf{r}}$$

$$\text{Let } \mathbf{A} = f(\mathbf{r})\nabla g(\mathbf{r}) - g(\mathbf{r})\nabla f(\mathbf{r})$$

$$\int_V d^3r \nabla \cdot (f(\mathbf{r})\nabla g(\mathbf{r}) - g(\mathbf{r})\nabla f(\mathbf{r})) = \oint_S d^2r (f(\mathbf{r})\nabla g(\mathbf{r}) - g(\mathbf{r})\nabla f(\mathbf{r})) \cdot \hat{\mathbf{r}}$$



$$\int_V d^3r (f(\mathbf{r})\nabla^2 g(\mathbf{r}) - g(\mathbf{r})\nabla^2 f(\mathbf{r}))$$

Here we derive the equations stated on the previous slides.

“Derivation” of Green's Theorem

Poisson equation: $\nabla^2 \Phi(\mathbf{r}) = -\frac{\rho(\mathbf{r})}{\epsilon_0}$

Green's relation: $\nabla'^2 G(\mathbf{r}, \mathbf{r}') = -4\pi\delta^3(\mathbf{r} - \mathbf{r}')$.

$$\int_V d^3r \left(f(\mathbf{r}) \nabla^2 g(\mathbf{r}) - g(\mathbf{r}) \nabla^2 f(\mathbf{r}) \right) = \oint_S d^2r \left(f(\mathbf{r}) \nabla g(\mathbf{r}) - g(\mathbf{r}) \nabla f(\mathbf{r}) \right) \cdot \hat{\mathbf{r}}$$

$$f(\mathbf{r}) \leftrightarrow \Phi(\mathbf{r}) \qquad g(\mathbf{r}) = G(\mathbf{r}, \mathbf{r}')$$

$$\Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V d^3r' \rho(\mathbf{r}') G(\mathbf{r}, \mathbf{r}') +$$

$$\frac{1}{4\pi} \int_S d^2r' \left[G(\mathbf{r}, \mathbf{r}') \nabla' \Phi(\mathbf{r}') - \Phi(\mathbf{r}') \nabla' G(\mathbf{r}, \mathbf{r}') \right] \cdot \hat{\mathbf{r}}'.$$

1/19/2022

PHY 712 Spring 2022 -- Lecture 4

10

Derivation continued.

Example of charge density and potential varying in one dimension

Consider the following one dimensional charge distribution:

$$\rho(x) = \begin{cases} 0 & \text{for } x < -a \\ -\rho_0 & \text{for } -a < x < 0 \\ +\rho_0 & \text{for } 0 < x < a \\ 0 & \text{for } x > a \end{cases}$$

We want to find the electrostatic potential such that

$$\frac{d^2\Phi(x)}{dx^2} = -\frac{\rho(x)}{\varepsilon_0},$$

with the boundary condition $\Phi(-\infty) = 0$ and $\frac{d\Phi}{dx}(-\infty) = 0$

Simple one-dimensional example of a particular charge distribution.

Electrostatic field solution

The solution to the Poisson equation is given by:


$$\Phi(x) = \begin{cases} 0 & \text{for } x < -a \\ \frac{\rho_0}{2\varepsilon_0}(x+a)^2 & \text{for } -a < x < 0 \\ -\frac{\rho_0}{2\varepsilon_0}(x-a)^2 + \frac{\rho_0 a^2}{\varepsilon_0} & \text{for } 0 < x < a \\ \frac{\rho_0}{\varepsilon_0}a^2 & \text{for } x > a \end{cases} .$$

Laplace
Poisson
Poisson
Laplace

The electrostatic field is given by:

$$E(x) = \begin{cases} 0 & \text{for } x < -a \\ -\frac{\rho_0}{\varepsilon_0}(x+a) & \text{for } -a < x < 0 \\ \frac{\rho_0}{\varepsilon_0}(x-a) & \text{for } 0 < x < a \\ 0 & \text{for } x > a \end{cases} .$$

Resultant potential and electric field.



Graph of results for this example

Comment about the example and solution

This particular example is one that is used to model semiconductor junctions where the charge density is controlled by introducing charged impurities near the junction.

The solution of the Poisson equation for this case can be determined by piecewise solution within each of the four regions. Alternatively, from Green's theorem in one-dimension, one can use the Green's function

$$\Phi(x) = \frac{1}{4\pi\epsilon_0} \int_{-\infty}^{\infty} G(x, x') \rho(x') dx' \quad \text{where} \quad G(x, x') = 4\pi x_{<}$$

$x_{<}$ should be taken as the smaller of x and x' .

Comment and generalization.

Notes on the one-dimensional Green's function

The Green's function for the one-dimensional Poisson equation can be defined as a solution to the equation: $\nabla^2 G(x, x') = -4\pi\delta(x - x')$

Here the factor of 4π is not really necessary, but ensures consistency with your text's treatment of the 3-dimensional case. The meaning of this expression is that x' is held fixed while taking the derivative with respect to x .

Some details.

Construction of a Green's function in one dimension

Consider two independent solutions to the homogeneous equation

$$\nabla^2 \phi_i(x) = 0$$

where $i = 1$ or 2 . Let

$$G(x, x') = \frac{4\pi}{W} \phi_1(x_<) \phi_2(x_>).$$

This notation means that $x_<$ should be taken as the smaller of x and x' and $x_>$ should be taken as the larger.

W is defined as the "Wronskian":

$$W \equiv \frac{d\phi_1(x)}{dx} \phi_2(x) - \phi_1(x) \frac{d\phi_2(x)}{dx}.$$

Details continued for one dimensional Poisson equation.

Summary

$$\nabla^2 G(x, x') = -4\pi\delta(x - x')$$

$$G(x, x') = \frac{4\pi}{W} \phi_1(x_<) \phi_2(x_>)$$

$$W \equiv \frac{d\phi_1(x)}{dx} \phi_2(x) - \phi_1(x) \frac{d\phi_2(x)}{dx}$$

$$\left. \frac{dG(x, x')}{dx} \right|_{x=x'+\epsilon} - \left. \frac{dG(x, x')}{dx} \right|_{x=x'-\epsilon} = -4\pi$$

Summary for one dimensional Poisson equation.

One dimensional Green's function in practice

$$\begin{aligned}\Phi(x) &= \frac{1}{4\pi\epsilon_0} \int_{-\infty}^{\infty} G(x, x') \rho(x') dx' \\ &= \frac{1}{4\pi\epsilon_0} \left\{ \int_{-\infty}^x G(x, x') \rho(x') dx' + \int_x^{\infty} G(x, x') \rho(x') dx' \right\}\end{aligned}$$

For the one-dimensional Poisson equation, we can construct the Green's function by choosing $\phi_1(x) = x$ and $\phi_2(x) = 1; W = 1$:

$$\Phi(x) = \frac{1}{\epsilon_0} \left\{ \int_{-\infty}^x x' \rho(x') dx' + x \int_x^{\infty} \rho(x') dx' \right\}.$$

$$G(x, x') = 4\pi x_{<}$$

This expression gives the same result as previously obtained for the example $\rho(x)$ and more generally is appropriate for any neutral charge distribution.

Some general comments.

Question -- How do we know which one of x and x' is the $x_<$ term?

$$G(x, x') = 4\pi x_<$$

$$\Phi(x) = \frac{1}{\epsilon_0} \left\{ \int_{-\infty}^x \underset{x' < x}{x' \rho(x') dx'} + x \int_x^{\infty} \underset{x' > x}{\rho(x') dx'} \right\}.$$

Orthogonal function expansions and Green's functions

Suppose we have a “complete” set of orthogonal functions $\{u_n(x)\}$ defined in the interval $x_1 \leq x \leq x_2$ such that

$$\int_{x_1}^{x_2} u_n(x) u_m(x) dx = \delta_{nm}.$$

We can show that the completeness of this functions implies that

$$\sum_{n=1}^{\infty} u_n(x) u_n(x') = \delta(x - x').$$

This relation allows us to use these functions to represent a Green's function for our system. For the 1-dimensional Poisson equation, the Green's function satisfies

$$\frac{\partial^2}{\partial x^2} G(x, x') = -4\pi\delta(x - x').$$

Now we will discuss another approach to analyzing Green's functions based on expansion in terms of a complete set of orthogonal functions.

Orthogonal function expansions –continued

Therefore, if

$$\frac{d^2}{dx^2} u_n(x) = -\alpha_n u_n(x),$$

where $\{u_n(x)\}$ also satisfy the appropriate boundary conditions, then we can write the Green's functions as

$$G(x, x') = 4\pi \sum_n \frac{u_n(x) u_n(x')}{\alpha_n}.$$

Some details for orthogonal function expansion method.

Example

For example, consider the example discussed earlier in the interval $-a \leq x \leq a$ with

$$\rho(x) = \begin{cases} 0 & \text{for } x < -a \\ -\rho_0 & \text{for } -a < x < 0 \\ +\rho_0 & \text{for } 0 < x < a \\ 0 & \text{for } x > a \end{cases} \quad (24)$$

We want to solve the Poisson equation with boundary condition $d\Phi(-a)/dx = 0$ and $d\Phi(a)/dx = 0$. For this purpose, we may choose

$$u_n(x) = \sqrt{\frac{1}{a}} \sin\left(\frac{[2n+1]\pi x}{2a}\right). \quad (25)$$

The Green's function for this case as:


$$G(x, x') = \frac{4\pi}{a} \sum_{n=0}^{\infty} \frac{\sin\left(\frac{[2n+1]\pi x}{2a}\right) \sin\left(\frac{[2n+1]\pi x'}{2a}\right)}{\left(\frac{[2n+1]\pi}{2a}\right)^2}. \quad (26)$$

Application to our example.

Example – continued

$$\Phi(x) = \frac{\rho_0 a^2}{\epsilon_0} \left(16 \sum_{n=0}^{\infty} \frac{\sin\left(\frac{[2n+1]\pi x}{2a}\right)}{([2n+1]\pi)^3} + \frac{1}{2} \right).$$

Constant shift to allow $\Phi(0) = 0$.



1/19/2022

PHY 712 Spring 2022 -- Lecture 4

23

Graph of potential (green) and expansion for a few terms. Note that it was necessary to shift the potential by a constant.