

PHY 712 Electrodynamics

10-10:50 AM MWF in Olin 103

Lecture Notes for Lecture 15:

Start reading Chapter 6

- 1. Maxwell's full equations; effects of time varying fields and sources**
- 2. Gauge choices and transformations**
- 3. Green's function for vector and scalar potentials**

	Lecture date	JDJ Reading	Topic	HW	Due date
1	Mon: 01/9/2023	Chap. 1 & Appen.	Introduction, units and Poisson equation	#1	01/13/2023
2	Wed: 01/11/2023	Chap. 1	Electrostatic energy calculations	#2	01/18/2023
3	Fri: 01/13/2023	Chap. 1	Electrostatic energy calculations thanks to Ewald	#3	01/18/2023
	Mon: 01/16/2023		MLK Holiday -- no class		
4	Wed: 01/18/2023	Chap. 1 & 2	Electrostatic potentials and fields	#4	01/20/2023
5	Fri: 01/20/2023	Chap. 1 - 3	Poisson's equation in 2 and 3 dimensions	#5	01/23/2023
6	Mon: 01/23/2023	Chap. 1 - 3	Brief introduction to numerical methods	#6	01/25/2023
7	Wed: 01/25/2023	Chap. 2 & 3	Image charge constructions	#7	01/30/2023
8	Fri: 01/27/2023	Chap. 2 & 3	Cylindrical and spherical geometries		
9	Mon: 01/30/2023	Chap. 3 & 4	Spherical geometry and multipole moments	#8	02/01/2023
10	Wed: 02/01/2023	Chap. 4	Dipoles and Dielectrics	#9	02/03/2023
11	Fri: 02/03/2023	Chap. 4	Dipoles and Dielectrics	#10	02/06/2023
12	Mon: 02/06/2023	Chap. 5	Magnetostatics	#11	02/08/2023
13	Wed: 02/08/2023	Chap. 5	Magnetic dipoles and hyperfine interaction	#12	02/10/2023
14	Fri: 02/10/2023	Chap. 5	Magnetic dipoles and dipolar fields	#13	02/13/2023
15	Mon: 02/13/2023	Chap. 6	Maxwell's Equations	#14	02/17/2023
16	Wed: 02/15/2023	Chap. 6	Electromagnetic energy and forces		
17	Fri: 02/17/2023	Chap. 7	Electromagnetic plane waves		
18	Mon: 02/20/2023	Chap. 7	Electromagnetic plane waves		
19	Wed: 02/22/2023	Chap. 7	Optical effects of refractive indices		
20	Fri: 02/24/2023	Chap. 8	Brief introduction to wave guides		

PHY 712 – Problem Set #14

Start reading Chapter 6 in **Jackson**

1. In deriving the Liénard Wiechert potentials, the retarded time

$$t_r = t - \frac{|\mathbf{r} - \mathbf{R}_q(t_r)|}{c}$$

was introduced. Note that this expression for the retarded time t_r may be difficult to evaluate in practice since the right hand side of the equation also depends on t_r . Here \mathbf{r}, t represent the position and time at which the field is measured, and $\mathbf{R}_q(t')$ represents the trajectory of the charged particle as a function of its time denoted by t' . The velocity of the particle is given by

$$\mathbf{v} \equiv \mathbf{v}(t') \equiv \frac{d\mathbf{R}_q(t')}{dt'}$$

Demonstrate the following identities:

(a)

$$\frac{\partial t_r}{\partial t} = \frac{1}{1 - \frac{\mathbf{v}(t_r) \cdot (\mathbf{r} - \mathbf{R}_q(t_r))}{c|\mathbf{r} - \mathbf{R}_q(t_r)|}}$$

(b)

$$-c\nabla t_r = \frac{\frac{\mathbf{r} - \mathbf{R}_q(t_r)}{|\mathbf{r} - \mathbf{R}_q(t_r)|}}{1 - \frac{\mathbf{v}(t_r) \cdot (\mathbf{r} - \mathbf{R}_q(t_r))}{c|\mathbf{r} - \mathbf{R}_q(t_r)|}}$$

Full electrodynamics with time varying fields and sources

Maxwell's equations




Image of statue of
James Clerk-Maxwell
(1831-1879) in Edinburgh

"From a long view of the history of mankind - seen from, say, ten thousand years from now - there can be little doubt that the most significant event of the 19th century will be judged as Maxwell's discovery of the laws of electrodynamics"

Richard P Feynman

<http://www.clerkmaxwellfoundation.org/>

Maxwell's equations

Coulomb's law :

$$\nabla \cdot \mathbf{D} = \rho_{free}$$

Ampere - Maxwell's law :

$$\nabla \times \mathbf{H} - \frac{\partial \mathbf{D}}{\partial t} = \mathbf{J}_{free}$$

Faraday's law :

$$\nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0$$

No magnetic monopoles :

$$\nabla \cdot \mathbf{B} = 0$$

Maxwell's equations

Microscopic or vacuum form ($\mathbf{P} = 0$; $\mathbf{M} = 0$):

Coulomb's law : $\nabla \cdot \mathbf{E} = \rho / \epsilon_0$

Ampere - Maxwell's law : $\nabla \times \mathbf{B} - \frac{1}{c^2} \frac{\partial \mathbf{E}}{\partial t} = \mu_0 \mathbf{J}$

Faraday's law : $\nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0$

No magnetic monopoles : $\nabla \cdot \mathbf{B} = 0$

$$\Rightarrow c^2 = \frac{1}{\epsilon_0 \mu_0}$$

Formulation of Maxwell's equations in terms of vector and scalar potentials

$$\nabla \cdot \mathbf{B} = 0 \quad \Rightarrow \quad \mathbf{B} = \nabla \times \mathbf{A}$$

$$\nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0 \quad \Rightarrow \quad \nabla \times \left(\mathbf{E} + \frac{\partial \mathbf{A}}{\partial t} \right) = 0$$

$$\mathbf{E} + \frac{\partial \mathbf{A}}{\partial t} = -\nabla \Phi$$

$$\text{or } \mathbf{E} = -\nabla \Phi - \frac{\partial \mathbf{A}}{\partial t}$$

Formulation of Maxwell's equations in terms of vector and scalar potentials -- continued

$$\nabla \cdot \mathbf{E} = \rho / \epsilon_0 :$$

$$-\nabla^2 \Phi - \frac{\partial(\nabla \cdot \mathbf{A})}{\partial t} = \rho / \epsilon_0$$

$$\nabla \times \mathbf{B} - \frac{1}{c^2} \frac{\partial \mathbf{E}}{\partial t} = \mu_0 \mathbf{J}$$

$$\nabla \times (\nabla \times \mathbf{A}) + \frac{1}{c^2} \left(\frac{\partial(\nabla \Phi)}{\partial t} + \frac{\partial^2 \mathbf{A}}{\partial t^2} \right) = \mu_0 \mathbf{J}$$

Formulation of Maxwell's equations in terms of vector and scalar potentials -- continued

General form for the scalar and vector potential equations:

$$-\nabla^2\Phi - \frac{\partial(\nabla \cdot \mathbf{A})}{\partial t} = \rho / \epsilon_0$$

$$\nabla \times (\nabla \times \mathbf{A}) + \frac{1}{c^2} \left(\frac{\partial(\nabla\Phi)}{\partial t} + \frac{\partial^2 \mathbf{A}}{\partial t^2} \right) = \mu_0 \mathbf{J}$$

Coulomb gauge form -- require $\nabla \cdot \mathbf{A}_C = 0$

$$-\nabla^2\Phi_C = \rho / \epsilon_0$$

$$-\nabla^2 \mathbf{A}_C + \frac{1}{c^2} \frac{\partial^2 \mathbf{A}_C}{\partial t^2} + \frac{1}{c^2} \frac{\partial(\nabla\Phi_C)}{\partial t} = \mu_0 \mathbf{J}$$

Note that $\mathbf{J} = \mathbf{J}_l + \mathbf{J}_t$ with $\nabla \times \mathbf{J}_l = 0$ and $\nabla \cdot \mathbf{J}_t = 0$

Formulation of Maxwell's equations in terms of vector and scalar potentials -- continued

Coulomb gauge form -- require $\nabla \cdot \mathbf{A}_C = 0$

$$-\nabla^2 \Phi_C = \rho / \epsilon_0$$

$$-\nabla^2 \mathbf{A}_C + \frac{1}{c^2} \frac{\partial^2 \mathbf{A}_C}{\partial t^2} + \frac{1}{c^2} \frac{\partial(\nabla \Phi_C)}{\partial t} = \mu_0 \mathbf{J}$$

Note that $\mathbf{J} = \mathbf{J}_l + \mathbf{J}_t$ with $\nabla \times \mathbf{J}_l = 0$ and $\nabla \cdot \mathbf{J}_t = 0$

Continuity equation for charge and current density :

$$\begin{aligned} \frac{\partial \rho}{\partial t} + \nabla \cdot \mathbf{J}_l = 0 &\quad \Rightarrow \quad \frac{\partial \rho}{\partial t} = -\nabla \cdot \mathbf{J}_l = -\epsilon_0 \nabla \cdot \frac{\partial(\nabla \Phi_C)}{\partial t} \\ &\quad \Rightarrow \quad \frac{1}{c^2} \frac{\partial(\nabla \Phi_C)}{\partial t} = \epsilon_0 \mu_0 \frac{\partial(\nabla \Phi_C)}{\partial t} = \mu_0 \mathbf{J}_l \end{aligned}$$

$$-\nabla^2 \mathbf{A}_C + \frac{1}{c^2} \frac{\partial^2 \mathbf{A}_C}{\partial t^2} = \mu_0 \mathbf{J}_t$$

Formulation of Maxwell's equations in terms of vector and scalar potentials -- continued

Review of the general equations:

$$-\nabla^2\Phi - \frac{\partial(\nabla \cdot \mathbf{A})}{\partial t} = \rho / \epsilon_0$$

$$\nabla \times (\nabla \times \mathbf{A}) + \frac{1}{c^2} \left(\frac{\partial(\nabla\Phi)}{\partial t} + \frac{\partial^2 \mathbf{A}}{\partial t^2} \right) = \mu_0 \mathbf{J}$$

Lorentz gauge form -- require $\nabla \cdot \mathbf{A}_L + \frac{1}{c^2} \frac{\partial\Phi_L}{\partial t} = 0$

$$-\nabla^2\Phi_L + \frac{1}{c^2} \frac{\partial^2\Phi_L}{\partial t^2} = \rho / \epsilon_0$$

$$-\nabla^2\mathbf{A}_L + \frac{1}{c^2} \frac{\partial^2\mathbf{A}_L}{\partial t^2} = \mu_0 \mathbf{J}$$

Recall that

$$\nabla \times (\nabla \times \mathbf{A}) = \nabla(\nabla \cdot \mathbf{A}) - \nabla^2 \mathbf{A}$$

Formulation of Maxwell's equations in terms of vector and scalar potentials -- continued

Lorentz gauge form - - require $\nabla \cdot \mathbf{A}_L + \frac{1}{c^2} \frac{\partial \Phi_L}{\partial t} = 0$

$$-\nabla^2 \Phi_L + \frac{1}{c^2} \frac{\partial^2 \Phi_L}{\partial t^2} = \rho / \epsilon_0$$

$$-\nabla^2 \mathbf{A}_L + \frac{1}{c^2} \frac{\partial^2 \mathbf{A}_L}{\partial t^2} = \mu_0 \mathbf{J}$$

Alternate potentials: $\mathbf{A}'_L = \mathbf{A}_L + \nabla \Lambda$ and $\Phi'_L = \Phi_L - \frac{\partial \Lambda}{\partial t}$

Yields same physics provided that: $\nabla^2 \Lambda - \frac{1}{c^2} \frac{\partial^2 \Lambda}{\partial t^2} = 0$

Solution of Maxwell's equations in the Lorentz gauge

$$\nabla^2 \Phi_L - \frac{1}{c^2} \frac{\partial^2 \Phi_L}{\partial t^2} = -\rho / \epsilon_0$$

$$\nabla^2 \mathbf{A}_L - \frac{1}{c^2} \frac{\partial^2 \mathbf{A}_L}{\partial t^2} = -\mu_0 \mathbf{J}$$

Consider the general form of the 3 - dimensional wave equation :

$$\nabla^2 \Psi - \frac{1}{c^2} \frac{\partial^2 \Psi}{\partial t^2} = -4\pi f$$

$\Psi(\mathbf{r}, t) \Rightarrow$ wave field

$f(\mathbf{r}, t) \Rightarrow$ source

Solution of Maxwell's equations in the Lorentz gauge -- continued

Let Ψ represent Φ, A_x, A_y, A_z Let f represent ρ, J_x, J_y, J_z

$$\nabla^2 \Psi(\mathbf{r}, t) - \frac{1}{c^2} \frac{\partial^2 \Psi(\mathbf{r}, t)}{\partial t^2} = -4\pi f(\mathbf{r}, t)$$

Green's function :

$$\left(\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) G(\mathbf{r}, t; \mathbf{r}', t') = -4\pi \delta^3(\mathbf{r} - \mathbf{r}') \delta(t - t')$$

Formal solution for field $\Psi(\mathbf{r}, t)$:

$$\Psi(\mathbf{r}, t) = \Psi_{f=0}(\mathbf{r}, t) + \int d^3 r' \int dt' G(\mathbf{r}, t; \mathbf{r}', t') f(\mathbf{r}', t')$$

Solution of Maxwell's equations in the Lorentz gauge -- continued

Determination of the form for the Green's function :

$$\left(\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) G(\mathbf{r}, t; \mathbf{r}', t') = -4\pi \delta^3(\mathbf{r} - \mathbf{r}') \delta(t - t')$$

For the case of isotropic boundary values at infinity :

$$G(\mathbf{r}, t; \mathbf{r}', t') = \frac{1}{|\mathbf{r} - \mathbf{r}'|} \delta\left(t' - \left(t - \frac{1}{c} |\mathbf{r} - \mathbf{r}'| \right) \right)$$

Formal solution for field $\Psi(\mathbf{r}, t)$:

$$\Psi(\mathbf{r}, t) = \Psi_{f=0}(\mathbf{r}, t) + \int d^3 r' \int dt' \frac{1}{|\mathbf{r} - \mathbf{r}'|} \delta\left(t' - \left(t - \frac{1}{c} |\mathbf{r} - \mathbf{r}'| \right) \right) f(\mathbf{r}', t')$$

Solution of Maxwell's equations in the Lorentz gauge -- continued

Analysis of the Green's function:

$$\left(\nabla^2 - \frac{1}{c^2} \frac{\partial^2}{\partial t^2} \right) G(\mathbf{r}, t; \mathbf{r}', t') = -4\pi \delta^3(\mathbf{r} - \mathbf{r}') \delta(t - t')$$

"Proof" -- Fourier analysis in the time domain -- note that

$$\delta(t - t') = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega e^{i\omega(t-t')}$$

Define:

$$G(\mathbf{r}, t; \mathbf{r}', t') = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega e^{i\omega(t-t')} \tilde{G}(\mathbf{r}, \mathbf{r}', \omega)$$

$$\Rightarrow \left(\nabla^2 + \frac{\omega^2}{c^2} \right) \tilde{G}(\mathbf{r}, \mathbf{r}', \omega) = -4\pi \delta^3(\mathbf{r} - \mathbf{r}')$$

Solution of Maxwell's equations in the Lorentz gauge -- continued

Analysis of the Green's function (continued):

$$\left(\nabla^2 + \frac{\omega^2}{c^2} \right) \tilde{G}(\mathbf{r}, \mathbf{r}', \omega) = -4\pi\delta^3(\mathbf{r} - \mathbf{r}')$$

For the case of isotropic boundary values at infinity:

$$\tilde{G}(\mathbf{r}, \mathbf{r}', \omega) = \tilde{G}(\mathbf{r} - \mathbf{r}', \omega)$$

Further assuming that $\tilde{G}(\mathbf{r} - \mathbf{r}', \omega)$ is isotropic in $|\mathbf{r} - \mathbf{r}'| \equiv R$:

$$\left(\frac{1}{R} \frac{d^2}{dR^2} R + \frac{\omega^2}{c^2} \right) \tilde{G}(\mathbf{r}, \mathbf{r}', \omega) = -4\pi\delta^3(\mathbf{r} - \mathbf{r}')$$

$$\text{Solution: } \tilde{G}(\mathbf{r}, \mathbf{r}', \omega) = \frac{1}{R} e^{\pm i\omega R/c}$$

Solution of Maxwell's equations in the Lorentz gauge -- continued

Analysis of the Green's function (continued):

$$\tilde{G}(\mathbf{r}, \mathbf{r}', \omega) = \frac{1}{|\mathbf{r} - \mathbf{r}'|} e^{\pm i\omega |\mathbf{r} - \mathbf{r}'|/c}$$

$$G(\mathbf{r}, t; \mathbf{r}', t') = \frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega e^{i\omega(t-t')} \tilde{G}(\mathbf{r}, \mathbf{r}', \omega)$$

$$= \frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega e^{i\omega(t-t')} \frac{1}{|\mathbf{r} - \mathbf{r}'|} e^{\pm i\omega |\mathbf{r} - \mathbf{r}'|/c}$$

$$= \frac{1}{|\mathbf{r} - \mathbf{r}'|} \left(\frac{1}{2\pi} \int_{-\infty}^{\infty} d\omega e^{i\omega(t-t' \pm |\mathbf{r} - \mathbf{r}'|/c)} \right)$$

$$= \frac{1}{|\mathbf{r} - \mathbf{r}'|} \delta(t - t' \pm |\mathbf{r} - \mathbf{r}'|/c) = \frac{1}{|\mathbf{r} - \mathbf{r}'|} \delta(t' - t \mp |\mathbf{r} - \mathbf{r}'|/c)$$

Solution of Maxwell's equations in the Lorentz gauge -- continued

$$G(\mathbf{r}, t; \mathbf{r}', t') = \frac{1}{|\mathbf{r} - \mathbf{r}'|} \delta\left(t' - \left(t \pm |\mathbf{r} - \mathbf{r}'| / c\right)\right)$$

Solution for field $\Psi(\mathbf{r}, t)$:

$$\Psi(\mathbf{r}, t) = \Psi_{f=0}(\mathbf{r}, t) +$$

$$\int d^3 r' \int dt' \frac{1}{|\mathbf{r} - \mathbf{r}'|} \delta\left(t' - \left(t - \frac{1}{c} |\mathbf{r} - \mathbf{r}'|\right)\right) f(\mathbf{r}', t')$$

Solution of Maxwell's equations in the Lorentz gauge -- continued

Liènard-Wiechert potentials and fields --

Determination of the scalar and vector potentials for a moving point particle (also see Landau and Lifshitz *The Classical Theory of Fields*, Chapter 8.)

Consider the fields produced by the following source: a point charge q moving on a trajectory $\mathbf{R}_q(t)$.

Charge density: $\rho(\mathbf{r}, t) = q\delta^3(\mathbf{r} - \mathbf{R}_q(t))$

Current density: $\mathbf{J}(\mathbf{r}, t) = q\dot{\mathbf{R}}_q(t)\delta^3(\mathbf{r} - \mathbf{R}_q(t))$, where $\dot{\mathbf{R}}_q(t) \equiv \frac{d\mathbf{R}_q(t)}{dt}$.



Solution of Maxwell's equations in the Lorentz gauge -- continued

$$\Phi(\mathbf{r}, t) = \frac{1}{4\pi\epsilon_0} \iint d^3r' dt' \frac{\rho(\mathbf{r}', t')}{|\mathbf{r} - \mathbf{r}'|} \delta(t' - (t - |\mathbf{r} - \mathbf{r}'|/c))$$

$$\mathbf{A}(\mathbf{r}, t) = \frac{1}{4\pi\epsilon_0 c^2} \iint d^3r' dt' \frac{\mathbf{J}(\mathbf{r}', t')}{|\mathbf{r} - \mathbf{r}'|} \delta(t' - (t - |\mathbf{r} - \mathbf{r}'|/c)).$$

We performing the integrations over first d^3r' and then dt' making use of the fact that for any function of t' ,

$$\int_{-\infty}^{\infty} dt' f(t') \delta(t' - (t - |\mathbf{r} - \mathbf{R}_q(t')|/c)) = \frac{f(t_r)}{1 - \frac{\dot{\mathbf{R}}_q(t_r) \cdot (\mathbf{r} - \mathbf{R}_q(t_r))}{c |\mathbf{r} - \mathbf{R}_q(t_r)|}},$$

where the "retarded time" is defined to be

$$t_r \equiv t - \frac{|\mathbf{r} - \mathbf{R}_q(t_r)|}{c}.$$

Solution of Maxwell's equations in the Lorentz gauge -- continued

Resulting scalar and vector potentials:

$$\Phi(\mathbf{r}, t) = \frac{q}{4\pi\epsilon_0} \frac{1}{R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}},$$

$$\mathbf{A}(\mathbf{r}, t) = \frac{q}{4\pi\epsilon_0 c^2} \frac{\mathbf{v}}{R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}},$$

Notation: $\mathbf{R} \equiv \mathbf{r} - \mathbf{R}_q(t_r)$

$$\mathbf{v} \equiv \dot{\mathbf{R}}_q(t_r),$$

$$t_r \equiv t - \frac{|\mathbf{r} - \mathbf{R}_q(t_r)|}{c}.$$

Comment on Lienard-Wiechert potential results

$$\int_{-\infty}^{\infty} dt' f(t') \delta(t' - (t - |\mathbf{r} - \mathbf{R}_q(t')|/c)) = \frac{f(t_r)}{1 - \frac{\dot{\mathbf{R}}_q(t_r) \cdot (\mathbf{r} - \mathbf{R}_q(t_r))}{c|\mathbf{r} - \mathbf{R}_q(t_r)|}},$$

where the “retarded time” is defined to be

$$t_r \equiv t - \frac{|\mathbf{r} - \mathbf{R}_q(t_r)|}{c}.$$

Note that for any function $F(x)$:

$$\int_{-\infty}^{\infty} F(x) \delta(x - x_0) dx = F(x_0)$$

Now consider a function $p(x)$, for which $p(x_i) = 0$ for $i = 1, 2, \dots$

$$\begin{aligned} \int_{-\infty}^{\infty} F(x) \delta(p(x)) dx &= \int_{-\infty}^{\infty} F(x) \left(\sum_i \delta \left((x - x_i) \frac{dp}{dx} \Big|_{x_i} \right) \right) dx \\ &= \sum_i \frac{F(x_i)}{\left| \frac{dp}{dx} \Big|_{x_i} \right|} \end{aligned}$$

Comment on Lienard-Wiechert potential results -- continued

$$\int_{-\infty}^{\infty} dt' f(t') \delta(t' - (t - |\mathbf{r} - \mathbf{R}_q(t')|/c)) = \frac{f(t_r)}{1 - \frac{\dot{\mathbf{R}}_q(t_r) \cdot (\mathbf{r} - \mathbf{R}_q(t_r))}{c|\mathbf{r} - \mathbf{R}_q(t_r)|}},$$

where the “retarded time” is defined to be

$$t_r \equiv t - \frac{|\mathbf{r} - \mathbf{R}_q(t_r)|}{c}.$$

In this case we have:
$$\int_{-\infty}^{\infty} f(t') \delta(p(t')) dt' = \frac{f(t_r)}{\left| 1 - \frac{\dot{\mathbf{R}}_q(t_r) \cdot (\mathbf{r} - \mathbf{R}_q(t_r))}{c|\mathbf{r} - \mathbf{R}_q(t_r)|} \right|}$$

where:
$$p(t') \equiv t' - \left(t - \frac{|\mathbf{r} - \mathbf{R}_q(t')|}{c} \right)$$

$$\frac{dp(t')}{dt'} = 1 - \frac{\frac{d\mathbf{R}_q(t')}{dt'} \cdot (\mathbf{r} - \mathbf{R}_q(t'))}{c|\mathbf{r} - \mathbf{R}_q(t')|} \equiv 1 - \frac{\dot{\mathbf{R}}_q(t') \cdot (\mathbf{r} - \mathbf{R}_q(t'))}{c|\mathbf{r} - \mathbf{R}_q(t')|}$$

Summary of results for fields due to moving charge – Liénard Wiechert potentials

Resulting scalar and vector potentials:

$$\Phi(\mathbf{r}, t) = \frac{q}{4\pi\epsilon_0} \frac{1}{R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}}$$
$$\mathbf{A}(\mathbf{r}, t) = \frac{q}{4\pi\epsilon_0 c^2} \frac{\mathbf{v}}{R - \frac{\mathbf{v} \cdot \mathbf{R}}{c}}$$

Notation: $\mathbf{R} \equiv \mathbf{r} - \mathbf{R}_q(t_r)$

$$\mathbf{v} \equiv \dot{\mathbf{R}}_q(t_r),$$

$$t_r \equiv t - \frac{|\mathbf{r} - \mathbf{R}_q(t_r)|}{c}.$$