

PHY 712 Electrodynamics

10-10:50 AM MWF Olin 103

Class notes for Lecture 3:

Reading: Chapter 1 (especially 1.11) in JDJ;

- 1. Continued discussion/derivation of Ewald summation methods**
- 2. Example for CsCl**

PHY 712 Electrodynamics

MWF 10-10:50 AM Olin 103 Webpage: <http://www.wfu.edu/~natalie/s23phy712/>

Instructor: [Natalie Holzwarth](#) Office: 300 OPL e-mail: natalie@wfu.edu

Course schedule for Spring 2023

(Preliminary schedule -- subject to frequent adjustment.)

| | Lecture date | JDJ Reading | Topic | HW | Due date |
|---|-----------------|------------------|---|--------------------|------------|
| 1 | Mon: 01/9/2023 | Chap. 1 & Appen. | Introduction, units and Poisson equation | #1 | 01/13/2023 |
| 2 | Wed: 01/11/2023 | Chap. 1 | Electrostatic energy calculations | #2 | 01/18/2023 |
| 3 | Fri: 01/13/2023 | Chap. 1 | Electrostatic energy calculations thanks to Ewald | #3 | 01/18/2023 |
| | Mon: 01/16/2023 | | MLK Holiday -- no class | | |

PHY 712 -- Assignment #3

January 13, 2023

Continue reading Chap. 1 in **Jackson**.

1. Using the Ewald summation methods developed in class, find the electrostatic interaction energy of a NaCl lattice having a cubic lattice constant a . Check that your result does not depend of the Ewald parameter η . You are welcome to copy (and modify) the maple file used in class. A FORTRAN code is also available upon request.

The details are presented in the pdf file Ewaldnotes.pdf

PHY 712 Electrodynamics

MWF 10-10:50 AM Olin 103 Webpage: <http://www.wfu.edu/~natalie/s23phy712/>

Instructor: [Natalie Holzwarth](#) Office: 300 OPL e-mail: natalie@wfu.edu

Lecture Notes

- Lecture 1 -- Introduction and electrostatics [PP slides](#) [PDF](#) [Extra PP](#) [Extra PDF](#)
- Lecture 2 -- Evaluation of electrostatic energy [PP slides](#) [PDF](#) [Ewald notes PDF](#) [Extra PP](#) [Extra PDF](#)
- Lecture 3 -- Ewald summation methods [PP slides](#) [PDF](#) [Ewald notes PDF](#) [CsCI Maple input](#) [CsCI PDF](#)

[Return to main web page for PHY 712](#)

Last modified: Thursday, 12-Jan-2023 11:12:51 EST

Your questions –

From Zezhong -- For CsCl structure graph demonstrations, is there any specific reason that Cl atoms are connected by bonds but Cs atoms are not connected? I found that it was the same for the graph demonstration of CsCl in lecture 2.

Short answer – The bonds are drawn arbitrarily. The notion of bonds has a quantum mechanical origin and Coulombic interactions are very long range. However, the bonds do help with visualizing the structure.

Last time, we argued that we need to carry out the following summation of all pairs of ions using Ewald's very clever trick with the erf and erfc functions:

$$W = \frac{1}{8\pi\epsilon_0} \sum_{i \neq j} \frac{q_i q_j}{|\mathbf{r}_i - \mathbf{r}_j|} = \frac{1}{8\pi\epsilon_0} \left(\sum_{i \neq j} \frac{q_i q_j \operatorname{erf}(\frac{1}{2}\sqrt{\eta}|\mathbf{r}_i - \mathbf{r}_j|)}{|\mathbf{r}_i - \mathbf{r}_j|} + \sum_{i \neq j} \frac{q_i q_j \operatorname{erfc}(\frac{1}{2}\sqrt{\eta}|\mathbf{r}_i - \mathbf{r}_j|)}{|\mathbf{r}_i - \mathbf{r}_j|} \right)$$



Must be evaluated in
“reciprocal” space



Can be converged
as is (in real space)

In order to make progress, we need to systematically enumerate all of the ion positions in the system

Using CsCl as our example system

● Cs


● Cl

Notice that a unit cell contains one Cl and one Cs


$$\mathbf{r}_i = \boldsymbol{\tau}_\alpha + \mathbf{T}$$

for $\alpha = \text{Cl or Cs}$

$$\text{and } \mathbf{T} = n_1 \mathbf{T}_1 + n_2 \mathbf{T}_2 + n_3 \mathbf{T}_3$$



For CsCl --



$$\tau_{\text{Cs}} = 0 \text{ and } \tau_{\text{Cl}} = \frac{a}{2}(\hat{x} + \hat{y} + \hat{z}).$$


$$\mathbf{T}_1 = a\hat{x} \quad \mathbf{T}_2 = a\hat{y} \quad \mathbf{T}_3 = a\hat{z}.$$

Note that in general --

$$\sum_{ij} = N \sum_{\alpha\beta\mathbf{T}}$$

Note that, using $\mathbf{r}_i = \boldsymbol{\tau}_\alpha + \mathbf{T}$

$$W = \frac{1}{8\pi\epsilon_0} \sum_{i \neq j} \frac{q_i q_j}{|\mathbf{r}_i - \mathbf{r}_j|} = \frac{N}{8\pi\epsilon_0} \sum_{\alpha, \beta, \mathbf{T}} \frac{q_\alpha q_\beta}{|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|}$$




omit term $\alpha = \beta$

when $\mathbf{T}=0$

$$\boldsymbol{\tau}_{\alpha\beta} \equiv \boldsymbol{\tau}_\alpha - \boldsymbol{\tau}_\beta$$

$$\frac{W}{N} = \frac{1}{8\pi\epsilon_0} \sum'_{\alpha, \beta, \mathbf{T}} \frac{q_\alpha q_\beta}{|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|}$$



Note – this works for CsCl and more generally

$$\frac{W}{N} = \frac{1}{8\pi\epsilon_0} \sum_{\alpha, \beta, \mathbf{T}} \frac{q_\alpha q_\beta}{|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|} = w_1 + w_2$$

$$w_2 \equiv \frac{1}{8\pi\epsilon_0} \sum_{\alpha, \beta, \mathbf{T}} \frac{q_\alpha q_\beta \operatorname{erfc}\left(\frac{1}{2}\sqrt{\eta}|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|\right)}{|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|} \equiv \frac{e^2}{8\pi\epsilon_0} \mathcal{T}_2$$

For CsCl:

$$\tau_{\text{Cs}} = 0 \text{ and } \tau_{\text{Cl}} = \frac{a}{2}(\hat{x} + \hat{y} + \hat{z}).$$

$$q_{\text{Cs}} = e$$

$$q_{\text{Cl}} = -e$$

$$\mathbf{T}_1 = a\hat{x} \quad \mathbf{T}_2 = a\hat{y} \quad \mathbf{T}_3 = a\hat{z}.$$

$$\mathcal{T}_2 \equiv \sum_{\mathbf{T} \neq \mathbf{0}} 2 \frac{\operatorname{erfc}\left(\frac{1}{2}\sqrt{\eta}|\mathbf{T}|\right)}{|\mathbf{T}|} - \sum_{\mathbf{T}} 2 \frac{\operatorname{erfc}\left(\frac{1}{2}\sqrt{\eta}|\tau_{\text{Cl}} + \mathbf{T}|\right)}{|\tau_{\text{Cl}} + \mathbf{T}|}.$$

For evaluating w_1 which needs to be evaluated in reciprocal space, additional considerations are needed --

It can be shown that --

$$\sum_{\mathbf{T}} \delta^3(\mathbf{r} - \mathbf{T}) = \frac{1}{\Omega} \sum_{\mathbf{G}} e^{i\mathbf{G}\cdot\mathbf{r}},$$



Volume of unit cell
in real space

In this discussion, will assume we have a 3-dimensional periodic system. It can be easily generalized to 1- or 2- dimensional systems. In general, a translation vector can be described a linear combination of the three primitive translation vectors \mathbf{T}_1 , \mathbf{T}_2 , and \mathbf{T}_3 :

$$\mathbf{T} = n_1\mathbf{T}_1 + n_2\mathbf{T}_2 + n_3\mathbf{T}_3, \quad (12)$$

where $\{n_1, n_2, n_3\}$ are integers. Note that the unit cell volume Ω can be expressed in terms of the primitive translation vectors according to:

$$\Omega = |\mathbf{T}_1 \cdot (\mathbf{T}_2 \times \mathbf{T}_3)|. \quad (13)$$

Preview – what happens if $\sum_{\mathbf{T}} \delta^3(\mathbf{r} - \mathbf{T}) = \frac{1}{\Omega} \sum_{\mathbf{G}} e^{i\mathbf{G}\cdot\mathbf{r}}$,
is true?

Recall:
$$\frac{W}{N} = \frac{1}{8\pi\epsilon_0} \sum_{\alpha,\beta,\mathbf{T}} \frac{q_\alpha q_\beta}{|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|} = w_1 + w_2$$

$$w_1 \equiv \frac{1}{8\pi\epsilon_0} \sum_{\alpha,\beta,\mathbf{T}} \frac{q_\alpha q_\beta \operatorname{erf}\left(\frac{1}{2}\sqrt{\eta}|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|\right)}{|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|}$$

$$= \frac{1}{8\pi\epsilon_0} \left(\sum_{\alpha,\beta,\mathbf{T}} \frac{q_\alpha q_\beta \operatorname{erf}\left(\frac{1}{2}\sqrt{\eta}|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|\right)}{|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|} - \sqrt{\frac{\eta}{\pi}} \sum_{\alpha} q_\alpha^2 \right)$$

Subtract out self-energy
contribution

Continued --

Preview – what happens if
is true?

$$\sum_{\mathbf{T}} \delta^3(\mathbf{r} - \mathbf{T}) = \frac{1}{\Omega} \sum_{\mathbf{G}} e^{i\mathbf{G}\cdot\mathbf{r}},$$

$$\sum_{\mathbf{T}} \frac{\text{erf}(\frac{1}{2}\sqrt{\eta}|\tau_{\alpha\beta} + \mathbf{T}|)}{|\tau_{\alpha\beta} + \mathbf{T}|} = \int d^3r \sum_{\mathbf{T}} \delta^3(\mathbf{r} - \mathbf{T}) \frac{\text{erf}(\frac{1}{2}\sqrt{\eta}|\tau_{\alpha\beta} + \mathbf{r}|)}{|\tau_{\alpha\beta} + \mathbf{r}|}.$$

This becomes,

$$\frac{1}{\Omega} \sum_{\mathbf{G}} \int d^3r e^{i\mathbf{G}\cdot\mathbf{r}} \frac{\text{erf}(\frac{1}{2}\sqrt{\eta}|\tau_{\alpha\beta} + \mathbf{r}|)}{|\tau_{\alpha\beta} + \mathbf{r}|} = \frac{4\pi}{\Omega} \left(\sum_{\mathbf{G}\neq\mathbf{0}} \frac{e^{-i\mathbf{G}\cdot\tau_{\alpha\beta}} e^{-G^2/\eta^2}}{G^2} + \frac{1}{2} \int_0^{\frac{1}{2}\sqrt{\eta}} \frac{du}{u^3} \right),$$

where the last term, which is infinite, comes from the $\mathbf{G} = \mathbf{0}$ contribution.

Back to “proof” --

The reciprocal lattice vectors \mathbf{G} can generally be written as a linear combination of the three primitive reciprocal lattice vectors \mathbf{G}_1 , \mathbf{G}_2 , and \mathbf{G}_3 :

$$\mathbf{G} = m_1 \mathbf{G}_1 + m_2 \mathbf{G}_2 + m_3 \mathbf{G}_3, \quad (14)$$

where $\{m_1, m_2, m_3\}$ are integers. The primitive reciprocal lattice vectors are determined from the primitive translation vectors according to the identities:

$$\mathbf{G}_i \cdot \mathbf{T}_j = 2\pi \delta_{ij}. \quad (15)$$

Note that the “volume” of the primitive reciprocal lattice is given by

$$|\mathbf{G}_1 \cdot (\mathbf{G}_2 \times \mathbf{G}_3)| = \frac{(2\pi)^3}{\Omega}. \quad (16)$$

CsCl structure

There are two kinds of sites – $\tau_{\text{Cs}} = 0$ and $\tau_{\text{Cl}} = \frac{a}{2}(\hat{x} + \hat{y} + \hat{z})$.

Example --


$$\mathbf{T}_1 = a\hat{x} \quad \mathbf{T}_2 = a\hat{y} \quad \mathbf{T}_3 = a\hat{z}.$$

$$\mathbf{G}_1 = \frac{2\pi}{a}\hat{x} \quad \mathbf{G}_2 = \frac{2\pi}{a}\hat{y} \quad \mathbf{G}_3 = \frac{2\pi}{a}\hat{z}.$$

Consider the summation;

$$\sum_{k=-M}^{+M} e^{ik(\mathbf{G}_1 \cdot \mathbf{r})} = \frac{\sin\left(\left(M + \frac{1}{2}\right)\mathbf{G}_1 \cdot \mathbf{r}\right)}{\sin(\mathbf{G}_1 \cdot \mathbf{r}/2)}.$$

Behavior of function for various values of M :



$x = \mathbf{G}_1 \cdot \mathbf{r}$ $\sum_{k=-M}^M e^{ikx}$ has large peaks at $x = \text{multiples of } 2\pi$

“Proof” of identity --

Consider the geometric series

$$\sum_{k=-M}^{+M} e^{ik(\mathbf{G}_1 \cdot \mathbf{r})} = \frac{\sin\left(\left(M + \frac{1}{2}\right)\mathbf{G}_1 \cdot \mathbf{r}\right)}{\sin(\mathbf{G}_1 \cdot \mathbf{r}/2)}. \quad (17)$$

The behavior of the right hand side of Eq. (17) is that it is small in magnitude very except when the denominator vanishes. This occurs whenever $\mathbf{G}_1 \cdot \mathbf{r}/2 = n_1\pi$, where n_1 represents any integer. If we take the limit $M \rightarrow \infty$, we find that the function represents the behavior of a sum of delta functions:

$$\lim_{M \rightarrow \infty} \frac{\sin\left(\left(M + \frac{1}{2}\right)\mathbf{G}_1 \cdot \mathbf{r}\right)}{\sin(\mathbf{G}_1 \cdot \mathbf{r}/2)} = 2\pi \sum_{n_1} \delta(\mathbf{G}_1 \cdot (\mathbf{r} - n_1 \mathbf{T}_1)). \quad (18)$$

The summation over all lattice translations $n_1 \mathbf{T}_1$ is due to the fact that $\sin(\mathbf{G}_1 \cdot \mathbf{r}/2) = 0$ whenever $\mathbf{r} = n_1 \mathbf{T}_1$. Carrying out the geometric summations in the right hand side of Eq. 7 for all three reciprocal lattice vectors and taking the limit as in Eq. 18,

$$\sum_{\mathbf{G}} e^{i\mathbf{G} \cdot \mathbf{r}} = (2\pi)^3 \sum_{n_1, n_2, n_3} \delta(\mathbf{G}_1 \cdot (\mathbf{r} - n_1 \mathbf{T}_1)) \delta(\mathbf{G}_2 \cdot (\mathbf{r} - n_2 \mathbf{T}_2)) \delta(\mathbf{G}_3 \cdot (\mathbf{r} - n_3 \mathbf{T}_3)). \quad (19)$$

Final result --

$$\sum_{\mathbf{G}} e^{i\mathbf{G}\cdot\mathbf{r}} = \frac{(2\pi)^3}{|\mathbf{G}_1 \cdot (\mathbf{G}_2 \times \mathbf{G}_3)|} \sum_{\mathbf{T}} \delta^3(\mathbf{r} - \mathbf{T}) = \Omega \sum_{\mathbf{T}} \delta^3(\mathbf{r} - \mathbf{T})$$

$$\sum_{\mathbf{T}} \delta^3(\mathbf{r} - \mathbf{T}) = \frac{1}{\Omega} \sum_{\mathbf{G}} e^{i\mathbf{G}\cdot\mathbf{r}},$$

$$\frac{W}{N} = \frac{1}{8\pi\epsilon_0} \sum_{\alpha, \beta, \mathbf{T}} \frac{q_\alpha q_\beta}{|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|} = w_1 + w_2$$

$$w_1 \equiv \frac{1}{8\pi\epsilon_0} \sum_{\alpha, \beta, \mathbf{T}} \frac{q_\alpha q_\beta \operatorname{erf}\left(\frac{1}{2}\sqrt{\eta}|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|\right)}{|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|} \equiv \frac{e^2}{8\pi\epsilon_0} \mathcal{I}_1$$

$$\sum_{\alpha, \beta, \mathbf{T}} \frac{q_\alpha q_\beta \operatorname{erf}\left(\frac{1}{2}\sqrt{\eta}|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|\right)}{|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|}$$

$$= \sum_{\alpha, \beta} \left(\sum_{\mathbf{T}} \frac{q_\alpha q_\beta \operatorname{erf}\left(\frac{1}{2}\sqrt{\eta}|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|\right)}{|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|} - \delta_{\alpha\beta} q_\alpha^2 \lim_{x \rightarrow 0} \left(\frac{\operatorname{erf}\left(\frac{1}{2}\sqrt{\eta}x\right)}{x} \right) \right)$$

$$\lim_{x \rightarrow 0} \left(\frac{\operatorname{erf}\left(\frac{1}{2}\sqrt{\eta}x\right)}{x} \right) = \sqrt{\frac{\eta}{\pi}}$$

$$\sum_{\mathbf{T}} \frac{\text{erf}(\frac{1}{2}\sqrt{\eta}|\tau_{\alpha\beta} + \mathbf{T}|)}{|\tau_{\alpha\beta} + \mathbf{T}|} = \int d^3r \sum_{\mathbf{T}} \delta^3(\mathbf{r} - \mathbf{T}) \frac{\text{erf}(\frac{1}{2}\sqrt{\eta}|\tau_{\alpha\beta} + \mathbf{r}|)}{|\tau_{\alpha\beta} + \mathbf{r}|}$$

$$\frac{1}{\Omega} \sum_{\mathbf{G}} \int d^3r e^{i\mathbf{G}\cdot\mathbf{r}} \frac{\text{erf}(\frac{1}{2}\sqrt{\eta}|\tau_{\alpha\beta} + \mathbf{r}|)}{|\tau_{\alpha\beta} + \mathbf{r}|} =$$

$$\frac{4\pi}{\Omega} \left(\sum_{\mathbf{G} \neq \mathbf{0}} \frac{e^{-i\mathbf{G}\cdot\tau_{\alpha\beta}} e^{-G^2/\eta^2}}{G^2} + \frac{1}{2} \int_0^{\frac{1}{2}\sqrt{\eta}} \frac{du}{u^3} \right)$$



divergent results

Derivation includes :

$$\text{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt$$

+ appropriate changes of variables
and order of integration

Summary --

$$\sum_{\alpha, \beta, \mathbf{T}} \frac{q_\alpha q_\beta \operatorname{erf}\left(\frac{1}{2} \sqrt{\eta} |\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|\right)}{|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|}$$

$$= \sum_{\alpha, \beta} \left(\sum_{\mathbf{T}} \frac{q_\alpha q_\beta \operatorname{erf}\left(\frac{1}{2} \sqrt{\eta} |\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|\right)}{|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|} - \delta_{\alpha\beta} q_\alpha^2 \sqrt{\frac{\eta}{\pi}} \right)$$

$$= \sum_{\alpha, \beta} q_\alpha q_\beta \left(\frac{4\pi}{\Omega} \sum_{\mathbf{G} \neq 0} \frac{e^{-i\mathbf{G} \cdot \boldsymbol{\tau}_{\alpha\beta}} e^{-G^2/\eta}}{G^2} + \frac{4\pi}{\Omega} \int_0^{\frac{1}{2}\sqrt{\eta}} \frac{du}{u^3} - \delta_{\alpha\beta} q_\alpha^2 \sqrt{\frac{\eta}{\pi}} \right)$$



Divergent term

Note that for a neutral system

$\sum_{\alpha, \beta} q_\alpha q_\beta = 0$ and divergent term vanishes

Problem – Electrostatic energy of a periodic non-neutral system is not defined!

Solution -- Add uniform compensating charge and find the electrostatic energy of periodic neutral system, carefully analyzing diverging terms to determine a convergent expression.

$$\frac{W}{N} = \sum_{\alpha\beta} \frac{q_\alpha q_\beta}{8\pi\epsilon_0} \left(\frac{4\pi}{\Omega} \sum_{\mathbf{G} \neq \mathbf{0}} \frac{e^{-i\mathbf{G} \cdot \boldsymbol{\tau}_{\alpha\beta}} e^{-G^2/\eta}}{G^2} - \sqrt{\frac{\eta}{\pi}} \delta_{\alpha\beta} + \sum_{\mathbf{T}}' \frac{\text{erfc}(\frac{1}{2}\sqrt{\eta}|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|)}{|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|} \right) - \frac{4\pi Q^2}{8\pi\epsilon_0\Omega\eta}$$

Where $Q \equiv \sum_{\alpha} q_{\alpha}$

Some details

$$\frac{W'}{N} = \frac{1}{8\pi\epsilon_0} \left(\sum_{\alpha,\beta,\mathbf{T}} \frac{q_\alpha q_\beta}{|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|} - \sum_\alpha q_\alpha \frac{Q}{\Omega} 4\pi \int_0^\infty \frac{r^2}{r} dr \right)$$

$$- \sum_\alpha q_\alpha \frac{Q}{\Omega} 4\pi \int_0^\infty \frac{r^2}{r} dr = -\frac{4\pi Q^2}{\Omega} \int_0^\infty r dr = -\frac{4\pi Q^2}{\Omega} \int_0^\infty \frac{du}{u^3}$$

$$= -\frac{4\pi Q^2}{\Omega} \left(\int_0^{\sqrt{\eta}/2} \frac{du}{u^3} + \int_{\sqrt{\eta}/2}^\infty \frac{du}{u^3} \right)$$


$$= -\frac{4\pi Q^2}{\Omega} \left(\int_0^{\sqrt{\eta}/2} \frac{du}{u^3} + \frac{2}{\eta} \right)$$


Cancels
divergence

Extra
contribution

$$\frac{W}{N} = \sum_{\alpha\beta} \frac{q_\alpha q_\beta}{8\pi\epsilon_0} \left(\frac{4\pi}{\Omega} \sum_{\mathbf{G} \neq 0} \frac{e^{-i\mathbf{G} \cdot \boldsymbol{\tau}_{\alpha\beta}} e^{-G^2/\eta}}{G^2} - \sqrt{\frac{\eta}{\pi}} \delta_{\alpha\beta} + \sum'_{\mathbf{T}} \frac{\text{erfc}(\frac{1}{2}\sqrt{\eta}|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|)}{|\boldsymbol{\tau}_{\alpha\beta} + \mathbf{T}|} \right) - \frac{4\pi Q^2}{8\pi\epsilon_0 \Omega \eta}$$

For CsCl --



$$\tau_{Cs} = 0 \text{ and } \tau_{Cl} = \frac{a}{2}(\hat{x} + \hat{y} + \hat{z}).$$

$$\mathbf{T}_1 = a\hat{x} \quad \mathbf{T}_2 = a\hat{y} \quad \mathbf{T}_3 = a\hat{z}.$$

$$q_{Cs} = e$$

$$q_{Cl} = -e$$

$$\frac{W}{N} = \sum_{\alpha\beta} \frac{q_\alpha q_\beta}{8\pi\epsilon_0} \left(\frac{4\pi}{\Omega} \sum_{\mathbf{G} \neq \mathbf{0}} \frac{e^{-i\mathbf{G} \cdot \tau_{\alpha\beta}} e^{-G^2/\eta}}{G^2} - \sqrt{\frac{\eta}{\pi}} \delta_{\alpha\beta} + \sum_{\mathbf{T}}' \frac{\text{erfc}(\frac{1}{2}\sqrt{\eta}|\tau_{\alpha\beta} + \mathbf{T}|)}{|\tau_{\alpha\beta} + \mathbf{T}|} \right) - \frac{4\pi Q^2}{8\pi\epsilon_0\Omega\eta}$$

$$\frac{W}{N} = \frac{e^2}{8\pi\epsilon_0} (\mathcal{T}_1 + \mathcal{T}_2)$$

$$\mathcal{T}_1 \equiv \frac{4\pi}{\Omega} \sum_{\mathbf{G} \neq \mathbf{0}} 2 \frac{(1 - e^{-i\mathbf{G} \cdot \tau_{Cl}}) e^{-|\mathbf{G}|^2/\eta}}{|\mathbf{G}|^2} - 2\sqrt{\frac{\eta}{\pi}}$$

$$\mathcal{T}_2 \equiv \sum_{\mathbf{T} \neq \mathbf{0}} 2 \frac{\text{erfc}(\frac{1}{2}\sqrt{\eta}|\mathbf{T}|)}{|\mathbf{T}|} - \sum_{\mathbf{T}} 2 \frac{\text{erfc}(\frac{1}{2}\sqrt{\eta}|\tau_{Cl} + \mathbf{T}|)}{|\tau_{Cl} + \mathbf{T}|}.$$

➔ Can be evaluated using Maple or Mathematica or other programming language

Using Maple example, we find that for a CsCl lattice constant a :

$$\frac{W}{N} = \frac{e^2}{8\pi\epsilon_0 a} (-4.0707)$$

This result is consistent with the Madelung constant.