



PHY 712 Electrodynamics

10-10:50 AM MWF in Olin 103

Class notes for Lecture 5:

Reading: Chapter 1 - 3 in JDJ

Electrostatic potentials

- 1. One, two, and three dimensions
(Cartesian coordinates)**
- 2. Mean value theorem for the
electrostatic potential**

PHY 712 Electrodynamics

MWF 10-10:50 AM Olin 103 Webpage: <http://www.wfu.edu/~natalie/s24phy712/>

Instructor: [Natalie Holzwarth](#) Office:300 OPL e-mail:natalie@wfu.edu

Course schedule for Spring 2024

(Preliminary schedule -- subject to frequent adjustment.)

	Lecture date	JDJ Reading	Topic	HW	Due date
1	Wed: 01/17/2024	Chap. 1 & Appen.	Introduction, units and Poisson equation	#1	01/19/2024
2	Fri: 01/19/2024	Chap. 1	Electrostatic energy calculations	#2	01/29/2024
3	Mon: 01/22/2024	Chap. 1	Electrostatic energy calculations	#3	01/29/2024
4	Wed: 01/24/2024	Chap. 1 & 2	Electrostatic potentials and fields	#4	01/29/2024
5	Fri: 01/26/2024	Chap. 1 - 3	Poisson's equation in 2 and 3 dimensions	#5	01/29/2024
6	Mon: 01/29/2024	Chap. 1 - 3	Brief introduction to numerical methods		

PHY 712 – Problem Set #5

Assigned: 01/26/2024

Due: 01/29/2024

Continue reading Chapter 1 & 2 in **Jackson**

1. Consider a one-dimensional charge distribution of the form:

$$\rho(x) = \begin{cases} 0 & \text{for } x < -a \\ \rho_0 \sin(\pi x/a) & \text{for } -a \leq x \leq a \\ 0 & \text{for } x > a, \end{cases}$$

where ρ_0 and a are constants.

- Solve the Poisson equation for the electrostatic potential $\Phi(x)$ with the boundary conditions $\Phi(-a) = 0$ and $\frac{d\Phi}{dx}(a) = 0$.
- Find the corresponding electrostatic field $E(x)$.
- Plot $\Phi(x)$ and $E(x)$.
- Discuss your results in terms of possible electronic devices.



Poisson Equation

$$\nabla^2 \Phi_P(\mathbf{r}) = -\frac{\rho(\mathbf{r})}{\epsilon_0}$$

Solution to Poisson equation using Green's function $G(\mathbf{r}, \mathbf{r}')$:

$$\Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V d^3 r' \rho(\mathbf{r}') G(\mathbf{r}, \mathbf{r}') +$$
$$\frac{1}{4\pi} \int_S d^2 r' [G(\mathbf{r}, \mathbf{r}') \nabla' \Phi(\mathbf{r}') - \Phi(\mathbf{r}') \nabla' G(\mathbf{r}, \mathbf{r}')] \cdot \hat{\mathbf{r}}'.$$

Poisson equation for one-dimensional system

$$\frac{d^2\Phi_P(x)}{dx^2} = -\frac{\rho(x)}{\epsilon_0}$$

Example solution (assuming the boundary values vanish):

$$\Phi_P(x) = \frac{1}{4\pi\epsilon_0} \int_{-\infty}^{\infty} G(x, x') \rho(x') dx' + C_1 + C_2 x$$

where $G(x, x') = 4\pi x_{<}^{-1}$ where $x_{<}$ is the smaller of x and x' ;
 C_1 and C_2 are constants.

Check:

$$\Phi_P(x) = \frac{1}{\epsilon_0} \left\{ \int_{-\infty}^x x' \rho(x') dx' + x \int_x^{\infty} \rho(x') dx' \right\} + C_1 + C_2 x$$

$$\frac{d\Phi_P(x)}{dx} = \frac{1}{\epsilon_0} \int_x^{\infty} \rho(x') dx' + C_2 \quad \Rightarrow \frac{d^2\Phi_P(x)}{dx^2} = -\frac{\rho(x)}{\epsilon_0}$$

Note that

$$\begin{aligned} & \frac{d}{dx} \left(\int_{A(x)}^{B(x)} f(x, x') dx' \right) \\ &= f(x, B(x)) \frac{dB(x)}{dx} - f(x, A(x)) \frac{dA(x)}{dx} + \int_{A(x)}^{B(x)} \frac{\partial f(x, x')}{\partial x} dx' \end{aligned}$$

Question

Example solution:

Why these extra terms?



$$\Phi_P(x) = \frac{1}{4\pi\epsilon_0} \int_{-\infty}^{\infty} G(x, x') \rho(x') dx' + C_1 + C_2 x$$

where $G(x, x') = 4\pi x_{<}$ where $x_{<}$ is the smaller of x and x' ;
 C_1 and C_2 are constants.

$$\frac{d^2 \Phi_P(x)}{dx^2} = -\frac{\rho(x)}{\epsilon_0} ?$$

Checking:

$$\frac{d^2 C_1}{dx^2} = 0 \quad \frac{d^2 C_2 x}{dx^2} = 0$$

$$\frac{d^2 G(x, x')}{dx^2} = -4\pi\delta(x - x')$$

$$\begin{aligned} \frac{d^2}{dx^2} \int_{-\infty}^{\infty} G(x, x') \rho(x') dx' \\ &= -4\pi \int_{-\infty}^{\infty} \delta(x - x') \rho(x') dx' \\ &= -4\pi \rho(x) \end{aligned}$$



General procedure for constructing Green's function for one-dimensional system using 2 independent solutions of the homogeneous equations

Consider two independent solutions to the homogeneous equation

$$\nabla^2 \phi_i(x) = 0$$

where $i = 1$ or 2 . Let

$$G(x, x') = \frac{4\pi}{W} \phi_1(x_{<}) \phi_2(x_{>}).$$

This notation means that $x_{<}$ should be taken as the smaller of x and x' and $x_{>}$ should be taken as the larger.

"Wronskian": $W \equiv \frac{d\phi_1(x)}{dx} \phi_2(x) - \phi_1(x) \frac{d\phi_2(x)}{dx}.$

Beautiful method; but only works in one dimension.



Orthogonal function expansions and Green's functions

Suppose we have a “complete” set of orthogonal functions $\{u_n(x)\}$ defined in the interval $x_1 \leq x \leq x_2$ such that

$$\int_{x_1}^{x_2} u_n(x) u_m(x) dx = \delta_{nm}.$$

We can show that the completeness of this functions implies that

$$\sum_{n=1}^{\infty} u_n(x) u_n(x') = \delta(x - x').$$

This relation allows us to use these functions to represent a Green’s function for our system. For the 1-dimensional Poisson equation, the Green’s function satisfies

$$\frac{\partial^2}{\partial x^2} G(x, x') = -4\pi \delta(x - x').$$

Orthogonal function expansion -- continued

Suppose the orthogonal functions satisfy an eigenvalue equation:

$$\frac{d^2}{dx^2} u_n(x) = -\alpha_n u_n(x)$$

where the functions $u_n(x)$ also satisfy the appropriate boundary conditions, then we can construct the Green's function:

$$G(x, x') = 4\pi \sum_n \frac{u_n(x)u_n(x')}{\alpha_n}.$$

Check:

$$\begin{aligned} \frac{d^2}{dx^2} G(x, x') &= 4\pi \sum_n \frac{(-\alpha_n u_n(x))u_n(x')}{\alpha_n} = -4\pi \sum_n u_n(x)u_n(x') \\ &= -4\pi \delta(x - x') \end{aligned}$$

Example

For example, consider the previous example in the interval

$-a \leq x \leq a$:

$$\rho(x) = \begin{cases} 0 & \text{for } x < -a \\ -\rho_0 & \text{for } -a < x < 0 \\ +\rho_0 & \text{for } 0 < x < a \\ 0 & \text{for } x > a \end{cases}$$

We want to solve the Poisson equation with boundary condition

$\Phi(-a) = 0$ and $d\Phi(-a)/dx = 0$. We may choose

$u_n(x) = \sqrt{\frac{1}{a}} \sin\left(\frac{[2n+1]\pi x}{2a}\right)$ and the corresponding Green's function

$$G(x, x') = \frac{4\pi}{a} \sum_{n=0}^{\infty} \frac{\sin\left(\frac{[2n+1]\pi x}{2a}\right) \sin\left(\frac{[2n+1]\pi x'}{2a}\right)}{\left(\frac{[2n+1]\pi}{2a}\right)^2}.$$

Note that this is a convenient choice, but not necessarily compatible with the boundary values.

Example -- continued

This form of the one-dimensional Green's function only allows us to find a solution to the Poisson equation within the interval $-a \leq x \leq a$ from

$$\Phi(x) = \frac{1}{4\pi\epsilon_0} \int_{-a}^a dx' G(x, x') \rho(x') + C_1$$
$$\Rightarrow \Phi(x) = \frac{\rho_0 a^2}{\epsilon_0} \left(16 \sum_{n=0}^{\infty} \frac{\sin\left(\frac{[2n+1]\pi x}{2a}\right)}{([2n+1]\pi)^3} + \frac{1}{2} \right),$$

choosing C_1 so that $\Phi(-a) = 0$.

Exact result: $\Phi(x) = \begin{cases} 0 & \text{for } x < -a \\ \frac{\rho_0}{2\epsilon_0} (x+a)^2 & \text{for } -a < x < 0 \\ -\frac{\rho_0}{2\epsilon_0} (x-a)^2 + \frac{\rho_0 a^2}{\epsilon_0} & \text{for } 0 < x < a \\ \frac{\rho_0}{\epsilon_0} a^2 & \text{for } x > a \end{cases}$


Some details --

$$\rho(x) = \begin{cases} 0 & \text{for } x < -a \\ -\rho_0 & \text{for } -a < x < 0 \\ +\rho_0 & \text{for } 0 < x < a \\ 0 & \text{for } x > a \end{cases}$$

$$G(x, x') = \frac{4\pi}{a} \sum_{n=0}^{\infty} \frac{\sin\left(\frac{[2n+1]\pi x}{2a}\right) \sin\left(\frac{[2n+1]\pi x'}{2a}\right)}{\left(\frac{[2n+1]\pi}{2a}\right)^2}$$

$$\frac{1}{4\pi\epsilon_0} \int_{-a}^a dx' G(x, x') \rho(x') = \frac{\rho_0 a^2}{\epsilon_0} \left(16 \sum_{n=0}^{\infty} \frac{\sin\left(\frac{[2n+1]\pi x}{2a}\right)}{([2n+1]\pi)^3} \right); \quad S(x, N) \equiv 16 \sum_{n=0}^N \frac{\sin\left(\frac{[2n+1]\pi x}{2a}\right)}{([2n+1]\pi)^3}$$

N~






Example -- continued

$$\Phi(x) = \frac{\rho_0 a^2}{\epsilon_0} \left(16 \sum_{n=0}^{\infty} \frac{\sin\left(\frac{[2n+1]\pi x}{2a}\right)}{([2n+1]\pi)^3} + \frac{1}{2} \right)$$

Needed for boundary values.





Orthogonal function expansions in 2 and 3 dimensions – for cartesian coordinates:

$$\nabla^2 \Phi(\mathbf{r}) \equiv \frac{\partial^2 \Phi(\mathbf{r})}{\partial x^2} + \frac{\partial^2 \Phi(\mathbf{r})}{\partial y^2} + \frac{\partial^2 \Phi(\mathbf{r})}{\partial z^2} = -\rho(\mathbf{r}) / \epsilon_0.$$

Let $\{u_n(x)\}$, $\{v_n(y)\}$, $\{w_n(z)\}$ denote complete orthogonal function sets in the x , y , and z dimensions, respectively. The Green's function construction becomes:

$$G(x, x', y, y', z, z') = 4\pi \sum_{lmn} \frac{u_l(x)u_l(x')v_m(y)v_m(y')w_n(z)w_n(z')}{\alpha_l + \beta_m + \gamma_n},$$

where


$$\frac{d^2}{dx^2} u_l(x) = -\alpha_l u_l(x), \quad \frac{d^2}{dy^2} v_m(y) = -\beta_m v_m(y), \quad \text{and} \quad \frac{d^2}{dz^2} w_n(z) = -\gamma_n w_n(z).$$

(See Eq. 3.167 in Jackson for example.)



Details of a two-dimensional example --

Example:



Two dimensional box with sides a and b with boundary conditions: $\Phi(0,y)=\Phi(a,y)=\Phi(x,0)=\Phi(x,b)=0$

$$\nabla^2 \Phi(\mathbf{r}) \equiv \frac{\partial^2 \Phi(\mathbf{r})}{\partial x^2} + \frac{\partial^2 \Phi(\mathbf{r})}{\partial y^2} = -\rho(\mathbf{r})/\epsilon_0.$$


$$G(x, x', y, y') = 4\pi \sum_{lm} \frac{u_l(x)u_l(x')v_m(y)v_m(y')}{\alpha_l + \beta_m},$$

$$\text{where } \frac{d^2}{dx^2}u_l(x) = -\alpha_l u_l(x), \quad \frac{d^2}{dy^2}v_m(y) = -\beta_m v_m(y)$$



Two dimensional example continued --

b




$$u_l(x) = \sqrt{\frac{2}{a}} \sin\left(\frac{l\pi x}{a}\right) \quad v_m(y) = \sqrt{\frac{2}{b}} \sin\left(\frac{m\pi y}{b}\right) \text{ with } \alpha_l = \left(\frac{l\pi}{a}\right)^2 \quad \beta_m = \left(\frac{m\pi}{b}\right)^2$$

$$\begin{aligned} G(x, x', y, y') &= 4\pi \sum_{lm} \frac{u_l(x)u_l(x')v_m(y)v_m(y')}{\alpha_l + \beta_m} \\ &= \frac{16}{\pi ab} \sum_{lm} \frac{\sin\left(\frac{l\pi x}{a}\right)\sin\left(\frac{l\pi x'}{a}\right)\sin\left(\frac{m\pi y}{b}\right)\sin\left(\frac{m\pi y'}{b}\right)}{\left(\frac{l}{a}\right)^2 + \left(\frac{m}{b}\right)^2} \end{aligned}$$



Example two-dimensional system continued -- Two



dimensional box
with sides a and
 b with boundary
conditions:

$$\Phi(0,y)=\Phi(a,y)=0$$
$$\Phi(x,0)=\Phi(x,b)=0$$

$$\Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V d^3 r' \rho(\mathbf{r}') G(\mathbf{r}, \mathbf{r}') +$$

Don't know this term

$$\frac{1}{4\pi} \int_S d^2 r' [G(\mathbf{r}, \mathbf{r}') \nabla' \Phi(\mathbf{r}') - \Phi(\mathbf{r}') \nabla' G(\mathbf{r}, \mathbf{r}')] \cdot \hat{\mathbf{r}}'.$$

Know this term=0

→ In this case it is prudent to design $G(\mathbf{r}, \mathbf{r}')$ to vanishes on boundary and the surface integral is trivial.



The integral that is needed in this case:

$$\Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V d^3 r' \rho(\mathbf{r}') G(\mathbf{r}, \mathbf{r}')$$

For example:

$$G(x, x', y, y') = \frac{16}{\pi ab} \sum_{lm} \frac{\sin\left(\frac{l\pi x}{a}\right) \sin\left(\frac{l\pi x'}{a}\right) \sin\left(\frac{m\pi y}{b}\right) \sin\left(\frac{m\pi y'}{b}\right)}{\left(\frac{l}{a}\right)^2 + \left(\frac{m}{b}\right)^2}$$

Note that in this case, the eigenfunctions are compatible with the given boundary values.

Example charge densities:

Example #1: $\rho(x, y) = \rho_0 \sin\left(\frac{\pi x}{a}\right) \sin\left(\frac{\pi y}{b}\right)$

Example #2: $\rho(x, y) = \rho_0$

Worked out in
textbook.

Combined orthogonal function expansion and homogeneous solution construction of Green's function in 2 and 3 dimensions.

An alternative method of finding Green's functions for a second order ordinary differential equations (in 1 dimension) is based on a product of two independent solutions of the homogeneous equation, $\phi_1(x)$ and $\phi_2(x)$:

$$G(x, x') = K \phi_1(x_{<}) \phi_2(x_{>}), \text{ where } K \equiv \frac{4\pi}{\frac{d\phi_1}{dx} \phi_2 - \phi_1 \frac{d\phi_2}{dx}},$$

where $x_{<}$ denotes the smaller of x and x' .

For the two and three dimensional cases, we can use this technique in one of the dimensions in order to reduce the number of summation terms. These ideas are discussed in Section 3.11 of Jackson.



Green's function construction -- continued

For the two dimensional case, for example, we can assume that the Green's function can be written in the form:

$$G(x, x', y, y') = \sum_n u_n(x) u_n(x') g_n(y, y') \text{ where } \frac{d^2}{dx^2} u_n(x) = -\alpha_n u_n(x)$$

The y dependence of this equation will have the required

behavior, if we choose: $\left[-\alpha_n + \frac{\partial^2}{\partial y^2} \right] g_n(y, y') = -4\pi\delta(y - y'),$

which in turn can be expressed in terms of the two independent solutions $v_{n_1}(y)$ and $v_{n_2}(y)$ of the homogeneous equation:

$$\left[\frac{d^2}{dy^2} - \alpha_n \right] v_{n_1}(y) = 0, \quad \left[\frac{d^2}{dy^2} - \alpha_n \right] v_{n_2}(y) = 0$$

and the Wronskian constant: $K_n \equiv \frac{dv_{n_1}}{dy} v_{n_2} - v_{n_1} \frac{dv_{n_2}}{dy}$



$$\left[-\alpha_n + \frac{\partial^2}{\partial y^2} \right] g_n(y, y') = -4\pi\delta(y - y'),$$

$$g_n(y, y') = \frac{4\pi}{K_n} v_{n_1}(y_<) v_{n_2}(y_>)$$

where: $\left[\frac{d^2}{dy^2} - \alpha_n \right] v_{n_i}(y) = 0,$


and $K_n \equiv \frac{dv_{n_1}}{dy} v_{n_2} - v_{n_1} \frac{dv_{n_2}}{dy}$

For example, choose $v_{n_1}(y) = \sinh(\sqrt{\alpha_n} y)$ and $v_{n_2}(y) = \sinh(\sqrt{\alpha_n} (b - y))$

where $K_n = \sqrt{\alpha_n} \sinh(\sqrt{\alpha_n} b)$

using the identity: $\cosh(r)\sinh(s) + \sinh(r)\cosh(s) = \sinh(r + s)$

Example:



Two dimensional box with sides a and b with boundary conditions: $\Phi(0,y)=\Phi(a,y)=\Phi(x,0)=\Phi(x,b)=0$

$$\Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V d^3 r' \rho(\mathbf{r}') G(\mathbf{r}, \mathbf{r}') +$$


Don't know this term

$$\frac{1}{4\pi} \int_S d^2 r' [G(\mathbf{r}, \mathbf{r}') \nabla' \Phi(\mathbf{r}') - \Phi(\mathbf{r}') \nabla' G(\mathbf{r}, \mathbf{r}')] \cdot \hat{\mathbf{r}}'.$$

Know this term

$$G(x, x', y, y') = \sum_n u_n(x) u_n(x') \frac{4\pi}{K_n} v_{n_1}(y_-) v_{n_2}(y_+).$$

Example:



Two dimensional box with sides a and b with boundary conditions: $\Phi(0,y)=\Phi(a,y)=\Phi(x,0)=\Phi(x,b)=0$

For this type of problem, it is necessary to construct $G(x, x', y, y')$ so that it vanishes on the boundary:

$$G(x, x', y, 0) = G(x, x', y, b) = G(x, 0, y, y') = G(x, a, y, y') = 0$$

$$G(x, x', y, y') = \sum_n u_n(x) u_n(x') \frac{4\pi}{K_n} v_{n_1}(y_<) v_{n_2}(y_>).$$

$$\frac{d^2}{dx^2} u_n(x) = -\alpha_n^2 u_n(x) \quad \text{where} \quad u_n(0) = u_n(a) = 0$$

$$\Rightarrow u_n(x) = \sqrt{\frac{2}{a}} \sin\left(\frac{n\pi x}{a}\right) \quad \alpha_n = \left(\frac{n\pi}{a}\right)^2$$

$$\left[\frac{d^2}{dy^2} - \left(\frac{n\pi}{a}\right)^2 \right] v_{n_i}(y) = 0$$

$$v_{n_1}(y) = \sinh\left(\frac{n\pi}{a}y\right) \quad v_{n_2}(y) = \sinh\left(\frac{n\pi}{a}(b-y)\right)$$

$$K_n = \frac{n\pi}{a} \sinh\left(\frac{n\pi b}{a}\right)$$





Green's function construction -- continued

$$G(x, x', y, y') = \sum_n u_n(x) u_n(x') K_n v_{n_1}(y_<) v_{n_2}(y_>).$$

For example, a Green's function for a two-dimensional rectangular system with $0 \leq x \leq a$ and $0 \leq y \leq b$, which vanishes on the rectangular boundaries:

$$G(x, x', y, y') = 8 \sum_{n=1}^{\infty} \frac{\sin\left(\frac{n\pi x}{a}\right) \sin\left(\frac{n\pi x'}{a}\right) \sinh\left(\frac{n\pi y_<}{a}\right) \sinh\left(\frac{n\pi}{a}(b - y_>)\right)}{n \sinh\left(\frac{n\pi b}{a}\right)}.$$





$$\Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V d^3 r' \rho(\mathbf{r}') G(\mathbf{r}, \mathbf{r}') + \mathbf{=0}$$

(for our case)

$$\frac{1}{4\pi} \int_S d^2 r' [G(\mathbf{r}, \mathbf{r}') \nabla' \Phi(\mathbf{r}') - \Phi(\mathbf{r}') \nabla' G(\mathbf{r}, \mathbf{r}')] \cdot \hat{\mathbf{r}}'.$$

$$G(x, x', y, y') = 8 \sum_{n=1}^{\infty} \frac{\sin\left(\frac{n\pi x}{a}\right) \sin\left(\frac{n\pi x'}{a}\right) \sinh\left(\frac{n\pi y_-}{a}\right) \sinh\left(\frac{n\pi}{a}(b - y_+)\right)}{n \sinh\left(\frac{n\pi b}{a}\right)}.$$



$$G(x, x', y, y') = 8 \sum_{n=1}^{\infty} \frac{\sin\left(\frac{n\pi x}{a}\right) \sin\left(\frac{n\pi x'}{a}\right) \sinh\left(\frac{n\pi y_<}{a}\right) \sinh\left(\frac{n\pi}{a}(b - y_>)\right)}{n \sinh\left(\frac{n\pi b}{a}\right)}$$

Example: $\rho(x, y) = \rho_0 \sin\left(\frac{\pi x}{a}\right) \sin\left(\frac{\pi y}{b}\right)$

$$\Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V d^3 r' \rho(\mathbf{r}') G(\mathbf{r}, \mathbf{r}')$$

In this example, only $n=1$ contributes because

$$\int_0^a dx' \sin\left(\frac{\pi x'}{a}\right) \sin\left(\frac{n\pi x'}{a}\right) = \frac{a}{2} \delta_{1n}$$

$$\Phi(x, y) = \frac{8\rho_0}{4\pi\epsilon_0} \frac{a}{2 \sinh(\pi b/a)} \sin\left(\frac{\pi x}{a}\right) \times \\ \left(\sinh\left(\frac{\pi(b-y)}{a}\right) \int_0^y dy' \sin\left(\frac{\pi y'}{b}\right) \sinh\left(\frac{\pi y'}{a}\right) + \sinh\left(\frac{\pi y}{a}\right) \int_y^b dy' \sin\left(\frac{\pi y'}{b}\right) \sinh\left(\frac{\pi(b-y')}{a}\right) \right)$$

(somewhat painful solution to example problem)

After asking Maple to perform this integral – (and after the dust clears), we find for

$$\rho(x, y) = \rho_0 \sin\left(\frac{\pi x}{a}\right) \sin\left(\frac{\pi y}{b}\right)$$

$$\Phi(x, y) = \frac{\rho_0}{\epsilon_0} \frac{\sin\left(\frac{\pi x}{a}\right) \sin\left(\frac{\pi y}{b}\right)}{\left(\left(\frac{\pi}{a}\right)^2 + \left(\frac{\pi}{b}\right)^2\right)}$$

Check -- $\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) \Phi(x, y) \stackrel{?}{=} -\frac{\rho_0}{\epsilon_0} \sin\left(\frac{\pi x}{a}\right) \sin\left(\frac{\pi y}{b}\right)$



Another example: $\rho(x, y) = \rho_0$ for $0 \leq x \leq a$ and $0 \leq y \leq b$

$$\Phi(\mathbf{r}) = \frac{1}{4\pi\epsilon_0} \int_V d^3 r' \rho(\mathbf{r}') G(\mathbf{r}, \mathbf{r}')$$

Again -- $G(x, x', y, y') = 8 \sum_{n=1}^{\infty} \frac{\sin\left(\frac{n\pi x}{a}\right) \sin\left(\frac{n\pi x'}{a}\right) \sinh\left(\frac{n\pi y_-}{a}\right) \sinh\left(\frac{n\pi}{a}(b - y_+)\right)}{n \sinh\left(\frac{n\pi b}{a}\right)}$.

$$\int_0^a dx' \sin\left(\frac{n\pi x'}{a}\right) = \begin{cases} 0 & n = \text{even} \\ \frac{2a}{n\pi} & n = \text{odd} \end{cases}$$

$$\Phi(x, y) =$$

$$\frac{16\rho_0 a}{4\pi^2 \epsilon_0} \sum_{n \text{ (odd)}} \frac{\sin\left(\frac{n\pi x}{a}\right)}{n^2 \sinh\left(\frac{n\pi b}{a}\right)} \left(\sinh\left(\frac{n\pi(b-y)}{a}\right) \int_0^y dy' \sinh\left(\frac{n\pi y'}{a}\right) + \sinh\left(\frac{n\pi y}{a}\right) \int_y^b dy' \sinh\left(\frac{n\pi(b-y')}{a}\right) \right)$$

$$= \frac{16\rho_0 a^2}{4\pi^3 \epsilon_0} \sum_{n \text{ (odd)}} \frac{\sin\left(\frac{n\pi x}{a}\right)}{n^3 \sinh\left(\frac{n\pi b}{a}\right)} \left(\sinh\left(\frac{n\pi b}{a}\right) - \sinh\left(\frac{n\pi y}{a}\right) - \sinh\left(\frac{n\pi(b-y)}{a}\right) \right)$$

=

A complicated expression that converges rapidly.

Note that up to now, we have been considering Cartesian coordinates in 1, 2, and 3 dimensions. We will need additional considerations for systems that are more naturally described in curvilinear coordinates.

Cylindrical coordinates: $\Phi(r, \varphi, z)$

$$\nabla^2 \Phi(r, \varphi, z) = \left(\frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2}{\partial \varphi^2} + \frac{\partial^2}{\partial z^2} \right) \Phi(r, \varphi, z)$$

Spherical polar coordinates: $\Phi(r, \theta, \varphi)$

$$\nabla^2 \Phi(r, \theta, \varphi) = \left(\frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + \frac{1}{r^2} \left(\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{\sin^2 \theta} \frac{\partial^2}{\partial \varphi^2} \right) \right) \Phi(r, \theta, \varphi)$$

→ Additional eigenfunction expansions will be useful



A useful theorem for electrostatics

The mean value theorem (Problem 1.10 in Jackson)

The “mean value theorem” value theorem (problem 1.10 of your textbook) states that the value of $\Phi(\mathbf{r})$ at the arbitrary (charge-free) point \mathbf{r} is equal to the average of $\Phi(\mathbf{r}')$ over the surface of any sphere centered on the point \mathbf{r} (see Jackson problem #1.10). One way to prove this theorem is the following. Consider a point $\mathbf{r}' = \mathbf{r} + \mathbf{u}$, where \mathbf{u} will describe a sphere of radius R about the fixed point \mathbf{r} . We can make a Taylor series expansion of the electrostatic potential $\Phi(\mathbf{r}')$ about the fixed point \mathbf{r} :

$$\Phi(\mathbf{r} + \mathbf{u}) = \Phi(\mathbf{r}) + \mathbf{u} \cdot \nabla \Phi(\mathbf{r}) + \frac{1}{2!} (\mathbf{u} \cdot \nabla)^2 \Phi(\mathbf{r}) + \frac{1}{3!} (\mathbf{u} \cdot \nabla)^3 \Phi(\mathbf{r}) + \frac{1}{4!} (\mathbf{u} \cdot \nabla)^4 \Phi(\mathbf{r}) + \dots \quad (1)$$

According to the premise of the theorem, we want to integrate both sides of the equation 1 over a sphere of radius R in the variable \mathbf{u} :

$$\int_{\text{sphere}} dS_u = R^2 \int_0^{2\pi} d\phi_u \int_{-1}^{+1} d \cos(\theta_u). \quad (2)$$



Mean value theorem – continued

We note that

$$R^2 \int_0^{2\pi} d\phi_u \int_{-1}^{+1} d \cos(\theta_u) 1 = 4\pi R^2,$$

$$R^2 \int_0^{2\pi} d\phi_u \int_{-1}^{+1} d \cos(\theta_u) \mathbf{u} \cdot \nabla = 0,$$

$$R^2 \int_0^{2\pi} d\phi_u \int_{-1}^{+1} d \cos(\theta_u) (\mathbf{u} \cdot \nabla)^2 = \frac{4\pi R^4}{3} \nabla^2,$$

$$R^2 \int_0^{2\pi} d\phi_u \int_{-1}^{+1} d \cos(\theta_u) (\mathbf{u} \cdot \nabla)^3 = 0,$$

and

$$R^2 \int_0^{2\pi} d\phi_u \int_{-1}^{+1} d \cos(\theta_u) (\mathbf{u} \cdot \nabla)^4 = \frac{4\pi R^6}{5} \nabla^4.$$

Since $\nabla^2 \Phi(\mathbf{r}) = 0$, the only non-zero term of the average is thus the first term:

$$R^2 \int_0^{2\pi} d\phi_u \int_{-1}^{+1} d \cos(\theta_u) \Phi(\mathbf{r} + \mathbf{u}) = 4\pi R^2 \Phi(\mathbf{r}),$$


or

$$\Phi(\mathbf{r}) = \frac{1}{4\pi R^2} R^2 \int_0^{2\pi} d\phi_u \int_{-1}^{+1} d \cos(\theta_u) \Phi(\mathbf{r} + \mathbf{u}) \equiv \frac{1}{4\pi R^2} \int_{\text{sphere}} dS_u \Phi(\mathbf{r} + \mathbf{u}).$$

Since this result is independent of the radius R , we see that we have the theorem.

Summary: Mean value theorem

$$\Phi(\mathbf{r}) = \frac{1}{4\pi R^2} \int R^2 d\Omega_u \Phi(\mathbf{r} + \mathbf{u})$$



Next week we will take a short digression to discuss two different “numerical” methods for solving the Poisson equation, following your textbook.