

Stability of Solitons for the KdV equation in H^s , $0 \leq s < 1$ *Preliminary Version*

S. Raynor and G. Staffilani

ABSTRACT. We study the long-time stability of soliton solutions to the Korteweg-deVries equation. We consider solutions u to the KdV with initial data in H^s , $0 \leq s < 1$, that are initially close in H^s norm to a soliton. We prove that the possible orbital instability of these ground states is at most polynomial in time. This is an analogue to the H^s orbital instability result of [7], and obtains the same maximal growth rate in t . Our argument is based on the “ I -method” used in [7] and other papers of Colliander, Keel, Staffilani, Takaoka and Tao, which pushes these H^s functions to the H^1 norm.

1. Introduction

We will consider the long-time stability of soliton solutions to the Korteweg-deVries Equation. The KdV equation, which was developed as a model for one-dimensional waves in shallow water, is as follows:

$$(1.1) \quad u_t + u_{xxx} + (u^2)_x = 0.$$

We will consider the initial value problem for the KdV with initial data $u_0 \in H^s$, $0 \leq s < 1$. Local well-posedness (that is, short-time existence, uniqueness and uniform continuity with regard to initial data) for the Cauchy problem is known. (See [1] and [11] for the most recent results.) Moreover, the KdV equation has an infinite sequence of conservation laws which hold for any solution which is sufficiently smooth. The first few are:

$$\begin{aligned} G(u) &= \int_{\mathbb{R}} u(x, t) dx = \int_{\mathbb{R}} u(x, 0) dx, \\ \|u(t)\|_{L^2}^2 &= \int_{\mathbb{R}} |u(x, t)|^2 dx = \int_{\mathbb{R}} |u(x, 0)|^2 dx, \\ H(u) &= \int_{\mathbb{R}} (|\partial_x u(x, t)|^2 - \frac{2}{3} u(x, t)^3) dx = \int_{\mathbb{R}} (|\partial_x u(x, 0)|^2 - \frac{2}{3} u(x, 0)^3) dx. \end{aligned}$$

Using the local well-posedness arguments, these conservation laws, and iteration arguments, global well-posedness can be deduced for $s \geq 0$.¹

It is known that the KdV equation admits traveling wave solutions called solitons which satisfy $Q(x, t) = \psi(x - Ct)$, and ψ therefore is a solution to the following ODE:

$$(1.2) \quad \psi_{xx} - C\psi + \psi^2 = 0.$$

There exists a unique even, positive solution ψ_0 to this equation. This soliton is smooth and rapidly decreasing as $|x| \rightarrow \infty$. In fact,

$$(1.3) \quad \psi_0(x) = \frac{3}{2} C \operatorname{sech}^2\left(\frac{1}{2} C^{\frac{1}{2}} x\right).$$

1991 *Mathematics Subject Classification.* 35Q53, 42B35, 37K10, 37B25.

Key words and phrases. Korteweg-de Vries equation, nonlinear dispersive equations, multilinear estimates, stability. G.S. is supported in part by N.S.F. Grant DMS 0100375 and a grant by the Sloan Foundation.

¹Global well-posedness also holds for $s > -\frac{3}{4}$ [10].

For simplicity we will consider only the case $C = 1$, as the others can be recovered by scaling. We will define $\Sigma = \{\psi_0(x - x_0) | x_0 \in \mathbb{R}\}$ to be the one-parameter space of all solitons moving with speed 1. Note that the KdV flow preserves Σ and that each element of Σ is a solution to (1.2).

It was proven by Benjamin [3] in 1972 that soliton solutions are stable in the following sense: if u is a solution to the KdV which is initially close to a soliton in H^1 norm, then for all time u is close to a soliton. Some corrections and extensions of his result were offered by Bona [4]. More recently, Weinstein [14] has offered a general theory which proves the stability of soliton solutions to generalized KdV equations as well as a class of non-linear Schrödinger equations. In [7], Colliander, Keel, Staffilani, Takaoka, and Tao exploited Weinstein's result to prove that the instability of soliton solutions to the NLS in $H^s, 0 \leq s < 1$ grows at most polynomially in t . They made use of a multiplier operator which they had developed in their proof of global well-posedness for dispersive equations with initial data in $H^s, 0 \leq s < 1$. [10] This multiplier operator allowed them to work with H^1 norms, which they could then control using Weinstein's result.

In this paper, we will again exploit the multiplier operator which they developed, as well as the original proof of H^1 stability of solitons for the KdV. We will prove that in $H^s, 0 \leq s < 1$, soliton solutions to the KdV are at most polynomially unstable. Our main result is:

THEOREM 1.1. *Let $0 \leq s < 1$, Let $\sigma = \text{dist}_{H^s}(u_0, \Sigma) \ll 1$, and let u be the solution to the KdV such that $u(\cdot, 0) = u_0$. Then $\text{dist}_{H^s}(u(t), \Sigma) \leq t^{1-s+\epsilon} \sigma$, for all t such that $t \ll \sigma^{-\frac{1}{1-s+\epsilon}}$.*

To prove this, we will employ the Lyapunov functional introduced by Benjamin [3]:

$$\mathcal{L}(u) = \|u\|_{L^2}^2 + H(u) = \int_{\mathbb{R}} |u_x|^2 + |u|^2 - \frac{2}{3}|u|^3.$$

It can be shown using the Gagliardo-Nirenberg inequality that $\mathcal{L} \geq 0$. Note that if u is a solution to the KdV equation with $u \in H^s$ and $s \geq 1$, then $\mathcal{L}(u)$ is conserved. In fact, we have the equation

$$\partial_t \mathcal{L}(u) = 2 \int_{\mathbb{R}} u_t (-u_{xx} + u - u^2) dx,$$

which vanishes if u is a solution to (1.1) by integration by parts. This calculation also shows that solitons, which are solutions to (1.2), are critical points of the functional \mathcal{L} . In [3] (see also [14]), Benjamin proved that they are minimizers and moreover that, for all $u \in H^1$ such that $\text{dist}_{H^1}(u, \Sigma) \ll 1$,

$$(1.4) \quad \mathcal{L}(u) - \mathcal{L}(Q) \sim \text{dist}_{H^1}(u, \Sigma)^2.$$

This then implies the stability of the solitons because $\mathcal{L}(u)$ is conserved in t .

We will extend this result to $H^s, 0 \leq s < 1$, finding the possible growth in time of the distance between u and the solitons to be at most polynomial. To do so, we will exploit the fact that the quantity $\mathcal{L}(Iu)$ is almost conserved in time, where I is a smoothing operator that maps H^s to H^1 . This technique was used by Colliander, Keel, Staffilani, Takaoka, and Tao in [7] to prove polynomial stability bounds for soliton solutions to the Schrödinger Equation. We will follow the technique developed in that paper in general outline, making the necessary estimates for the KdV equation. We will also follow the structure of that paper, giving progressively more sophisticated arguments that get closer to Theorem 1.1 with each iteration.

Several interesting open questions remain. It is not known whether the power of t which we obtain in the theorem is sharp. Moreover, we have not completed the estimates for the modified KdV equation

$$u_t + u_{xxx} + (u^3)_x = 0$$

and it is not known whether such stability results hold in that case. Finally, a recent paper of Merle and Vega [13] has concluded that in fact KdV solitons are stable in L^2 .² The authors are currently studying whether this result and the I -method exploited in this paper can be extended to prove polynomial stability bounds below L^2 .

²The reader may think that some sort of “interpolation” between the H^1 and L^2 stability should give an even better result than the authors obtain, but unfortunately it is no obvious how to “interpolate.”

The structure of this paper is as follows. In section 2, we define our notation and quote some important estimates that will be used in the following sections. In section 3, we make our first attempt at proving the main theorem, obtaining a weaker form of the estimate. In section 4, we refine the techniques of section 3 but still miss the main theorem by an ϵ power in $\text{dist}_{H^s}(u_0, \Sigma)$. Finally, in section 5 we complete the proof of the theorem.

2. Notation and Set-Up

We will use the notation $A \lesssim B$ to mean that $A \leq cB$ where c is a constant depending on s that may vary from line to line, and similarly for the notation $A \sim B$. We will use $\langle \xi \rangle$ to denote $1 + |\xi|$.

We define the spatial Fourier transform by

$$\hat{f}(\xi) = \int_{\mathbb{R}} e^{-ix\xi} f(x) dx$$

and the spacetime Fourier transform by

$$\tilde{u}(\xi, \tau) = \int_{\mathbb{R}} e^{-i(x\xi + t\tau)} f(x, t) dx dt.$$

We define the $X^{s,b}$ space, as in [1], by the norm

$$\|u\|_{X^{s,b}} = \|\langle \xi \rangle^s \langle \tau - \xi^3 \rangle^b \tilde{u}(\xi, \tau)\|_{L^2_{\xi, \tau}}.$$

We will also use the notation

$$X_I^{s,b} = \{u|_{\mathbb{R} \times I} : u \in X^{s,b}\}$$

with the norm

$$\|u\|_{X_I^{s,b}} = \inf\{\|v\|_{X^{s,b}} : v|_{\mathbb{R} \times I} = u\}.$$

We will use the notation

$$\Lambda_n(m(\xi_1, \dots, \xi_n); f_1, \dots, f_n) = \int_{\xi_1 + \dots + \xi_n = 0} [m(\xi_1, \dots, \xi_n)] \hat{f}_1(\xi_1) \cdot \dots \cdot \hat{f}_n(\xi_n) d\xi_1 \dots d\xi_n,$$

where $[m(\xi_1, \dots, \xi_n)]$ is the symmetrization of the multiplier m in the ξ_i variables. Note that we will not always work with the symmetrized multiplier if it does not matter, but that occasionally symmetrization will be necessary to obtain the appropriate estimates.

For $N \gg 1$ and fixed, we define the operator $I = I_N$ to be a smooth even multiplier operator such that

$$(2.1) \quad \widehat{(I_N u)} = \begin{cases} \hat{u} & |\xi| < N \\ \frac{\xi^{s-1}}{N^{s-1}} \hat{u} & |\xi| > 10N \end{cases}.$$

We generally omit the subscript N unless it is necessary for clarity. We also use the notation N_i for a dyadic block in the frequency space of the function u_i , that is, in the domain of the variable ξ_i . Note that N_i is not necessarily positive. We will write u_{i, N_i} for the function obtained from u_i by restricting it to its components with frequency in N_i . That is, if ϕ_{N_i} is a smooth cutoff function which is the identity in $[N_i, 2N_i]$ and which has support in $[N_i - 1, 2N_i + 1]$, then $\widehat{u_{i, N_i}} = \phi_{N_i} \hat{u}_i$.

Finally, we will denote by $W(t)$ the solution operator for the linear KdV equation, $u_t + u_{xxx} = 0$.

We will also use the following estimates:

(1) By the Plancherel Theorem and Cauchy-Schwartz, we have³ and the proof of Lemma 8.1 in [8]:

$$(2.2) \quad \int \int |u_1 u_2| dx dt = \int \int \left(\frac{|\widehat{u_1}|}{\langle \tau - \xi^3 \rangle^{b+\epsilon}} \right) (\langle \tau - \xi^3 \rangle^{b+\epsilon} |\widehat{u_2}|) d\xi d\tau \leq \|u_1\|_{X^{0, -b-\epsilon}} \|u_2\|_{X^{0, b+\epsilon}}$$

(2) The KdV bilinear estimate [11]:

$$(2.3) \quad \|\partial_x(u_1 u_2)\|_{X^{0,b}} \leq \|u_1\|_{X^{0, b+1}} \|u_2\|_{X^{0, b+1}},$$

for $b = -\frac{1}{2} + \epsilon$, for any $\epsilon > 0$.

³For a more precise proof of this estimate on a finite interval in t , see [9]

(3) The Strichartz Estimate [12]:

$$(2.4) \quad \|D_x^{\frac{\theta\alpha}{2}} u\|_{L_t^q L_x^p} \leq C(\theta, \alpha) \|u\|_{X^{0, \frac{1}{2}+\epsilon}},$$

for all $(\theta, \alpha) \in [0, 1] \times [0, \frac{1}{2}]$, with $p = \frac{2}{1-\theta}$ and $q = \frac{6}{\theta(\alpha+1)}$. Here, as elsewhere, the norm $\|\cdot\|_{L_t^q L_x^p}$ will mean to take the L^p norm with respect to x first and then to take the L^q norm with respect to t .

3. A First Pass at the Theorem

In this section, we obtain a weaker version of the main result of this paper. As mentioned in the introduction, we will follow the structure of [7] because we believe that in this way the argument can be better understood. Even though the structure is the same we have to repeat most of the arguments because the estimates are different. In addition, in following sections we will use the estimates proven here.

PROPOSITION 3.1. *Let $0 \leq s < 1$. Let $\sigma = \text{dist}_{H^s}(u_0, \Sigma) \ll 1$, and let u be the solution to the KdV such that $u(\cdot, 0) = u_0$. Then*

$$\text{dist}_{H^s}(u(t), \Sigma) \leq C t^{\frac{1-s}{3-2s-\epsilon}} \sigma^{\frac{1}{3-2s-\epsilon}}$$

for some small $\epsilon > 0$ and for all t such that $t \ll \sigma^{-\frac{1}{(1-s)-\epsilon}}$.

PROOF Fix s , u_0 , and σ .

Let $N \gg 1$. We will fix N later subject to some future constraints. Let I_N be the multiplier operator discussed in 2.1 with cutoff point N . From now on we will refer to I_N simply as I unless that is unclear.

Define

$$(3.1) \quad E_N(t) = \mathcal{L}(Iu(t)).$$

Let ψ be a ground state such that $\|u_0 - \psi\|_{H^s} = \sigma$. Then $\|Iu_0 - I\psi\| \leq CN^{1-s}\sigma$. Moreover, because ψ is smooth, its Fourier transform is rapidly decreasing, so $\|I\psi - \psi\|_{H^1} \leq CN^{-C_1}$ for any C_1 we choose. So, if we require that $N \geq \sigma^{-\epsilon}$ for some $\epsilon > 0$, then we obtain $\|I\psi - \psi\|_{H^1} \leq CN^{1-s}\sigma$, so

$$\|Iu_0 - \psi\|_{H^1} \leq CN^{1-s}\sigma.$$

By (1.4) this implies (for σ sufficiently small with respect to N) that

$$|E_N(0) - \mathcal{L}(\psi)| \lesssim N^{2-2s}\sigma^2.$$

We will need the following lemma, the proof of which is postponed until later:

LEMMA 3.2. *If there is $t_0 \in \mathbb{R}$ such that $|E_N(t_0) - \mathcal{L}(\psi)| \ll 1$, then*

$$|E_N(t_0 + \delta) - E_N(t_0)| \leq \mathcal{O}\left(\frac{1}{N^{1-\epsilon}}\right)$$

where δ is an absolute constant depending only on s .⁴

For now we will assume the lemma. Once we have this lemma, by the same argument which appears in [7], we can iterate to say that

$$|E_N(t) - \mathcal{L}(\psi)| \lesssim N^{2-2s}\sigma^2,$$

for all t such that $t \ll N^{1-\epsilon}N^{2-2s}\sigma^2$. We may therefore conclude that, for all such t ,

$$\|u(t) - \psi\|_{H^s} \lesssim N^{1-s}\sigma.$$

We finally optimize N under the necessary constraints:

$$(3.2) \quad N^{3-2s-\epsilon}\sigma^2 \gg 1 \quad t \ll N^{1-\epsilon}N^{2-2s}\sigma^2 \quad N^{2-2s}\sigma^2 \ll 1$$

and conclude that

$$\|u - \psi\|_{H^s} \lesssim t^{\frac{1-s}{3-2s-\epsilon}} \sigma^{\frac{1}{3-2s-\epsilon}},$$

for all $t \ll \sigma^{\frac{-1}{1-s-\epsilon}}$. \square

⁴Note that δ may depend also on $\|u(t_0)\|_{H^s}$, but $\|u(t_0) - \psi\|_{H^s} \ll 1$ by [14] and $\|\psi\|_{H^s}$ depends only on s , so $\|u(t_0)\|_{H^s}$ also can be controlled by a constant dependent only on s .

It remains for us to prove the lemma:

PROOF (of Lemma 3.2) To prove Lemma 3.2, we first control $\|Iu\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}}$.⁵ We will then use this control to take a δ -step forward in time and measure the growth of $E_N(t)$ in this time step.

CLAIM 1. *There exists a $\delta > 0$ such that, for $0 < \epsilon \ll 1$,*

$$\|Iu\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} \lesssim 1.$$

PROOF (of Claim) First note that by (1.4), $\|Iu(t_0)\|_{H^1} \lesssim 1$, because $|E_N(t_0) - \mathcal{L}(\psi)| \ll 1$ and $\|\psi\|_{H^1}$ is a constant. Moreover, I commutes with differentiation and with $W(t)$. We may therefore apply the standard $X^{s,b}$ estimates (see, e.g., [11], pp. 587-8). Let $\phi(t)$ be a cutoff function with support in $[t_0 - 3, t_0 + 3]$, such that $\phi \equiv 1$ inside $[t_0 - 2, t_0 + 2]$. Then, u is a fixed point of the operator

$$Lu = \phi(t)W(t - t_0)u(t_0) + \phi(t) \int_{t_0}^t W(t - t') \partial_x(u(t')^2) dt'$$

on the interval $[t_0 - 2, t_0 + 2]$. Then, for $0 < \delta \ll 1$:

$$\begin{aligned} \|Iu\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} &\leq \|I\psi(t)W(t - t_0)u(t_0)\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} + \|I\psi(t) \int_{t_0}^t W(t - t') \partial_x(u(t')^2) dt'\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} \\ &\leq C\|Iu(t_0)\|_{H^1} + C\|\partial_x Iu^2\|_{X_{[t_0-\delta, t_0+\delta]}^{1, -\frac{1}{2}+\epsilon}} \\ &\leq C\|Iu(t_0)\|_{H^1} + C\delta^\epsilon \|\partial_x Iu^2\|_{X_{[t_0-\delta, t_0+\delta]}^{1, -\frac{1}{2}+2\epsilon}}. \end{aligned}$$

Now, by the bilinear estimate for the KdV (see [11]), we have, for $s \geq -\frac{3}{4}$:

$$\|\partial_x u^2\|_{X_{[t_0-\delta, t_0+\delta]}^{s, -\frac{1}{2}+2\epsilon}} \leq \|u\|_{X_{[t_0-\delta, t_0+\delta]}^{s, \frac{1}{2}+\epsilon}}^2.$$

Consider the multiplier operator I_1 , which is the same type of operator as I but with $N = 1$. It is clear that $\|f\|_{X^{s,b}} \sim \|I_1 f\|_{X^{1,b}}$, so

$$\|\partial_x I_1 u^2\|_{X_{[t_0-\delta, t_0+\delta]}^{1, -\frac{1}{2}+2\epsilon}} \sim \|\partial_x u^2\|_{X_{[t_0-\delta, t_0+\delta]}^{s, -\frac{1}{2}+2\epsilon}} \lesssim \|u\|_{X_{[t_0-\delta, t_0+\delta]}^{s, \frac{1}{2}+\epsilon}}^2 \sim \|I_1 u\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}}^2.$$

But then, by Lemma 12.1 of [6] it also follows for general N that

$$\|\partial_x I_N u^2\|_{X_{[t_0-\delta, t_0+\delta]}^{1, -\frac{1}{2}+2\epsilon}} \lesssim \|I_N u\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}}^2.$$

We may therefore conclude that

$$\|Iu\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} \leq C\|Iu_0\|_{H^1} + C\delta^\epsilon \|Iu\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}}^2 \leq C + C\delta^\epsilon \|Iu\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}}^2.$$

Therefore, by a continuity argument, there exists a $\delta > 0$ for which

$$\|Iu\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} \leq 2C \lesssim \|Iu(t_0)\|_{H^1} \lesssim 1.$$

This concludes the proof of the claim. \square

We now want to take a step forward in time. Let $f \in H^1$. Define $\Omega(f(t)) = \partial_t(\mathcal{L}(f(t)))$. Then:

$$\begin{aligned} \Omega(f(t)) &= \partial_t(\mathcal{L}(f)) = \partial_t \left(\int_{\mathbb{R}} (f_x^2 + f^2 - \frac{2}{3}f^3) dx \right) \\ &= 2 \int_{\mathbb{R}} (f_x f_{xt} + f f_t - f^2 f_t) dx \\ (3.3) \quad &= 2 \int_{\mathbb{R}} f_t (-f_{xx} + f - f^2) dx. \end{aligned}$$

⁵Due to the special features of the KdV equation, the $X^{s,b}$ norms have been found to be effective to work with.

In our case, we are interested in

$$\begin{aligned}
E_N(t_0 + \delta) - E_N(t_0) &= \int_{t_0}^{t_0 + \delta} \Omega(Iu(t)) \, dt \\
&= 2 \int_{t_0}^{t_0 + \delta} \int_{\mathbb{R}} Iu_t (-Iu_{xx} + Iu - (Iu)^2) \, dx \, dt \\
&= -2 \int_{t_0}^{t_0 + \delta} \int_{\mathbb{R}} (Iu_{xxx} + I(u^2)_x) (-Iu_{xx} + Iu - (Iu)^2) \, dx \, dt \\
&= -2 \int_{t_0}^{t_0 + \delta} \int_{\mathbb{R}} Iu_{xxx} ((Iu)^2 - Iu^2) \, dx \, dt + 2 \int_{t_0}^{t_0 + \delta} \int_{\mathbb{R}} (u^2)_x I^2 u \, dx \, dt + \\
&\quad - 2 \int_{t_0}^{t_0 + \delta} \int_{\mathbb{R}} I(u^2)_x (Iu)^2 \, dx \, dt \\
&= -2 \int_{t_0}^{t_0 + \delta} \Lambda_3 (\xi_1^3 m(\xi_1) (m(\xi_2) m(\xi_3) - m(\xi_2 + \xi_3)); u; u; u) \, dt + \\
&\quad + 4 \int_{t_0}^{t_0 + \delta} \Lambda_3 (\xi_1 m(\xi_3)^2); u; u; u) \, dt + \\
&\quad - 4 \int_{t_0}^{t_0 + \delta} \Lambda_4 (\xi_1 m(\xi_1 + \xi_2) m(\xi_3) m(\xi_4); u; u; u; u) \, dt.
\end{aligned}$$

We will prove the following more general estimates in order to control $E_N(t_0 + \delta) - E_N(t_0)$:

$$(3.4) \quad \left| \int_{t_0}^{t_0 + \delta} \Lambda_3 (\xi_1^3 m(\xi_1) (m(\xi_2) m(\xi_3) - m(\xi_2 + \xi_3)); u_1; u_2; u_3) \, dt \right| \lesssim N^{-1+\epsilon} \prod_{i=1}^3 \|Iu_i\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}},$$

$$(3.5) \quad \left| \int_{t_0}^{t_0 + \delta} \Lambda_3 (\xi_1 m(\xi_3)^2); u_1; u_2; u_3) \, dt \right| \lesssim N^{-1+\epsilon} \prod_{i=1}^3 \|Iu_i\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}},$$

$$(3.6) \quad \left| \int_{t_0}^{t_0 + \delta} \Lambda_4 ((\xi_1) m(\xi_1 + \xi_2) m(\xi_3) m(\xi_4); u_1; u_2; u_3; u_4) \, dt \right| \lesssim N^{-1+\epsilon} \prod_{i=1}^4 \|Iu_i\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}}.$$

Recall that $m(\xi)$ is the multiplier associated with the operator I , and it is identically 1 for $|\xi| \leq N$, and equals $\frac{\xi^{s-1}}{N^{s-1}}$ for $|\xi| > 10N$. Note that because our norms are of L^2 type, we may replace \hat{u} by $|\hat{u}|$ without affecting the estimates. For each estimate, we will divide the functions u_i into dyadic blocks N_i in frequency space and make appropriate estimates. We will then sum over these dyadic blocks to obtain the full estimate.

PROOF (of Estimate (3.4)) We consider the multiplier $N_1^3 m(N_1) (m(\xi_2) m(\xi_3) - m(\xi_2 + \xi_3))$. Recall that we have $N_1 + N_2 + N_3 = 0$ and note that we may assume $N_2 \geq N_3$ because of the symmetry, and that $N_2 > N$ or else the whole symbol is 0. We will consider two cases:

(1) $N_2 \gg N_3$: This implies that $N_1 \sim N_2$.

First suppose that $N_3 \leq N$. Then $m(N_3) = 1$, so

$$N_1^3 m(N_1) (m(N_2) m(N_3) - m(N_2 + N_3)) = N_1^3 m(N_1) (m(N_2) - m(N_2 + N_3)).$$

By the mean value theorem, this is $\leq N_1^3 m(N_1) m'(N_2) N_3$, so, since $N_1 \sim N_2$ and $m'(N_2) = \frac{m(N_2)}{N_2}$,

$$(3.7) \quad N_1^3 m(N_1) (m(N_2) m(N_3) - m(N_2 + N_3)) \leq N_1^3 m(N_1) m'(N_2) N_3 m(N_1) m(N_2) m(N_3).$$

Now, consider the whole integral, and use inequality (2.2):⁶

$$\begin{aligned} & N_1 N_2 N_3 m(N_1) m(N_2) m(N_3) \int_{t_0}^{t_0+\delta} \int |\hat{u}_{1,N_1} \hat{u}_{2,N_2} \hat{u}_{3,N_3}| d\xi dt \\ & \leq N_1 N_2 N_3 m(N_1) m(N_2) m(N_3) \|u_{1,N_1} u_{3,N_3}\|_{X^{0,-\frac{1}{2}+\epsilon}} \|u_{2,N_2}\|_{X^{0,\frac{1}{2}+\epsilon}}. \end{aligned}$$

Then, by the KdV bilinear estimate and because $\|\partial_x u_{1,N_1}\| \sim N_1 \|u_{1,N_1}\|$ and $N_3 \ll N_1$, we obtain:

$$\begin{aligned} & N_1 N_2 N_3 m(N_1) m(N_2) m(N_3) \int_{t_0}^{t_0+\delta} \int |\hat{u}_{1,N_1} \hat{u}_{2,N_2} \hat{u}_{3,N_3}| d\xi dt \\ & \leq N_1 N_2 N_3 m(N_1) m(N_2) m(N_3) \frac{1}{N_1} \|\partial_x(u_{1,N_1} u_{3,N_3})\|_{X^{0,-\frac{1}{2}+\epsilon}} \|u_2\|_{X^{0,\frac{1}{2}+\epsilon}} \\ & \leq N_2 N_3 m(N_1) m(N_2) m(N_3) \|u_{1,N_1}\|_{X^{0,\frac{1}{2}+\epsilon}} \|u_{2,N_2}\|_{X^{0,\frac{1}{2}+\epsilon}} \|u_{3,N_3}\|_{X^{0,\frac{1}{2}+\epsilon}}. \end{aligned}$$

But then, by definition of I and the $X^{s,b}$ spaces and because $N_1 \sim N_2$, this is controlled by

$$\frac{1}{N_1^{\frac{1}{2}-\epsilon}} \frac{1}{N_2^{\frac{1}{2}}} \frac{1}{N_3^{\epsilon}} \|Iu_{1,N_1}\|_{X^{1,\frac{1}{2}+\epsilon}} \|Iu_{2,N_2}\|_{X^{1,\frac{1}{2}+\epsilon}} \|Iu_{3,N_3}\|_{X^{1,\frac{1}{2}+\epsilon}}.$$

When we sum this in the N_i s, we will lose a power of ϵ , and obtain a term of size $\frac{1}{N^{1-\epsilon}}$ as claimed.

Now, suppose instead that $N_2 \gg N_3 > N$. Then

$$\begin{aligned} & N_1^3 m(N_1) (m(N_2) m(N_3) - m(N_2 + N_3)) = \\ & N_1^3 m(N_1) (m(N_2) m(N_3) - m(N_3) m(N_2 + N_3)) + N_1^3 m(N_1) (m(N_3) m(N_2 + N_3) - m(N_2 + N_3)) \\ & = M_1 + M_2 \end{aligned}$$

For estimate M_1 , use the mean value theorem (recall that $N_1 \sim N_2$):

$$M_1 \leq N_1 N_2 N_3 m(N_1) m(N_2) m(N_3).$$

Then the same calculation as before implies that the part of the left-hand side of (3.4) containing M_1 also sums to $\frac{1}{N^{1-\epsilon}}$ as desired.

On the other hand,

$$M_2 = N_1^3 m(N_1) m(N_2 + N_3) (m(N_3) - 1).$$

Note that $|m(N_3) - 1| \leq 2$, and $m(N_2 + N_3) \sim m(N_2)$ because $N_2 \gg N_3$. So,

$$M_2 \leq N_1^3 m(N_1) m(N_2) \frac{m(N_3)}{m(N_3)} \lesssim \frac{N_1^2 N_2 N_3 m(N_1) m(N_2) m(N_3)}{N_3 m(N_3)},$$

where $m(N_3) \sim \frac{N_3^{s-1}}{N_3^{2-1}}$, so $\frac{1}{N_3 m(N_3)} \sim \frac{1}{N_3^s N_3^{1-s}}$. Therefore, we find that

$$M_2 \lesssim \frac{1}{N^{1-s}} \frac{1}{N_3^s} N_1^2 N_2 N_3 m(N_1) m(N_2) m(N_3) N_1.$$

As before, we compute that

$$\begin{aligned} & \int_{t_0}^{t_0+\delta} \int |\hat{u}_{1,N_1} \hat{u}_{2,N_2} \hat{u}_{3,N_3}| d\xi dt \leq \|u_1 u_3\|_{X^{0,-\frac{1}{2}+\epsilon}} \|u_2\|_{X^{0,\frac{1}{2}+\epsilon}} \\ & \leq \frac{1}{N_1} \|u_{1,N_1}\|_{X^{0,\frac{1}{2}+\epsilon}} \|u_{2,N_2}\|_{X^{0,\frac{1}{2}+\epsilon}} \|u_{3,N_3}\|_{X^{0,\frac{1}{2}+\epsilon}}. \end{aligned}$$

And so, the part of the left-hand side of (3.4) containing M_2 is bounded by

$$\frac{1}{N^{1-s}} \frac{1}{N_3^s} \|Iu_{1,N_1}\|_{X^{1,\frac{1}{2}+\epsilon}} \|Iu_{2,N_2}\|_{X^{1,\frac{1}{2}+\epsilon}} \|Iu_{3,N_3}\|_{X^{1,\frac{1}{2}+\epsilon}}.$$

⁶Here we are ignoring that we are on a finite time interval. To be precise one should repeat the argument given in [8] during the proof of Lemma 8.1

To sum this, we use Cauchy-Schwartz and the fact that $N_1 \sim N_2$, to obtain the same estimate as before.

- (2) Now consider the case where $N_2 \sim N_3$. Then $N_1 = -(N_2 + N_3)$ may be smaller. We once again want to estimate the multiplier

$$N_1^3 m(N_1)(m(N_2)m(N_3) - m(N_2 + N_3)) = N_1^3 m(N_1)m(N_2)m(N_3) - N_1^3 m(N_1)^2 = M_3 + M_4.$$

We have

$$\begin{aligned} M_3 &= N_1^3 m(N_1)m(N_2)m(N_3) = N_1 N_2 N_3 m(N_1)m(N_2)m(N_3) \frac{N_1^2}{N_2 N_3} \\ &\lesssim N_1 N_2 N_3 m(N_1)m(N_2)m(N_3). \end{aligned}$$

Then, by the same argument as for the first part of the first case, this sums to $\mathcal{O}(\frac{1}{N^{1-\epsilon}})$. For M_4 , we have

$$\begin{aligned} M_4 &\lesssim N_1 N_2 N_3 m(N_1)m(N_2)m(N_3) \frac{N_1^2}{N_2 N_3} \frac{m(N_1)}{m(N_2)m(N_3)} \\ &= N_1 N_2 N_3 m(N_1)m(N_2)m(N_3) \frac{N_1^{1-s}}{N^{1-s} N_2^s N_3^s}. \end{aligned}$$

We then use the bilinear estimate as before to conclude that

$$\begin{aligned} N_1^3 m(N_1)^2 \int_{t_0}^{t_0+\delta} \int |\hat{u}_{1,N_1} \hat{u}_{2,N_2} \hat{u}_{3,N_3}| d\xi dt &\lesssim N_1 N_2 N_3 m(N_1)m(N_2)m(N_3) \frac{N_1^s}{N^{1-s} N_2^s N_3^s} \|u_{1,N_1}\|_{X^{0,\frac{1}{2}+\epsilon}} \|u_{2,N_2}\|_{X^{0,\frac{1}{2}+\epsilon}} \|u_{3,N_3}\|_{X^{0,\frac{1}{2}+\epsilon}} \\ &\lesssim \frac{1}{N^{1-s}} \frac{1}{N_1^{2\epsilon}} \frac{1}{N_2^{\frac{s}{2}-\epsilon}} \frac{1}{N_3^{\frac{s}{2}-\epsilon}} \|Iu_{1,N_1}\|_{X^{1,\frac{1}{2}+\epsilon}} \|Iu_{2,N_2}\|_{X^{1,\frac{1}{2}+\epsilon}} \|Iu_{3,N_3}\|_{X^{1,\frac{1}{2}+\epsilon}}, \end{aligned}$$

after using again the fact that $N_1 \leq N_2 \sim N_3$. Summing in the N_i s, we can see that this again gives $\mathcal{O}(\frac{1}{N^{1-\epsilon}})$.

This concludes the proof of estimate (3.4). \square

We next need to prove the estimate (3.5):

$$\left| \int_{t_0}^{t_0+\delta} \Lambda_3 (\xi_1 m(\xi_3)^2); u; u; u \right| dt \lesssim N^{-1+\epsilon} \prod_{i=1}^3 \|Iu_i\|_{X^{1,\frac{1}{2}+\epsilon}}$$

PROOF (of Estimate (3.5))

We will consider the multiplier $N_1 m(N_3)^2$. Note that if N_1 , N_2 , and N_3 are all less than N , then the operator given by the symmetrization of this multiplier is identically zero. So at least one of N_1 , N_2 , and N_3 must be greater than N . If $N_3 < N$, this multiplier is just N_1 , and, as above,

$$\begin{aligned} N_1 \int_{t_0}^{t_0+\delta} \int |\hat{u}_{1,N_1} \hat{u}_{2,N_2} \hat{u}_{3,N_3}| d\xi dt &\lesssim N_1 \|u_{1,N_1} u_{3,N_3}\|_{X^{0,-\frac{1}{2}+\epsilon}} \|u_{2,N_2}\|_{X^{0,\frac{1}{2}+\epsilon}} \\ &\lesssim \|u_{1,N_1}\|_{X^{0,\frac{1}{2}+\epsilon}} \|u_{2,N_2}\|_{X^{0,\frac{1}{2}+\epsilon}} \|u_{3,N_3}\|_{X^{0,\frac{1}{2}+\epsilon}} \\ &\lesssim \frac{1}{N_1 m(N_1) N_2 m(N_2) N_3 m(N_3)} \prod_{i=1}^3 \|Iu_{i,N_i}\|_{X^{1,\frac{1}{2}+\epsilon}}. \end{aligned}$$

Since at least one of N_1 , N_2 is greater than N , the quantity computed above sums to no more than $\mathcal{O}(\frac{1}{N^{1-\epsilon}})$.

Now, if $N_3 > N$, as above

$$N_1 m(N_3)^3 \int_{t_0}^{t_0+\delta} \int |\hat{u}_{1,N_1} \hat{u}_{2,N_2} \hat{u}_{3,N_3}| d\xi dt \lesssim \frac{m(N_3)}{N_1 m(N_1) N_2 m(N_2) N_3} \prod_{i=1}^3 \|Iu_{i,N_i}\|_{X^{1,\frac{1}{2}+\epsilon}},$$

which again sums to $\mathcal{O}(\frac{1}{N^{1-\epsilon}})$ in the worst cases. \square

Finally, we need to prove estimate (3.6):

$$\left| \int_{t_0}^{t_0+\delta} \Lambda_4((\xi_1)m(\xi_1 + \xi_2)m(\xi_3)m(\xi_4); u; u; u; u) dt \right| \lesssim N^{-1+\epsilon} \prod_{i=1}^4 \|Iu_i\|_{X^{1, \frac{1}{2}+\epsilon}}$$

PROOF (of Estimate (3.6)) We consider the multiplier $N_1 m(N_1 + N_2) m(N_3) m(N_4)$. Recall that we have $N_1 + N_2 + N_3 + N_4 = 0$ and by symmetry we may assume $N_3 \geq N_4$. Consider

$$\int |\hat{u}_{1,N_1} \hat{u}_{2,N_2} \hat{u}_{3,N_3} \hat{u}_{4,N_4}| d\xi \lesssim \prod_{i=1}^4 \|u_{i,N_i}\|_{L^4}.$$

We use the Strichartz estimate (2.4) with $(\theta, \alpha) = (\frac{1}{2}, 0)$ and $p = 4$, $q = 12$, obtaining

$$\|u\|_{L_t^{12} L_x^4} \leq C \|u\|_{X^{0, \frac{1}{2}+\epsilon}}.$$

In our case, therefore, we may conclude that

$$\begin{aligned} \int_{t_0}^{t_0+\delta} \int |\hat{u}_{1,N_1} \hat{u}_{2,N_2} \hat{u}_{3,N_3} \hat{u}_{4,N_4}| d\xi dt &\leq \int_{t_0}^{t_0+\delta} \prod_{i=1}^4 \|u_{i,N_i}\|_{L^4} dt \\ &\leq \|1\|_{L_t^{\frac{3}{2}}} \prod_{i=1}^4 \|u_{i,N_i}\|_{L_t^{12} L_x^4} \leq C \delta^{\frac{2}{3}} \prod_{i=1}^4 \|u_{i,N_i}\|_{X^{0, \frac{1}{2}+\epsilon}}. \end{aligned}$$

Therefore

$$\begin{aligned} N_1 m(N_1 + N_2) m(N_3) m(N_4) \int |\hat{u}_{1,N_1} \hat{u}_{2,N_2} \hat{u}_{3,N_3} \hat{u}_{4,N_4}| d\xi \\ \leq N_1 N_2 N_3 N_4 m(N_1) m(N_2) m(N_3) m(N_4) \frac{m(N_1 + N_2)}{N_2 N_3 N_4 m(N_1) m(N_2)} \prod_{i=1}^4 \|u_{i,N_i}\|_{X^{0, \frac{1}{2}+\epsilon}} \\ \leq \frac{m(N_1 + N_2)}{m(N_1) m(N_2)} \frac{1}{N_2 N_3 N_4} \prod_{i=1}^4 \|Iu_{i,N_i}\|_{X^{1, \frac{1}{2}+\epsilon}}. \end{aligned}$$

We will now estimate $\frac{m(N_1 + N_2)}{m(N_1) m(N_2)} \frac{1}{N_2 N_3 N_4}$, considering several cases (recall that $N_3 \geq N_4$):

(1) First assume $N_1 \gg N_2$.

If $N_1 \leq N$, $m(N_1 + N_2) = m(N_1) = m(N_2) = 1$. Note that if N_1, N_2, N_3, N_4 are all less than N , then the operator is identically zero by symmetrization. Hence at least one of the dyadic blocks must be at least N for the operator to be nontrivial. Therefore, if $N_1 \leq N$, then $N_3 > N$. Hence the multiplier, which reduces to $\frac{1}{N_2 N_3 N_4}$ in this case clearly sums to no more than $\mathcal{O}(\frac{1}{N^{1-\epsilon}})$.

So we may assume that $N_1 > N$. Then $m(N_1 + N_2) \sim m(N_1)$ because $N_1 \gg N_2$. So our multiplier reduces to $\frac{1}{m(N_2)} \frac{1}{N_2 N_3 N_4}$. If $N_2 < N$, this is again $\frac{1}{N_2 N_3 N_4}$. But now, because $N_1 + N_2 + N_3 + N_4 = 0$, $N_1 \sim N_3$. Hence we may write

$$\frac{1}{N_2 N_3 N_4} \leq \frac{1}{N_1^{\frac{1}{2}} N_2 N_3^{\frac{1}{2}} N_4},$$

which sums to $\mathcal{O}(\frac{1}{N^{2-\epsilon}})$.

Finally, if $N_2 > N$ as well, we have $m(N_2) \sim \frac{N_2^{s-1}}{N^{s-1}}$. Therefore, because $N_1 \sim N_3$, the multiplier is controlled by

$$\frac{1}{N^{1-s} N_2^s N_1^{\frac{1}{2}} N_3^{\frac{1}{2}} N_4},$$

which sums to $\mathcal{O}(\frac{1}{N^{3-\epsilon}})$.

(2) $N_1 \ll N_2$.

Then $m(N_1 + N_2) \sim m(N_2)$, so the multiplier is

$$\frac{1}{m(N_1)N_2N_3N_4}.$$

The case where N_1 and N_2 are both less than N is the same as before. So we consider first what happens when $N_1 < N$. Then we again have $\frac{1}{N_2N_3N_4}$. As before, the operator is trivial unless $N_3 > N$, and when $N_3 > N$ this sums to $\mathcal{O}(\frac{1}{N^{1-\epsilon}})$ as desired.

If instead $N_1 > N$, we have

$$\frac{N_1^{1-s}}{N^{1-s}N_2N_3N_4} \leq \frac{1}{N^{1-s}N_1^\epsilon N_2^{\frac{1+s-\epsilon}{2}} N_3^{\frac{1+s-\epsilon}{2}} N_4},$$

which sums to $\mathcal{O}(\frac{1}{N^{3-\epsilon}})$ as in the first case.

(3) Finally we consider the case where $N_1 \sim N_2$.

Once again the case where both N_1 and N_2 are less than N is the same as before. Therefore, we consider the case where $N_1 \sim N_2 > N$. Then $m(N_1 + N_2) \leq 1$, so the multiplier reduces to $\frac{1}{m(N_1)m(N_2)N_2N_3N_4}$. If $N_3 \sim N_1 \sim N_2$, then this is controlled by $\frac{1}{N^{2(1-s)}N_2^sN_3^sN_4}$ and since N_3 controls all the other quantities, we may again sum to conclude that this is bounded by $\mathcal{O}(\frac{1}{N^{2-\epsilon}})$.

We must at last consider the case $N_3 \ll N_1$. For this case we must reconsider the original calculations done at the beginning of this estimate. Instead of treating all four functions equally, we will write:

$$\int |\hat{u}_{1,N_1} \hat{u}_{2,N_2} \hat{u}_{3,N_3} \hat{u}_{4,N_4}| d\xi \leq \|u_{1,N_1} u_{3,N_3}\|_{L^2} \|u_{2,N_2}\|_{L^4} \|u_{4,N_4}\|_{L^4}.$$

Therefore, using the Strichartz estimate again, the fact that $\|f\|_{X^{0,0}} \leq \|f\|_{X^{0,\frac{1}{2}+\epsilon}}$ for any function f , and the KdV bilinear estimate:

$$\begin{aligned} \int_{t_0}^{t_0+\delta} \int |\hat{u}_{1,N_1} \hat{u}_{2,N_2} \hat{u}_{3,N_3} \hat{u}_{4,N_4}| d\xi dt &\leq \int_{t_0}^{t_0+\delta} \|u_{1,N_1} u_{3,N_3}\|_{L^2} \|u_{2,N_2}\|_{L^4} \|u_{4,N_4}\|_{L^4} dt \\ &\leq \|1\|_{L_t^3} \|u_{1,N_1} u_{3,N_3}\|_{L_t^2 L_x^2} \|u_{2,N_2}\|_{L_t^{12} L_x^4} \|u_{4,N_4}\|_{L_t^{12} L_x^4} \\ &\leq C\delta^{\frac{1}{3}} \|u_{1,N_1} u_{3,N_3}\|_{X^{0,\frac{1}{2}+\epsilon}} \|u_{2,N_2}\|_{X^{0,\frac{1}{2}+\epsilon}} \|u_{4,N_4}\|_{X^{0,\frac{1}{2}+\epsilon}} \\ &\leq C\delta^{\frac{1}{3}} \frac{1}{N_1 + N_3} \prod_{i=1}^4 \|u_{i,N_i}\|_{X^{0,\frac{1}{2}+\epsilon}}. \end{aligned}$$

Now, recall that $N_3 \ll N_1$, $N_3 \geq N_4$, and $N_1 + N_2 + N_3 + N_4 = 0$, so $N_3 + N_1 \sim N_2 \sim N_1$. Therefore, our entire operator may be estimated as follows:

$$\begin{aligned} N_1 m(N_1 + N_2) m(N_3) m(N_4) \int |\hat{u}_{1,N_1} \hat{u}_{2,N_2} \hat{u}_{3,N_3} \hat{u}_{4,N_4}| d\xi \\ \leq m(N_1 + N_2) m(N_3) m(N_4) \prod_{i=1}^4 \|u_{i,N_i}\|_{X^{0,\frac{1}{2}+\epsilon}} \\ \leq \frac{1}{N_1 N_2 N_3 N_4} \frac{m(N_1 + N_2)}{m(N_1) m(N_2)} \prod_{i=1}^4 \|I u_{i,N_i}\|_{X^{1,\frac{1}{2}+\epsilon}}. \end{aligned}$$

We therefore need only to sum

$$\frac{1}{N_1 N_2 N_3 N_4} \frac{m(N_1 + N_2)}{m(N_1) m(N_2)} \lesssim \frac{1}{N^{2(1-s)} N_1^s N_2^s N_3 N_4}$$

which as before is at worst $\mathcal{O}(\frac{1}{N^{2-\epsilon}})$.

This concludes the proof of estimate (3.6). \square

Having proved all three estimates, we note that

$$|E_N(t_0 + \delta) - E_N(t_0)| \lesssim 2((3.4) - (3.5) + (3.6)) \leq \mathcal{O}\left(\frac{1}{N^{1-\epsilon}}\right)$$

since we have already checked that $\|Iu\|_{X_{t_0-\delta, t_0+\delta}^{1, \frac{1}{2}+\epsilon}} \lesssim 1$. This concludes the proof of Lemma 3.2. \square

4. A Second Pass at the Theorem

In this section, we will improve the powers of t and of σ which appear in Proposition 3.1. We will do this by exploiting more carefully the fact that $\|u_0 - \psi\|_{H^s}$ is small.

PROPOSITION 4.1. *Let $0 \leq s < 1$ and suppose $\text{dist}_{H^s}(u_0, \Sigma) = \sigma \ll 1$. Then we have, for some small $\epsilon > 0$,*

$$\text{dist}_{H^s}(u(t), \Sigma) \leq t^{1-s+\epsilon} \sigma^{1+\epsilon}$$

for all t such that $1 < t \ll \sigma^{-\frac{1}{1-s}-\epsilon}$.

PROOF Fix s, u_0 , and σ . We retain the definition of $E_N(t)$ (see 3.1), and the set-up of the previous proposition. The main difference will be a sharper estimate for $E_N(t_0 + \delta) - E_N(t_0)$:

LEMMA 4.2. *If there is a $t_0 \in \mathbb{R}$ and $\tilde{\sigma}$ with $N^{-C} < \tilde{\sigma} \ll 1$ for some arbitrary constant C , such that for some solution to (1.2) ψ , $|E_N(t_0) - \mathcal{L}(\psi)| \leq \tilde{\sigma}^2$ then we have, for some $\delta > 0$ depending only on s ,*

$$E_N(t_0 + \delta) = E_N(t_0) + \mathcal{O}\left(\frac{1}{N^{1-\epsilon}} \tilde{\sigma}^2\right).$$

We will, as in the previous section, postpone the proof of the lemma until later. First we will complete the proof of Proposition 4.1 taking advantage of Lemma 4.2. We can again iterate the lemma. Let $\tilde{\sigma} = N^{1-s}\sigma$. We then obtain

$$|E_N(t) - \mathcal{L}(\psi)| \lesssim N^{2-2s}\sigma^2,$$

for $1 \leq t \ll N^{1-\epsilon}$, and by 1.4 we can then conclude that for all such times t , $\text{dist}_{H^s}(u(t), \Sigma) \lesssim N^{1-s}\sigma$. But now, we may optimize N under the conditions

$$(4.1) \quad N^{-C} > \sigma \quad t \ll N^{1-\epsilon} \quad N^{2-2s}\sigma^2 \ll 1.$$

Contrast these conditions with (3.2). With this improvement, we obtain

$$\text{dist}_{H^s}(u(t), \Sigma) \lesssim t^{1-s+\epsilon} \sigma^{1+\epsilon},$$

for $1 \leq t \ll \sigma^{-\frac{1}{1-s+\epsilon}}$ as claimed. \square

It therefore remains only to prove the lemma:

PROOF (of Lemma 4.2)

By 1.4 and the calculations at the start of Lemma 3.2, there exists a $\psi \in \Sigma$ such that $\|Iu(t_0) - \psi\|_{H^1} \lesssim \tilde{\sigma}$. Let $Q(x, t) = \psi(x - t)$. Define

$$w(x, t) = u(x, t) - Q(x, t).$$

As before ψ is Schwartz and since $N^{-C} \lesssim \tilde{\sigma}$ for some C , we may conclude that $\|Iu(t_0) - I\psi\|_{H^1} \lesssim \tilde{\sigma}$, i.e. $\|w(t_0)\|_{H^1} \lesssim \tilde{\sigma}$.

$$\text{CLAIM 2. } \|Iw\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} \lesssim \tilde{\sigma}.$$

PROOF The function $w(t)$ obeys the following difference equation:

$$(4.2) \quad w_t + w_{xxx} + \partial_x(w(w + 2Q)) = 0.$$

We can therefore use the standard $X^{s,b}$ estimates as in Lemma 3.2 to conclude that

$$\|Iw\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} \leq \|Iw(t_0)\|_{H^1} + \delta^\epsilon \|I(\partial_x(w(w + 2Q)))\|_{X_{[t_0-\delta, t_0+\delta]}^{1, -\frac{1}{2}+\epsilon}}.$$

We then use the bilinear estimate as in Lemma 3.2, as well as the fact that Q is a Schwartz function in x , to conclude that

$$\|Iw\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} \leq \tilde{\sigma} + C\delta^\epsilon \|Iw\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} + \delta^\epsilon \|Iw\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}}^2$$

and therefore, by a continuity argument again, $\|Iw\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} \lesssim \tilde{\sigma}$ for some $\delta > 0$ sufficiently small.

This concludes the proof of the claim. \square

Finally, we must again take a δ -step forward in t . We will show that

$$E_N(t_0 + \delta) - E_N(t_0) = 2 \int_{t_0}^{t_0+\delta} \Omega(I(Q+w)(t)) dt = \mathcal{O}\left(\frac{1}{N^{1-\epsilon}} \tilde{\sigma}^2\right).$$

We will use Lemma 3.2 to do this, following the method of [7], rather than checking it directly.

Because $\tilde{\sigma} \gtrsim N^{-C}$ it will suffice to prove the more general bound

$$E_N(t_0 + \delta) - E_N(t_0) = \mathcal{O}\left(\frac{1}{N^{1-\epsilon}} \tilde{\sigma}^2\right) + \mathcal{O}\left(\frac{1}{N^{C+1}} \tilde{\sigma}\right) + \mathcal{O}\left(\frac{1}{N^{2C+1}}\right).$$

To do so, consider $\Omega(I(Q(t) + \frac{k}{\sigma} w(t)))$ for $|k| \leq 1$. Recall that if f is a solution to the KdV, then

$$\Omega(I f(t)) = \langle I f_{xxx}, (I f)^2 - I f^2 \rangle - \langle (f^2)_x, I^2 f \rangle + \langle I(f^2)_x, (I f)^2 \rangle,$$

where $\langle \cdot, \cdot \rangle$ denotes the L^2 inner product. Therefore, $\Omega(I(Q(t) + \frac{k}{\sigma} I w(t)))$ is a polynomial in k . In addition, from the estimates in Lemma 3.2, which applies to $I(Q(t) + \frac{k}{\sigma} w(t))$ because $\|\frac{k}{\sigma} I w(t)\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} \ll 1$ for

$|k| < 1$, we may conclude that the coefficients of the polynomial

$$P_\delta(k) = 2 \int_{t_0}^{t_0+\delta} \Omega(IQ(t) + \frac{k}{\sigma} I w(t)) dt$$

are $\mathcal{O}(\frac{1}{N^{1-\epsilon}})$ so all the terms of second order or higher will validate the desired inequality automatically.

We therefore need only to check that the constant term is $\mathcal{O}(\frac{1}{N^{2C+1}})$ and the linear terms are $\mathcal{O}(\frac{1}{N^{C+1}})$. The constant term comes from

$$\Omega(I(Q(t))) = \langle IQ_t, IQ - IQ_{xx} - (IQ)^2 \rangle = \langle IQ_t, IQ^2 - (IQ)^2 \rangle,$$

which is $\mathcal{O}(\frac{1}{N^{2C+1}})$ because $IQ^2 - (IQ)^2 = IQ(I-1)Q + Q(I-1)Q + (1-I)Q^2$. But now note that because Q is Schwartz and $m(\xi) \equiv 1$ for $|\xi| \leq N$, $m(\xi) \leq 1$ for all ξ , we may conclude that $(I-1)Q = \mathcal{O}(N^{-2C})$ for any C we choose because $Q(t)$ is Schwartz in x . The same is true for Q^2 .

For the linear term, note that the linear term of $E_N(t)$ is given by:

$$E_N(t) = 2\langle IQ(t)_x, Iw(t)_x \rangle + 2\langle IQ(t), Iw(t) \rangle - 2\langle (IQ)^2(t), w(t) \rangle,$$

so the linear term of $\Omega(t)$ is:

$$\begin{aligned} \Omega(t) &= \frac{d}{dt} E_N(t) \\ &= 2\langle w_t, I(IQ^2 - (IQ)^2) \rangle + 2\langle w, \frac{d}{dt} (I(IQ^2 - (IQ)^2)) \rangle + \text{higher order terms} \\ &= -2\langle w_{xxx}, I(IQ^2 - (IQ)^2) \rangle + 2\langle w, \frac{d}{dt} (I(IQ^2 - (IQ)^2)) \rangle + \text{higher order terms}. \end{aligned}$$

We can thus bound those linear terms by: (after integrating by parts)

$$\|w\|_{L^2} (\|I(IQ^2 - (IQ)^2)\|_{H^3} + \|\partial_t(I(IQ^2 - (IQ)^2))\|_{L^2})$$

But now note again that because Q is Schwartz we may conclude that $\|IQ^2 - (IQ)^2\|_{H^s} \lesssim N^{-C}$ for any $s > 0$. The same is true for Q_t . Therefore, the linear terms of $\Omega(t)$ are controlled by $\|w\|_{L^2} N^{-C-1}$, and so, also using the fact that $\|w\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} \leq \sigma$, we conclude that

$$|E_N(t_0 + \delta) - E_N(t_0)| \lesssim \mathcal{O}(N^{-2C-1}) + \mathcal{O}(N^{-C-1} \tilde{\sigma}) + \mathcal{O}(N^{-1+\epsilon} \tilde{\sigma}^2)$$

which concludes the proof of Lemma 4.2. \square

5. Final Proof of the Main Theorem

In this section we will at last obtain the full power of Theorem 1.1:

THEOREM 5.1. *Let $0 \leq s < 1$, Let $\sigma = \text{dist}_{H^s}(u_0, \Sigma) \ll 1$, and let u be the solution to the KdV such that $u(\cdot, 0) = u_0$. Then $\text{dist}_{H^s}(u(t), \Sigma) \leq t^{1-s+\epsilon} \sigma$, for all t such that $t \ll \sigma^{-\frac{1}{1-s+\epsilon}}$.*

To do so, we will need to refine the choice of the soliton Q to which u is close. In the previous section, we chose a ψ to which u was close at time 0, and then assumed that u remained close to the soliton evolution of ψ over time. This required us to make use of the fact that $I\psi$ is close to ψ , which in turn forced us to require the condition $\sigma \gtrsim N^{-C}$ for some large C . We must eliminate this condition in order to obtain the full force of the theorem. We will therefore find a ψ^t which is close to u for each t , and study the equation by which this ψ^t moves in time. Define $\psi_0(x)$ to be the standard ground state solution to equation (1.2) centered at 0.

We begin by restating 1.4 in a form which will be more convenient:

LEMMA 5.2 (Weinstein, [14]). *Let $\psi \in \Sigma$, and let $w \in H^1$ such that $\|w\|_{H^1} \ll 1$ and $\langle w, (\psi^2)_x \rangle = 0$. Then*

$$\mathcal{L}(\psi + w) - \mathcal{L}(\psi_0) = \mathcal{L}(\psi + w) - \mathcal{L}(\psi) \sim \|w\|_{H^1}^2.$$

We will use the next lemma to find an appropriate ground state ψ for each t such that u is close to ψ and $w = u - \psi$ satisfies an appropriate orthogonality condition. Note that, since we will be studying Iw , not w , we will require $\langle Iw, (\psi^2)_x \rangle = 0$ instead of $\langle w, (\psi^2)_x \rangle = 0$.

LEMMA 5.3. *Let $u \in H^s$, and suppose $\text{dist}_{H^s}(u, \Sigma) \ll N^{s-1}$ with N sufficiently large. Then $u = \psi + w$ where ψ is a ground state, $\langle w, I(\psi^2)_x \rangle = 0$ and $\|Iw\|_{H^1} \lesssim N^{1-s} \text{dist}_{H^s}(u, \Sigma) \ll 1$.*

PROOF Define $d(u, \Sigma) = \|I(u - \psi)\|_{H^1}$. Then $d(u, \Sigma) \leq N^{1-s} \text{dist}_{H^s}(u, \Sigma) \ll 1$. So, as in [7], there exists a ψ' which minimizes $d(u, \Sigma)$. By the translation invariance of the problem, we may assume that this minimum occurs at ψ_0 . Note that the tangent space to Σ at ψ_0 is spanned by $\psi_{0,x}$. Therefore, if we differentiate

$$d(u, \psi)^2 = \langle I(u - \psi), I(u - \psi) \rangle + \langle \partial_x I(u - \psi), \partial_x I(u - \psi) \rangle$$

in the $\psi_{0,x}$ direction, we will get 0:

$$\langle I(u - \psi_0), I\psi_{0,x} \rangle + \langle \partial_x I(u - \psi_0), \partial_x I\psi_{0,x} \rangle = 0.$$

Let $\tilde{w} = u - \psi_0$. Then, since $\psi_0 - \psi_{0,xx} - \psi_0^2 = 0$, after integration by parts we get

$$\langle \tilde{w}, I^2(\psi_0^2)_x \rangle = 0.$$

This is almost what we want; we would like to replace the I^2 in the above equation by I . To do so, we will perturb ψ_0 slightly. Write $\psi = \psi_0(x - x_0)$, $w = u - \psi$ and $q = \psi - \psi_0$. We want to solve $\langle w, I(\psi^2)_x \rangle = 0$. Using what we know— $\langle \tilde{w}, I^2(\psi_0^2)_x \rangle = 0$ —and some algebra, what we want to solve for is:

$$\langle q, I((\psi_0 + q)^2)_x \rangle = \langle \tilde{w}, I((\psi_0 + q)^2)_x - I(\psi_0^2)_x \rangle + \langle I\tilde{w}, (\psi_0^2)_x - I(\psi_0^2)_x \rangle.$$

Note that the last term is $\mathcal{O}(N^{-100} \|I\tilde{w}\|_{H^1}) = \mathcal{O}(N^{-99} \text{dist}_{H^s}(u, \Sigma))$, because $I - 1$ is almost the identity on ψ_0 . For the left-hand side, note that $q = \psi - \psi_0 = -x_0 \psi_{0,x} + \mathcal{O}_{H^2}(|x_0|^2)$, where \mathcal{O}_{H^2} denotes the order of the H^2 norm of a function. Moreover, $((\psi_0 + q)^2)_x - (\psi_0^2)_x = 2x_0(\psi_{0,x}^2 + \psi \psi_{0,xx}) + \mathcal{O}_{H^2}(|x_0|^2)$. Therefore, the equation we wish to solve is

$$\langle x_0 \psi_{0,x} + \mathcal{O}_{H^2}(|x_0|^2), I((\psi_0 + q)^2)_x \rangle - x_0 \langle \tilde{w}, I(2\psi_{0,x}^2 + \psi_0 \psi_{0,xx} + \mathcal{O}_{H^2}(|x_0|^2)) \rangle = \mathcal{O}(N^{-99} \text{dist}_{H^s}(u, \Sigma)).$$

Since $\psi_{0,x}^2 + \psi_0 \psi_{0,xx}$ is Schwartz,

$$\langle I\tilde{w}, 2\psi_{0,x}^2 + \psi_0 \psi_{0,xx} \rangle = \mathcal{O}(\|I\tilde{w}\|_{H^1}) = \mathcal{O}(d(u, \Sigma)) \ll 1.$$

On the other hand,

$$\langle \psi_{0,x}, I(\psi_0^2)_x \rangle \sim \|\psi_0\|_{W^{2,4}}^2 + \mathcal{O}(N^{-100}),$$

which is an absolute constant that is not close to zero. So in the end, we get

$$x_0(\langle \psi_{0,x}, I(\psi_0^2)_x \rangle - \langle I\tilde{w}, 2\psi_{0,x}^2 + \psi_0 \psi_{0,xx} \rangle) = \mathcal{O}(N^{-99} \text{dist}_{H^s}(u, \Sigma)) + \mathcal{O}(|x_0|^2),$$

where the coefficient of x_0 on the left-hand side is close to a constant independent of \tilde{w} . Therefore, by the inverse function theorem, we find that there is an $x_0 \sim \mathcal{O}(N^{-99} \text{dist}_{H^s}(u, \Sigma))$ which solves this equation, and then since $\|\psi - \psi_0\|_{H^2} = \mathcal{O}(N^{-99} \text{dist}_{H^s}(u, \Sigma))$ the functions $\psi = \psi_0(x - x_0)$ and $w = u - \psi$ will satisfy all the desired conditions. \square

We apply this lemma at each time t such that $\text{dist}_{H^s}(u, \Sigma) \ll N^{s-1}$ to write $u(x, t) = \psi^t(x) + w(x, t)$. We will redefine $Q(x, t)$ by:

$$u(x, t) = Q(x, t) + w(x, t) = \psi_0(x - t - x_0(t)) + w(x, t)$$

For this section, we will redefine $E_N(t)$ in order to eliminate our dependence on the closeness of ψ and $I\psi$ and to reflect the more precisely chosen error function $w(t)$ found in the above lemma. We therefore set⁷

$$(5.1) \quad E_N(t) = \mathcal{L}(Q(t) + Iw(t)).$$

Note that, by (5.3), for each t such that $\text{dist}_{H^s}(u, \Sigma) \ll N^{s-1}$, $\langle Iw(t), (Q^2(t))_x \rangle = 0$ and $\|Iw\|_{H^1} \ll 1$. Therefore, by 5.2, $|E_N(t) - \mathcal{L}(Q(t))| \sim \|Iw\|_{H^1}^2$. In particular, at $t = 0$, we have

$$|E_N(0) - \mathcal{L}(Q(0))| \sim \|Iw\|_{H^1}^2 \lesssim N^{2-2s}\sigma^2$$

To prove the theorem, we will need the following lemma, a refinement of Lemmas 3.2 and 4.2:

LEMMA 5.4. *Suppose there is a $t_0 \in \mathbb{R}$ and a $\tilde{\sigma}$ with $0 < \tilde{\sigma} \ll 1$ such that $|E_N(t_0) - \mathcal{L}(\psi_0)| \lesssim \tilde{\sigma}^2$. Then there exists a $\delta > 0$ depending only on s such that*

$$E_N(t_0 + \delta) - E_N(t_0) = \mathcal{O}\left(\frac{1}{N^{1-\epsilon}} \tilde{\sigma}^2\right).$$

We will assume this lemma for now and conclude the proof of the theorem:

PROOF (of Theorem 5.1) Once again, we set $\tilde{\sigma} = N^{1-s}\sigma$. As in the proofs of Propositions 3.1 and 4.1, we can iterate the result of Lemma 5.4. In this case, for $Q(x, t) = \psi_0(x - t - x_0(t))$, we obtain that $|E_N(t) - \mathcal{L}(Q(t))| \lesssim N^{2-2s}\sigma^2$ for all t such that $t \ll N^{1-\epsilon}$. So, by Lemma 5.2,

$$\text{dist}_{H^s}(u, \Sigma) \lesssim \|w\|_{H^s} \lesssim N^{1-s}\sigma,$$

for all $t \ll N^{1-\epsilon}$. We therefore can optimize for N under only the two conditions:

$$(5.2) \quad t \ll N^{1-\epsilon} \quad N^{2-2s}\sigma^2 \ll 1.$$

Contrast these conditions with (3.2) and (4.1). Note that we have now eliminated the condition $\sigma \ll N^{-C}$ and therefore we obtain

$$\text{dist}_{H^s}(u, \Sigma) \lesssim t^{1-s+\epsilon}\sigma,$$

for all $t \ll \sigma^{\frac{1}{1-s-\epsilon}}$, as claimed. \square

It thus remains only to prove Lemma 5.4:

PROOF (of Lemma 5.4) We write $Q(x, t) = \psi_0(x - t - x_0(t))$ and $w(x, t) = u(x, t) - Q(x, t)$. Then $w(t)$ satisfies the difference equation:

$$(5.3) \quad w_t + w_{xxx} + \partial_x(w(w + 2Q)) + \dot{x}_0 Q_x = 0^8$$

We know that $\|Iw(t_0)\|_{H^1} \lesssim \tilde{\sigma} = N^{1-s}\sigma$. As before, we start by proving that the $X^{1, \frac{1}{2}+\epsilon}$ norm of Iw is controlled.

CLAIM 3.

$$\|Iw\|_{X^{1, \frac{1}{2}+\epsilon}_{[t_0-\delta, t_0+\delta]}} \lesssim \tilde{\sigma}.$$

⁷Compare to (3.1).

⁸We use the notation \dot{x}_0 to mean the ordinary derivative $\frac{dx}{dt}$.

PROOF As in each of the two previous claims, we use the standard $X^{s,b}$ estimates to obtain:

$$\begin{aligned} \|Iw\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} &\lesssim \|Iw(t_0)\|_{H^1} + \delta^\epsilon \|I(w_t + w_{xxx})\|_{X_{[t_0-\delta, t_0+\delta]}^{1, -\frac{1}{2}+2\epsilon}} \\ &\lesssim \tilde{\sigma} + \delta^\epsilon \|I(w(w + 2Q))_x\|_{X_{[t_0-\delta, t_0+\delta]}^{1, -\frac{1}{2}+2\epsilon}} + \delta^\epsilon \|\dot{x}_0(t)Q_x\|_{X_{[t_0-\delta, t_0+\delta]}^{1, -\frac{1}{2}+2\epsilon}}. \end{aligned}$$

Note that the first term on the right-hand side is the same as in Claim 2 and can be estimated in exactly the same way. For the second term, we will prove that for each $t \in [t_0 - \delta, t_0 + \delta]$, $\|\dot{x}_0(t)\| \lesssim \|Iw(t)\|_{H_x^1}$. Then we will have

$$\begin{aligned} \|\dot{x}_0(t)Q_x\|_{X_{[t_0-\delta, t_0+\delta]}^{1, -\frac{1}{2}+2\epsilon}} &\lesssim \|Iw(t)\|_{H^1} Q_x\|_{X_{[t_0-\delta, t_0+\delta]}^{1, -\frac{1}{2}+2\epsilon}} \\ &= \left\| \left(\frac{\langle \xi \rangle}{\langle \tau - \xi^3 \rangle^{\frac{1}{2}-2\epsilon}} \left(\widehat{Iw}_{H_x^1} *_{\tau} \tilde{Q}_x(\xi) \right) (\tau) \right) \right\|_{L_{\tau}^2 L_{\xi}^2} \\ &\leq \| \widehat{Iw}_{H_x^1} \|_{L_{\tau}^1} \left\| \frac{\langle \xi \rangle}{\langle \tau - \xi^3 \rangle^{\frac{1}{2}-2\epsilon}} \tilde{Q}_x(\xi, \tau - a) \right\|_{L_a^\infty L_{\xi, \tau}^2} \\ &\leq C \|Iw\|_{L_{t, [t_0-\delta, t_0+\delta]}^\infty H_x^1} \\ &\leq C \|Iw\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}}. \end{aligned}$$

The third line makes use of Minkowski's inequality for integrals, and the fourth takes advantage of the fact that $Q(x, t)$ and all of its x -translates are uniformly bounded in $X^{1, -\frac{1}{2}+2\epsilon}$ space. The last step is due to the standard estimate $\|f\|_{L_t^\infty H_x^1} \lesssim \|f\|_{X^{1, \frac{1}{2}+\epsilon}}$. This argument allows us to conclude that:

$$\|Iw\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} \lesssim \tilde{\sigma} + \delta^\epsilon \|I(w(w + 2Q))_x\|_{X_{[t_0-\delta, t_0+\delta]}^{1, -\frac{1}{2}+2\epsilon}} + C\delta^\epsilon \|Iw\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}},$$

and we can then complete the proof of the claim via a continuity argument. Therefore, to check that $\|Iw\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} \lesssim \tilde{\sigma}$, we need only prove that, for each t , $|\dot{x}_0(t)| \lesssim \|Iw(t)\|_{H_x^1}$.

To do so, write

$$\theta(x, t) = w(x + t + x_0(t), t) = u(x + t + x_0(t), t) - Q(x + t + x_0(t), t) = u(x + t + x_0(t), t) - \psi_0(x)$$

Then θ satisfies:

$$\theta_t + \theta_{xxx} + (\theta(\theta + 2Q))_x = \dot{x}_0(t)u_x + \theta_x.$$

Recall that w satisfies $\langle w, I(Q^2)_x \rangle = 0$. Differentiating in time, we see that, for each t ,

$$\langle \theta_t, I(\psi_0^2)_x \rangle = 0$$

Plugging in for θ_t and simplifying, we obtain:

$$\dot{x}_0(t) \langle u_x, I(\psi_0^2)_x \rangle = \langle \theta_{xxx}, I(\psi_0^2)_x \rangle + \langle (\theta(\theta + 2\psi_0))_x, I(\psi_0^2)_x \rangle + \langle \theta_x, I(\psi_0^2)_x \rangle,$$

i.e.

$$\dot{x}_0(t) = \frac{1}{\langle \psi_{0,x} + \theta_x, I(\psi_0^2)_x \rangle} \left(\langle I\theta_x, (\psi_0^2)_{xxx} \rangle + \langle (\theta + 2\psi_0)(\psi_0^2)_x + (\psi_0^2)_x, \theta \rangle + \langle \theta, I(\theta + 2\psi_0)_x(\psi_0^2)_x \rangle \right).$$

Note that the numerator is controlled by $\|I\theta\|_{H^1}$ and that the denominator is of a size greater than an absolute constant. Therefore, we conclude that $\dot{x}_0(t)$ is indeed controlled by $\|I\theta(t)\|_{H^1} = \|Iw(t)\|_{H^1}$ as claimed. \square

The final step in the proof of the lemma is to take a δ step forward in t . We want to prove

$$E_N(t_0 + \delta) - E_N(t_0) = \mathcal{O}\left(\frac{1}{N^{1-\epsilon}} \tilde{\sigma}^2\right).$$

Recall that $E_N(t) = \mathcal{L}(Q + Iw) = \mathcal{L}(Q(x + t, t) + Iw(x + t, t))$. Also recall that $\Omega(f)(t) = \partial_t(\mathcal{L}(f)(t)) = 2\langle f_t, f - f_{xx} - f^2 \rangle$. Therefore,

$$\begin{aligned} \partial_t E_N(t) &= \Omega((Q + Iw)(x + t, t)) \\ &= \langle \partial_t((Q + Iw)(x + t, t)), Q + Iw - Q_{xx} - Iw_{xx} - 2QIw - (Iw)^2 - Q^2 \rangle \\ &= 2\langle -\dot{x}_0(t)Q_x + I(-w_{xxx} - (w(w + 2Q))_x + \dot{x}_0(t)Q_x), Iw - Iw_{xx} - (Iw)(2Q + Iw) \rangle \\ &= 2\langle \dot{x}_0(t)(IQ_x - Q_x), (Iw - Iw_{xx} - (Iw)(2Q + Iw)) \rangle + \\ &\quad \langle I(w_x - w_{xxx} - (w(w + 2Q))_x), Iw - Iw_{xx} - (Iw)(2Q + Iw) \rangle. \end{aligned}$$

By integration by parts and the fact that I is almost the identity on Q , the first term (when integrated in t) will be controlled by $CN^{-100}\|Iw\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}}^2$ and will therefore be fine for our estimates.

Note also that the second term is a polynomial of degree at least 2 in w . Therefore, as in Section 4, we will be done if we can prove that for all γ such that $\|\gamma\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} \leq 1$,

$$\int_{t_0}^{t_0+\delta} \langle I(\gamma_x - \gamma_{xxx} - (\gamma(\gamma + 2Q))_x), I\gamma - I\gamma_{xx} - (I\gamma)(2Q + I\gamma) \rangle \lesssim \frac{1}{N^{1-\epsilon}}.$$

To do so, let $v = \gamma + Q$. Then

$$\gamma_x - \gamma_{xxx} - (\gamma(\gamma + 2Q))_x = v_x - v_{xxx} - (v^2)_x$$

and

$$I\gamma - I\gamma_{xx} - (I\gamma)(2Q + I\gamma) = Iv - Iv_{xx} - (Iv)^2 + 2(Iv)(IQ - Q) - (IQ - Q)^2 - (IQ^2 - Q^2).$$

Therefore

$$\begin{aligned} &\int_{t_0}^{t_0+\delta} \langle I(\gamma_x - \gamma_{xxx} - (\gamma(\gamma + 2Q))_x), I(\gamma - \gamma_{xx} - \gamma(2Q + I\gamma w)) \rangle dt = \\ &= \int_{t_0}^{t_0+\delta} \langle I(v_x - v_{xxx} - (v^2)_x), ((Iv - Iv_{xx} - (Iv)^2) + 2(Iv)(I - 1)Q + ((I - 1)Q)^2 + (I - 1)Q^2) \rangle dt. \end{aligned}$$

Then, once again, because $I - 1$ is nearly 0 on Q , the second third and fourth terms are controlled. For the remaining term, note that, by integration by parts $\langle Iv_x, Iv - Iv_{xx} - (Iv)^2 \rangle$ is zero, so the last term to be estimated is

$$(5.4) \quad \int_{t_0}^{t_0+\delta} \langle I(-v_{xxx} - (v^2)_x), Iv - Iv_{xx} - (Iv)^2 \rangle dt.$$

But this is exactly the quantity estimated in Lemma 3.2. Recall that the multilinear estimates used to prove those estimates did not depend on the properties of the function u except that $\|u\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}} \lesssim 1$.

The conclusion was that

$$\int_{t_0}^{t_0+\delta} \langle I(-v_{xxx} - (v^2)_x), Iv - Iv_{xx} - (Iv)^2 \rangle dt = \mathcal{O}\left(\frac{1}{N^{1-\epsilon}}\right).$$

Since $\|v\|_{X_{[t_0-\delta, t_0+\delta]}^{1, \frac{1}{2}+\epsilon}}$ is indeed controlled by a constant, by the estimates in the proof of Lemma 3.2 the quantity (5.4) is also controlled by $\frac{1}{N^{1-\epsilon}}$. This concludes the proof of Lemma 5.4 and, at last, the main theorem. \square

References

- [1] J. Bourgain, *Fourier Restriction Phenomena for Certain Lattice Subsets and Applications to Nonlinear Evolution Equations II: The KdV Equation*, Geom. Funct. Anal., **3** (1993), no. 3, 209-262.
- [2] J. Bourgain, *Refinements of Strichartz' inequality and applications to 2D-NLS with critical nonlinearity*, International Mathematical Research Notices, **5** (1998), 253-283.
- [3] T. Benjamin, *The Stability of Solitary Waves*, Proc. R. Soc. Lond. A. **328** (1972), 153-183.
- [4] Bona, J. *On the Stability Theory of Solitary Waves*, Proc. R. Soc. Lond. A **344** (1975), 363-374.

- [5] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. *Global Well-Posedness for KdV in Sobolev Spaces of Negative Index*, J. Differential Equations (2001), no. 26, 7 pp. (electronic).
- [6] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. *Multilinear Estimates for Periodic KdV Equations, and Applications*, to appear in *J. Funct. Anal.*, 2002.
- [7] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. *Polynomial Upper Bounds for the Orbital Instability of the 1D Cubic NLS Below the Energy Norm*, to appear in *DCDS*, 2002.
- [8] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. *Global Well-Posedness for Schrödinger Equations with Derivatives*, SIAM J. Math. Anal. (2001), no. 3, 649-669.
- [9] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. *A refined global well-posedness result for Schrödinger equations with derivatives*, to appear in *SIAM, J. MATH. Anal.*, 2002.
- [10] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao. *Sharp global well-posedness for KdV and Modified KdV on \mathbb{R} and \mathbb{T}* , to appear in *J. of the AMS*, 2002.
- [11] C. Kenig, G. Ponce, and L. Vega. *A Bilinear Estimate with Applications to the KdV Equation*, J. Amer. Math. Soc. **9** (1996), 573-603.
- [12] C. Kenig, G. Ponce, and L. Vega. *Oscillatory Integrals and Regularity of Dispersive Equations*, Indiana University Math J., **40**(1991), 33-69.
- [13] F. Merle and L. Vega. *L^2 Stability of Solitons for KdV Equation*, preprint.
- [14] M. Weinstein. *Lyapunov Stability of Ground States of Nonlinear Dispersive Evolution Equations*, Comm. Pure Appl. Math. **39** (1986), 51-68.

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

E-mail address: sraynor@math.mit.edu and gigliola@math.mit.edu