

Barrier Islands: Coupling Anthropogenic Stability with Ecological Sustainability

Rusty A. Feagin^{†*}, William K. Smith[‡], Norbert P. Psuty[§], Donald R. Young^{††}, M. Luisa Martínez^{‡‡}, Gregory A. Carter^{§§}, Kelly L. Lucas^{§§}, James C. Gibeaut^{†††††}, Jane N. Gemma^{§§§}, and Richard E. Koske^{§§§}

[†]Department of Ecosystem Science & Management
Texas A&M University
College Station, TX 77845,
U.S.A.
feaginr@tamu.edu

[‡]Department of Biology
Wake Forest University
Winston-Salem, NC 27109,
U.S.A.

[§]Institute of Marine and Coastal Sciences
Rutgers University
Sandy Hook, NJ 07732, U.S.A.

^{††}Department of Biology
Virginia Commonwealth University
Richmond, VA 23284, U.S.A.

^{‡‡}Red de Ecología Funcional
Instituto de Ecología
Xalapa 91070, Mexico

^{§§}Gulf Coast Geospatial Center
University of Southern Mississippi
Gulfport, MS 39501, U.S.A.

^{†††}Harte Research Institute for Gulf of Mexico Studies
Texas A&M University-Corpus Christi
Corpus Christi, TX 78412,
U.S.A.

^{§§§}Bureau of Economic Geology
University of Texas
Austin, TX 78758, U.S.A.

^{§§§}Department of Biological Sciences
University of Rhode Island
Kingston, RI 02881, U.S.A.

ABSTRACT



FEAGIN, R.A.; SMITH, W.K.; PSUTY, N.P.; YOUNG, D.R.; MARTÍNEZ, M.L.; CARTER, G.A.; LUCAS, K.L.; GIBEAUT, J.C.; GEMMA, J.N., and KOSKE, R.E., 2010. Barrier islands: coupling anthropogenic stability with ecological sustainability. *Journal of Coastal Research*, 00(0), 000-000. West Palm Beach (Florida), ISSN 0749-0208.

Barrier islands provide a host of critical ecosystem services to heavily populated coastal regions of the world, yet they are quite vulnerable to ongoing sea level rise and a potential increase in the frequency and intensity of oceanic storms. These islands are being degraded at an alarming rate, in part because of anthropogenic attempts at stabilization. In this article, we outline a possible sustainability strategy that incorporates the natural degree of substrate instability on these sedimentary landscapes. We recommend placing the focus for managing barrier islands on maintaining ecosystem function and process development rather than emphasizing barrier islands as structural impediments to wave and storm energy.

ADDITIONAL INDEX WORDS: *Barrier islands, coastal erosion, storm surge, sea level rise, coastal management, vegetation.*

INTRODUCTION

Barrier islands are ecosystems that border coastal shorelines and physically separate the offshore oceanic province from inshore wetlands, bays, sounds, and estuaries. As their name implies, they form a protective barrier between continental shorelines and wave action that originates offshore. Barrier islands also provide the structural framework for the formation of an array of coastal and estuarine habitats that host a variety of native and migratory species, many of which have substantial economic value.

Coastlines fronted by barrier islands also include some of the greatest concentrations of human populations and accompanying anthropogenic development in the world (Schlacher *et*

al., 2007; Weinstein *et al.*, 2007). The native vegetation and geological stability of these ecosystems are coupled and vulnerable to erosion events, particularly when also disturbed by development. As a result, barrier islands are quite vulnerable to potential global warming impacts such as sea level rise (Intergovernmental Panel On Climate Change, 2007) and increase in the frequency and intensity of major oceanic storms (Emanuel, 2005; Webster *et al.*, 2004). These islands are some of the most valuable yet potentially vulnerable ecosystems on Earth (Pérez-Maqueo, Intralawan, and Martínez, 2007; U.S. Commission On Ocean Policy, 2004).

Our goal here is to outline a sustainability strategy that recognizes and incorporates the high degree of substrate instability found in these natural ecosystems. An ideal solution would involve a strategic compromise between anthropogenic development and preservation of the natural ecosystem.

DOI: 10.2112/09-1185.1 received 2 October 2009; accepted in revision 1 November 2009.

1

*

NATURAL INSTABILITY

Barrier islands have an extremely dynamic nature whereby major changes in geomorphology and hydrology can occur over days, or even hours, in response to extreme episodic storm events (EESes) such as tropical cyclones, hurricanes, and northeasters. Of most interest to the management community are changes that occur in response to these natural hazards. Barrier island width, dune elevation, tidal prism, wave energy, and storm surge energy influence the likelihood of overwash (Claudino-Sales, Wang, and Horwitz, 2008; Leatherman, Williams, and Fisher, 1977; Morton and Sallenger, 2003), transport of sediment offshore, and formation of new inlets during storms (Fitzgerald and Van Heteren, 1999). An array of endemic, indigenous, and migratory species are adapted specifically to such transient geological conditions (Brown and MacLachlan, 2002; Ehrenfeld, 1990; Shao, Shugart, and Hayden, 1996). However, little is known concerning the nature of the adaptive physiological mechanisms associated with this high level of transience, especially in response to EESes, although these species are expected to have unique suites of evolutionary, genetic traits that underlie these adaptations (Gutschick and Bassirirad, 2003).

It is well documented that coastal plants are specifically adapted to take advantage of the dynamic nature of sediment transport in response to long-term processes, such as sea level rise or changes in sediment budget, having become adept “ecological engineers” (Costanza, Mitsch, and Day, 2006; Crain and Bertness, 2006; Halpern *et al.*, 2007; Jones, Lawton, and Shachak, 1997). In classical works on both sand dunes (Cowles, 1899) and salt marshes (Redfield and Rubin, 1962), ecologists have shown that these plants capture sediment, elevate the substrate, and drive the successional process. For example, dunes are the result of sand accumulation, becoming larger as plant sizes, densities, and root depths increase (Tsoar, 2005) and mutualist interactions begin with mycorrhizal fungi (Koske *et al.*, 2004). As succession progresses, primary and secondary dunes aggrade through further sand deposition, eventually leading to the formation of maritime forests on elevated upland areas (Young, Shao, and Brinson, 1995). There is a great deal of research on the ecological engineering capabilities of coastal plants, yet only a few studies have explored the eco-geomorphological limits of these capabilities (Feagin, Sherman, and Grant, 2005; Kirwan and Murray, 2007; Morris *et al.*, 2002; Orson, Warren, and Niering, 1998) and none have evaluated explicitly how this engineering can mediate the quite different time scales of physical forcing experienced by barrier islands, *i.e.*, sea level rise vs. EESE.

ANTHROPOGENIC STABILITY

One of the most conspicuous boundaries between landscapes modified by humans and those formed by natural processes occurs at the seaward edge of private property adjacent to beaches (Mitteager, Burke, and Nordstrom, 2006). For example, in the United States, nearly all state laws created in response to the federal Coastal Zone Management Act of 1972 demarcate the public-private property division as the high tide line, resulting in a desire by landowners to “hold the line”

between the beach and their backyards (Archer *et al.*, 1994). In contrast, the *Ley de Costas* (Law of Coasts) in Spain states that there should be a minimum distance of 100 m between shoreline and human structures. When necessary for effective shoreline protection, another 100 m can be added to this limit. Under this law, all sandy beaches and coastal dunes are public domain (Ley 22/1988, de 28 de Julio, de Costas). In India, there is currently a similar political effort to legally define the coastal zone and place it under federal jurisdiction (Sridhar *et al.*, 2006). A similar situation occurs in Mexico, where the coastal zone is also clearly defined (although routinely defied by private stakeholders). Ultimately, static legal definitions of the coastal zone enforce linear restrictions to the natural interplay of sediments and represent a threat to ecosystem functioning. Also, inevitable conflicts ensue once sea levels rise or EESes strike.

While such laws attempt to manage human interactions with the land-sea interface, they are fundamentally rooted in the anthropogenic desire for stability. In the U.S. example, this leads to patchy vegetative stabilization that is most often managed privately and a lack of any governmental control over the larger landscape-scale sedimentary dynamics since dune formation would occur landward of the high tide line. In contrast, in Spain, because the *Ley de Costas* defines this line further landward, a large number of private houses (300,000) located at the beach are likely to be bought by the government to enable such control to be asserted. Returning to the India example, its laws are currently espoused as a response to the 2004 Indian Ocean tsunami and have facilitated a vast effort toward planting nonnative forests as “bioshields” for stabilization of the coastal areas (Mukherjee *et al.*, 2009). Ultimately, this approach to stabilization impairs the long-term functioning of the ecosystem, primarily because nonnative species are not adapted to the dynamism of the sedimentary environment and are unable to promote accretion (Feagin, Mukherjee, *et al.*, 2009). Furthermore, this new demarcation of the coastal zone by the government effectively results in the resale of land to private entities (Rodriguez *et al.*, 2008) and allows the undocumented and customary land rights of indigenous residents to disappear (Menon and Sridhar, 2007). In Mexico, development for tourism is intensive and extensive, and the laws that do exist are not enforced.

ECOLOGICAL SUSTAINABILITY

What is the minimum level of landscape stability needed for human occupation of these naturally dynamic ecosystems? Is there a compromise solution that will incorporate both anthropogenic desires for structural stability and natural substrate instability of the system (Nordstrom, 2008) and that will lead to ecosystem sustainability? While precise answers to these questions do not currently exist, functional solutions will come by acknowledging that the dynamism and transience of these ecosystems at the landscape scale are paramount and overriding (Figure 1). Yet, how do we implement such a management strategy?

To begin, we must stop biasing our decisions toward the use of human-engineered structures on coasts at the expense of using native vegetation to achieve the same purpose (Nord-




Figure 1. "Sustainability" and "stability" are often contradictory terms, as shown on a cross-section of a typical barrier island landscape. Native vegetation is adapted to the movement of sediments during extreme episodic storm events (EESE) and builds elevations through the process of succession, countering relative sea level rise (RSLR). Anthropogenic barriers such as paved surfaces, housing, hotels, bulkheads, and seawalls inhibit the natural sedimentary exchange between the land and the sea, interrupt ecosystem migration and processes of "ecological engineering" by plants, and result in net sediment and elevation loss.

strom, 2000; Psuty, 2004). It is much less expensive and easier to guide vegetative succession (manage productivity and species mix) and coupled sedimentary processes (accretion) to build the elevation and structure of a landscape and thus counter selected long-term processes, such as sea level rise.

Second, we should begin to investigate more deeply the hypothesis that vegetated barrier island ecosystems can modify and control the sedimentary dynamics in response to gradual forcings like sea level rise (as long as the rate is not too fast) but cannot resist discrete, intense forcings such as EESEs (Brinson, Christian, and Blum, 1995; Jentsch, Kreyling, and Beierkuhnlein, 2007; Michener *et al.*, 1997). Recent work supports this hypothesis for salt marsh ecosystems (Feagin, Lozada-Bernard, *et al.*, 2009). For sand dune or other ecosystems where vegetation has appeared to reduce landscape-scale erosion in response to EESEs (Figure 2), it is important to understand whether the mechanism involved is primarily the direct attenuation of wave energy by plant cover or is more directly a function of the higher substrate elevations engineered by plants prior to the storm. If plants reduce the surge water level during EESEs (not just the wave height-to-water depth ratio), then more inductive field work is needed to quantify this effect during and following EESEs (Feagin, Mukherjee, *et al.*, 2009). If the mechanism of protection or erosion mitigation is primarily a function of substrate elevation, then substrate accumulation (indirect protection effect by ecological engineering over time) is paramount and must receive more research focus (Feagin, 2008; Feagin, Lozada-Bernard, *et al.*, 2009). If the hypothesis that plants can manage gradual physical changes in the sedimentary environment up to some threshold force (which EESEs exceed) is generally correct for barrier island species, then ecosystem management should emphasize landscape-scale sedimentary modification rather than short-term, structural prevention of erosion at a specific location.




Figure 2. Did vegetation *directly* reduce erosion during Hurricane Katrina? Areas on Dauphin Island, Alabama, with native vegetation (red in top images, brown-green in bottom images) appeared to have less overwash (left images) than areas that had been disturbed by anthropogenic development (right images). Was there less erosion in the undeveloped areas primarily because of the wave energy attenuated by the plants during the storm, or was it primarily a function of the high dunes created by plant succession prior to the storm? The answer has serious management implications, as described in the text. (Courtesy of National Aeronautics and Space Administration and Alabama Department of Economic and Community Affairs.)

Third, from an economic perspective, vegetation management should not focus on stabilization during EESEs. We know that coastal ecosystems contribute 77% of global ecosystem services, a value of about \$33 trillion *per year* (Martínez *et al.*, 2007), and barrier islands are an integral part of approximately 12% of these coastal ecosystems (Pilkey and Fraser, 2003). Moreover, beaches and dunes support a U.S. tourism industry valued at \$322 billion *per year*, more than 25 times the contribution of the National Park Service system to the U.S. economy (Houston, 2008). Engineered protection structures, dikes, seawalls, houses, buildings, and plantations of nonnative species do provide excellent stabilization for this valuable landscape. Land is limited and extremely valuable on barrier islands, and if better stabilization can be had with a seawall or dike that occupies less space than natural defenses such as dunes or wetlands, and the land's property value outweighs its ecosystem service value, then barrier islands inevitably will be developed. For example, the substrate stability and property value of barrier islands could potentially increase if we backfilled them to above the height of the native vegetation and then designed parking lots and buildings to withstand

EESEs. However, these important ecosystems will be lost if all valuation is placed upon direct storm protection or stabilization value, simply because the land is worth so much to developers. Thus, we need to explore economic valuation techniques that can account quantitatively for the value of ecosystem services that manage the natural instability over the long term, thereby sustaining the entire island metastructure rather than only site-specific structural impediments for deflecting storm energy or stabilizing substrate (Williams *et al.*, 2009).

Fourth, we need to develop laws analogous to Section 404 of the Clean Water Act (which protects wetlands from being developed or backfilled) to protect undeveloped beaches, sand dunes, maritime forests, and other critical barrier island habitats. Because these landscapes are being developed at such a fast rate, and the economic calculus for their land use is currently biased toward anthropogenic development, their preservation for posterity should be a legislative priority.

Fifth, we need to bring about a change in legal mindset about how to actively manage developed barrier islands. We must reconfigure the legal conception of these landscapes, rather than placing a linear boundary between public and private property upon them. Setbacks are certainly an option (Pilkey and Young, 2005) in the context of modified state laws in the United States, similar to the Ley de Costas in Spain, but we need to think creatively about using ecologically defined boundaries. For example, the state of Texas defines the public-private property line as the ecological reality of the native vegetation line (Open Beaches Act, Texas Natural Resources Code §61.011; see Feagin, 2005). While this law has several gaps that need to be addressed, it allows a functioning dune ecotone to form that provides energy absorption between the beach and the leading edges of personal property, enhancing the ecological resistance and resilience (recovery rate) of a developed shoreline to EESEs. Further by law, when this sand dune plant zone and vegetation line shifts landward, so does the location of the public-private boundary. With this approach, land speculators and real estate developers must accommodate the reality of a retreating beach ecotone. Governmental entities could also provide tax incentives for more mobile living structures that will accommodate shifting substrates on at least a decadal scale. Statutory modification of the public-private line in currently undeveloped areas could be a major goal for coastal managers and policymakers.

Sixth, federal governments should purchase as many undeveloped islands and contiguous marginal properties as possible. In the United States, beyond the original Coastal Barrier Resources Act of 1982 that prevented federal assistance for activities supporting commercial development of barrier islands and designated certain parklands and national seashores to be preserved, no federal framework exists today for sustaining these ecosystems. In the United States, many barrier island ecosystems are still undeveloped. Extending federal ownership to these undeveloped properties will be less expensive than using taxpayer monies (under the National Flood Insurance Program and other granting programs, subsidies, and tax incentives; *e.g.*, Bagstad, Stapleton, and D'agostino, 2007; Cordes and Yezer, 1998) to subsidize recovery efforts for vacation homes that will inevitably be blown or eroded away again. To reverse this tragedy of the commons, a

large-scale declaration of eminent domain may be required, as is currently occurring in Spain. Undoubtedly, such a declaration will find steep resistance in a society founded upon a strong notion of individual property rights. Compromises could be made whereby we work to armor our most developed and valuable properties against repeated flood losses due to EESEs and simultaneously declare undeveloped barrier islands off-limits to new permanent structures and subsidized flood insurance. One possibility for the United States could include tying annual funds, grants, or subsidies related to the Coastal Zone Management Act, National Flood Insurance Program, and National Oceanic and Atmospheric Administration, by state, to the amount of barrier island land that has been contributed to federal ownership. In locations such as India where coastal zone laws are still being developed, such programs should respect the customary yet undocumented uses of land, pay a fair rate for the land, and occur only in undeveloped locations. Any acquired land should be used only for public good and should remain nontransferable to private entities.

Seventh, we need to begin rethinking our cultural view of these ecosystems. The challenge may require that we adopt only a new vision of these landscapes rather than any major restructuring of the landscapes. Why did we start calling these features "barrier islands" instead of "ephemeral islands" or "migrating islands"? Although we recognize that past lexicon is difficult to change, how might something as simple as a different name have affected our desires to stabilize these islands? We will need to deepen our understanding and alter our management practices to ensure their future sustainability.

CONCLUSIONS

Barrier islands are naturally unstable, migrating, and changing in response to factors such as sea level rise and EESEs. They are now being degraded at an alarming rate, in part because of human efforts to force stability upon them. We should aim for sustaining the natural sedimentary processes that occur across barrier island landscapes, rather than trying only to stabilize the substrate—we must find a way to adapt to the dynamism.

ACKNOWLEDGMENTS

This paper was generated from a workshop (April 14–16, 2006, in Biloxi, Mississippi, USA) that organized the Coastal Barrier Island Network, a multidisciplinary group concerned with the preservation of barrier islands (<http://www.coastalbarrierisland.org/>). Support was provided by the National Science Foundation, Research Coordination Networks in Biological Sciences (NSF-RCN, grant numbers IOB 0607921 and DBI 0741928).

LITERATURE CITED

- Archer, J.H.; Connors, D.L.; Laurence, K.; Columbia, S.C., and Bowen, R., 1994. *The Public Trust Doctrine and the Management of America's Coasts*. Boston: University of Massachusetts Press.
- Bagstad, K.J.; Stapleton, K., and D'Agostino, J.R., 2007. Taxes,

- subsidies, and insurance as drivers of United States coastal development. *Ecological Economics*, 63, 285–298.
- Brinson, M.M.; Christian, R.R., and Blum, L.K., 1995. Multiple states in the sea-level induced transition from terrestrial forest to estuary. *Estuaries*, 18, 648–659.
- Brown, A.C. and McLachlan, A., 2002. Sandy shore ecosystems and the threats facing them: some predictions for the year 2025. *Environmental Conservation*, 29, 62–77.
- Claudino-Sales, V.; Wang, P., and Horwitz, M.H., 2008. Factors controlling the survival of coastal dunes during multiple hurricane impacts in 2004 and 2005: Santa Rosa barrier island, Florida. *Geomorphology*, 95, 295–315.
- Cordes, J.J. and Yezer, A.M.J., 1998. In harm's way: does federal spending on beach enhancement and protection induce excessive development in coastal areas? *Land Economics*, 74, 128–145.
- Costanza, R.; Mitsch, W.J., and Day, J.W., Jr., 2006. A new vision for New Orleans and the Mississippi Delta: applying ecological economics and ecological engineering. *Frontiers in Ecology and the Environment*, 4, 465–472.
- Cowles, H.C., 1899. The ecological relations of the vegetation of the sand dunes of Lake Michigan. *Botanical Gazette*, 27, 95–117, 167–202, 281–308, 361–391.
- Crain, C.M. and Bertness, M.D., 2006. Ecosystem engineering across environmental gradients: implications for conservation and management. *Bioscience*, 56, 211–218.
- Ehrenfeld, J.G., 1990. Dynamics and processes of barrier island vegetation. *Aquatic Sciences*, 2, 437–480.
- Emanuel, K., 2005. Increasing destructiveness of tropical cyclones of the last 30 years. *Nature*, 436, 686–688.
- Feagin, R.A., 2005. Artificial dunes created to protect property in Galveston, Texas: the lessons learned. *Ecological Restoration*, 23, 89–94.
- Feagin, R.A., 2008. Vegetation's role in coastal protection. *Science*, 320, 176–177.
- Feagin, R.A.; Lozada-Bernard, S.M.; Ravens, T.; Moller, I.; Yeager, K.M., and Baird, A.H., 2009. Does vegetation prevent wave erosion of salt marsh edges? *Proceedings of the National Academy of Sciences USA*, 106, 10101–10113.
- Feagin, R.A.; Mukherjee, N.; Shanker, K.; Baird, A.H.; Cinner, J.; Kerr, A.M.; Koedam, N.; Sridhar, A.; Arthur, R.; Jayatissa, L.P.; Lo Seen, D.; Menon, M.; Rodriguez, S.; Shamsuddoha, M., and Dahdouh-Guebas, F., 2009. Shelter from the storm? Use and misuse of bioshields for managing natural disasters on the coast. *Conservation Letters*. In press. DOI: 10.1111/j.1755-263X.2009.00087.x
- Feagin, R.A.; Sherman, D.J., and Grant, W.E., 2005. Coastal erosion, global sea-level rise, and the loss of sand dune plant habitats. *Frontiers in Ecology and the Environment*, 3, 359–364.
- Fitzgerald, D.M. and Van Heteren, S., 1999. Classification of paragalacial barrier systems: coastal New England, USA. *Sedimentology*, 46, 1083–1108.
- Gutschick, V.P. and BassiriRad, H., 2003. Extreme events as shaping physiology, ecology, and evolution of plants: toward a unified definition and evaluation of their consequences. *New Phytologist*, 160, 21–42.
- Halpern, B.S.; Silliman, B.R.; Olden, J.D.; Bruno, J.F., and Bertness, M.D., 2007. Incorporating positive interactions in aquatic restoration and conservation. *Frontiers in Ecology and the Environment*, 5, 153–160.
- Houston, J.R., 2008. The economic value of beaches: A 2008 update. *Shore & Beach*, 76, 22–26.
- Intergovernmental Panel on Climate Change, 2007. Global climate projections. In: Solomon, S.; Qin, D.; Manning, M.; Chen, Z.; Marquis, M.; Avery, K.B.; Tignor, M., and Miller, H.L. (eds.), *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, United Kingdom: Cambridge University Press.
- Jentsch, A.; Kreyling, J., and Beierkuhnlein, C., 2007. A new generation of climate-change experiments: events, not trends. *Frontiers in Ecology and the Environment*, 5, 365–374.
- Jones, C.G.; Lawton, J.H., and Shachak, M., 1997. Positive and negative effects of organisms as physical ecosystem engineers. *Ecology*, 78, 1946–1957.
- Kirwan, M.L. and Murray, A.B., 2007. A coupled geomorphic and ecological model of tidal marsh evolution. *Proceedings of the National Academy of Sciences USA*, 104, 6118–6122.
- Koske, R.E.; Gemma, J.N.; Corkidi, L.; Siguenza, C., and Rincon, E., 2004. Arbuscular mycorrhizas in coastal sand dunes. In: Martínez, M.; Psuty, N., and Lubke, R. (eds.), *Coastal Sand Dunes: Ecology and Restoration*. New York: Springer-Verlag.
- Leatherman, S.P.; Williams, A.T., and Fisher, J.S., 1977. Over-wash sedimentation associated with a large-scale northeaster. *Marine Geology*, 24, 109–121.
- Martínez, M.L.; Intralawan, A.; Vazquez, G.; Pérez-Maqueo, O.; Sutton, P., and Landgrave, R., 2007. The coasts of our world: ecological, economic and social importance. *Ecological Economics*, 63, 254–272.
- Menon, M. and Sridhar, A., 2007. An Appraisal of Coastal Regulation Law in Tsunami-Affected Mainland India. Report to the UN Development Programme, Post-tsunami Environment Initiative, pp. 105–149.
- Michener, W.K.; Blood, E.R.; Bildstein, K.L.; Brinson, M.M., and Gardner, L.R., 1997. Climate change, hurricanes and tropical storms, and rising sea level in coastal wetlands. *Ecological Applications*, 7, 770–801.
- Mitteager, W.A.; Burke, A., and Nordstrom, K.F., 2006. Landscape features and restoration potential on private shorefront lots in New Jersey, USA. *Journal of Coastal Research*, Special Issue No. 39, pp. 891–898.
- Morris, J.T.; Sundareshwar, P.V.; Nietch, C.T.; Kjerfve, B., and Cahoon, D.R., 2002. Responses of coastal wetlands to rising sea level. *Ecology*, 83, 2869–2877.
- Morton, R.A. and Sallenger, A.H., Jr., 2003. Morphological impacts of extreme storms on sandy beaches and barriers. *Journal of Coastal Research*, 19, 560–573.
- Mukherjee, N.; Balakrishnan, M., and Shanker, K., 2009. Bioshields and ecological restoration in tsunami-affected areas in India. In: Dahl, E.; Moskness, E., and Stotturup, J. (eds.), *Integrated Coastal Zone Management*. Hoboken, New Jersey: Wiley Blackwell Publishing.
- Nordstrom, K.F., 2000. *Beaches and Dunes of Developed Coasts*. New York: Cambridge University Press.
- Nordstrom, K.F., 2008. *Beaches and Dune Restoration*. New York: Cambridge University Press.
- Orson, R.A.; Warren, R.S., and Niering, W.A., 1998. Interpreting sea level rise and rates of vertical marsh accretion in a southern New England tidal salt marsh. *Estuarine, Coastal and Shelf Science*, 47, 419–429.
- Pérez-Maqueo, O.; Intralawan, A., and Martínez, M.L., 2007. Coastal disasters from the perspective of ecological economics. *Ecological Economics*, 63, 273–284.
- Pilkey, O.H. and Fraser, M.E., 2003. *A Celebration of the World's Barrier Islands*. New York: Columbia University Press.
- Pilkey, O.H. and Young, R.S., 2005. Will Hurricane Katrina impact shoreline management? Here's why it should. *Journal of Coastal Research*, 21, iii–ix.
- Psuty, N.P., 2004. The coastal foredune: a morphological basis for regional coastal dune development. In: Martínez, M. and Psuty, N.P. (eds.), *Coastal Dunes: Ecology and Conservation*. Berlin: Springer-Verlag.
- Redfield, A.C. and Rubin, M., 1962. The age of salt marsh peat and its relation to recent changes in sea level at Barnstable, Massachusetts. *Proceedings of the National Academy of Sciences USA*, 48, 1728–1735.
- Rodriguez, S.; Balasubramanian, G.; Peter, S.M.; Duraiswamy, M., and Jaiprakash, P., 2008. Beyond the Tsunami: Community Perceptions of Resources, Policy and Development, Post Tsunami Interventions and Community Institutions in Tamil Nadu, India. Bangalore, India: Report to the UN Development Programme/ UNTRS, Chennai and ATREE, Bangalore.
- Schlacher, T.A.; Dugan, J.; Schoeman, D.S.; Lastra, M.; Jones, A.; Scapini, F.; McLachlan, A., and Defeo, O., 2007. Sandy beaches at the brink. *Diversity and Distributions*, 13, 556–560.

- Shao, G.; Shugart, H.H., and Hayden, B.P., 1996. Functional classifications of coastal barrier island vegetation. *Journal of Vegetation Science*, 7, 391–396.
- Sridhar, A.; Arthur, R.; Goenka, D.; Jairaj, B.; Mohan, T.; Rodriguez, S., and Shanker, K., 2006. Review of the Swaminathan Committee Report on the CRZ Notification. New Delhi, India: UN Development Programme, 32p.
- Tsoar, H., 2005. Sand dunes mobility and stability in relation to climate. *Physica A*, 357, 50–56.
- U.S. Commission on Ocean Policy, 2004. *Final Report: An Ocean Blueprint for the 21st Century*. Washington, DC: U.S. Commission on Ocean Policy.
- Webster, P.J.; Holland, G.J.; Curry, J.A., and Chang, H.R., 2005. Changes in tropical cyclone number, duration, and intensity in a warming environment. *Science*, 309, 1844–1846.
- Weinstein, M.P.; Baird, R.C.; Conover, D.O.; Gross, M.; Keulartz, J.; Loomis, D.K.; Naveh, Z.; Peterson, S.B.; Reed, D.J.; Roe, E.; Swanson, R.L.; Swart, J.A.A.; Teal, J.M.; Turner, R.E., and van der Windt, H.J., 2007. Managing coastal resources in the 21st century. *Frontiers in Ecology and the Environment*, 5, 43–48.
- Williams, A.M.; Feagin, R.A.; Smith, W.K., and Jackson, N.L., 2009. Ecosystem impacts of Hurricane Ike: perspectives of the Coastal Barrier Island Network (CBIN). *Shore & Beach*, 77, 71–76.
- Young, D.R.; Shao, G., and Brinson, M.M., 1995. The impact of the October 1991 northeaster storm on barrier island shrub thickets (*Myrica cerifera*). *Journal of Coastal Research*, 11, 1322–1328.

Authors Queries

Journal: **Journal of Coastal Research**

Paper: **coas-26-05-06**

Title: **Barrier Islands: Coupling Anthropogenic Stability with Ecological Sustainability**

Dear Author

During the preparation of your manuscript for publication, the questions listed below have arisen. Please attend to these matters and return this form with your proof. Many thanks for your assistance

Query Reference	Query	Remarks
1	Please supply * footnote contents	

AUTHORS: THE FOLLOWING QUERIES/COMMENTS ARE FOR EDITORIAL STAFF FROM PROOFREADERS. The editorial staff will contact you if they have any questions regarding these queries and comments.

EDITORIAL STAFF: Query markers appear in boxes in the margins as a question mark followed by a number. Please respond to each query in the margin of the corresponding proof page. Comment markers are suggested corrections that will also appear in the margin, but as a "C" followed by a number. If you wish to implement the suggested correction, please transfer the correction to the margin and text of the corresponding proof page.