

Homework #1: Solutions

#4.1

Show that if $f(p)$ has period p , the average value of f is the same over any interval of length p .

Solution:

Let a be any real number. It suffices to show that $\int_a^{a+p} f(x) dx = \int_0^p f(x) dx$. Calculating it follows that

$$\begin{aligned}\int_a^{a+p} f(x) dx &= \int_a^p f(x) dx + \int_p^{a+p} f(x) dx \\ &= \int_a^p f(x) dx + \int_0^a f(t+p) dt \\ &= \int_a^p f(x) dx + \int_0^a f(t) dt \\ &= \int_0^p f(x) dx.\end{aligned}$$

#5.5

Find the Fourier series for

$$f(x) = \begin{cases} 0, & -\pi < x < 0 \\ -1, & 0 < x < \frac{\pi}{2} \\ 1, & \frac{\pi}{2} < x < \pi \end{cases}$$

Solution:

$$\begin{aligned}a_0 &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx \\ &= \frac{1}{\pi} \int_0^{\pi/2} (-1) dx + \frac{1}{\pi} \int_{\pi/2}^{\pi} (1) dx \\ &= 0.\end{aligned}$$

$$\begin{aligned}a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} \cos(nx) f(x) dx \\ &= \frac{1}{\pi} \int_0^{\pi/2} (-1) \cos(nx) dx + \frac{1}{\pi} \int_{\pi/2}^{\pi} \cos(nx) dx \\ &= -\frac{1}{n\pi} \sin(nx) \Big|_0^{\pi/2} + \frac{1}{n\pi} \sin(nx) \Big|_{\pi/2}^{\pi} \\ &= -\frac{2}{n\pi} \sin\left(\frac{n\pi}{2}\right) \\ &= -\frac{2}{\pi} \left(1 - \frac{1}{3} + \frac{1}{5} - \dots\right)\end{aligned}$$

$$\begin{aligned}
b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} \sin(nx) f(x) dx \\
&= \frac{1}{\pi} \int_0^{\pi/2} (-1) \sin(nx) dx + \frac{1}{\pi} \int_{\pi/2}^{\pi} \sin(nx) dx \\
&= -\frac{1}{n\pi} \cos(nx) \Big|_0^{\pi/2} - \frac{1}{n\pi} \cos(nx) \Big|_{\pi/2}^{\pi} \\
&= \frac{1}{n\pi} \cos(n\pi/2) - \frac{1}{n\pi} - \frac{1}{n\pi} \cos(n\pi) + \frac{1}{n\pi} \cos(n\pi/2) \\
&= \frac{2}{n\pi} \cos(n\pi/2) - \frac{1}{n\pi} (1 + (-1)^n) \\
&= \frac{2}{\pi} \left(-\frac{1}{2} + \frac{1}{4} - \frac{1}{6} + \dots \right) - \frac{2}{\pi} \left(\frac{1}{2} + \frac{1}{4} + \frac{1}{6} + \dots \right) \\
&\equiv -\frac{4}{\pi} \left(\frac{1}{2} + \frac{1}{6} + \frac{1}{10} + \dots \right)
\end{aligned}$$

Therefore,

$$\begin{aligned}
f(x) &= -\frac{2}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{2n-1} \cos((2n-1)x) - \frac{4}{\pi} \sum_{n=1}^{\infty} \frac{1}{4n-2} \sin((4n-2)x) \\
&= \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{(-1)^n}{2n-1} \cos((2n-1)x) - \frac{2}{\pi} \sum_{n=1}^{\infty} \frac{1}{2n-1} \sin((4n-2)x).
\end{aligned}$$

#7.13

If $f(x) = \frac{1}{2}a_0 + \sum_{n=1}^{\infty} a_n \cos(nx) + \sum_{n=1}^{\infty} b_n \sin(nx) = \sum_{n=-\infty}^{\infty} c_n e^{inx}$, find a relationship between a_n , b_n , and c_n .

Solution:

$$\begin{aligned}
\sum_{n=-\infty}^{\infty} c_n e^{inx} &= \sum_{n=-\infty}^{-1} c_n e^{inx} + c_0 + \sum_{n=1}^{\infty} c_n e^{inx} \\
&= \sum_{n=1}^{\infty} c_{-n} e^{-inx} + c_0 + \sum_{n=1}^{\infty} c_n e^{inx} \\
&= c_0 + \sum_{n=1}^{\infty} [c_{-n} e^{-inx} + c_n e^{inx}] \\
&= c_0 + \sum_{n=1}^{\infty} [c_{-n} (\cos(nx) - i \sin(nx)) + c_n (\cos(nx) + i \sin(nx))] \\
&= c_0 + \sum_{n=1}^{\infty} ((c_{-n} + c_n) \cos(nx) + \sum_{n=1}^{\infty} i(c_n - c_{-n}) \sin(nx))
\end{aligned}$$

Therefore,

$$a_0 = c_0$$

$$a_n = (c_n + c_{-n})$$

$$b_n = i(c_n - c_{-n})$$

Inverting it follows that

$$c_0 = a_0$$

$$ia_n + b_n = 2ic_n$$

$$\Rightarrow c_n = \frac{a_n - ib_n}{2}$$

$$-ia_n + b_n = -2ic_{-n}$$

$$\Rightarrow c_{-n} = \frac{a_n + ib_n}{2}$$

#8.18

Find the Fourier series for $f(x) = x^2$ on the interval $0 < x < 10$.

Solution:


$$\begin{aligned}c_0 &= \frac{1}{20} \int_0^{10} x^2 dx \\&= \frac{1}{60} \cdot 10^3 \\&= \frac{5}{3} \cdot 10,000 \\&= \frac{50,000}{3}\end{aligned}$$

$$\begin{aligned}c_n &= \frac{1}{10} \int_0^{10} x^2 e^{-inx} dx \\&= \frac{1}{10} \left(\frac{x^2}{in\pi} e^{-inx} \Big|_0^{10} - \frac{5}{in\pi} \int_0^{10} 2x e^{-inx} dx \right) \\&= \frac{1}{10} \left(\frac{500}{in\pi} - \frac{50}{in\pi} \int_0^{10} x e^{-inx} dx \right) \\&= \frac{5}{in\pi} - \frac{1}{in\pi} \left(\frac{5}{in\pi} x e^{-inx} \Big|_0^{10} - \frac{5}{in\pi} \int_0^{10} e^{-inx} dx \right) \\&= \frac{5}{in\pi} + \frac{50}{n^2\pi^2}\end{aligned}$$

$$\Rightarrow f(x) = \frac{50}{3} + \frac{5}{\pi} \sum_{n=-\infty}^{\infty} \left(\frac{1}{in\pi} + \frac{1}{n^2\pi^2} \right) e^{inx/5}$$

#9.23

Find the Fourier sine series for the following function:



Solution:

Let $f(x)$ be defined by:

$$f(x) = \begin{cases} \frac{2h}{l}x, & 0 \leq x \leq l/2 \\ -\frac{2h}{l}x + 2h, & l/2 < x < l \end{cases}$$

Then,

$$\begin{aligned}
 b_n &= \frac{2}{l} \int_0^l f(x) \sin\left(\frac{n\pi x}{l}\right) dx \\
 &= \frac{2}{l} \int_0^{l/2} \frac{2h}{l}x \sin\left(\frac{n\pi x}{l}\right) dx + \frac{2}{l} \int_{l/2}^l \left(-\frac{2h}{l}x + 2h\right) \sin\left(\frac{n\pi x}{l}\right) dx \\
 &= \frac{4h}{l^2} \int_0^{l/2} x \sin\left(\frac{n\pi x}{l}\right) dx - \frac{4h}{l^2} \int_{l/2}^l x \sin\left(\frac{n\pi x}{l}\right) dx + \frac{4h}{l} \int_{l/2}^l \sin\left(\frac{n\pi x}{l}\right) dx \\
 &= \frac{4h}{l^2} \left[\frac{l}{n\pi} x \left(-\cos\left(\frac{n\pi x}{l}\right) \right) \Big|_0^{l/2} + \frac{1}{n\pi} \int_0^{l/2} \cos\left(\frac{n\pi x}{l}\right) dx \right] \\
 &\quad - \frac{4h}{l^2} \left[\frac{l}{n\pi} x \left(-\cos\left(\frac{n\pi x}{l}\right) \right) \Big|_{l/2}^l + \frac{1}{n\pi} \int_{l/2}^l \cos\left(\frac{n\pi x}{l}\right) dx \right] \\
 &\quad + \frac{4h}{n\pi} \left. \cos\left(\frac{n\pi x}{l}\right) \right|_{l/2}^l \\
 &= \frac{4h}{l^2} \left[\frac{l^2}{2n\pi} \left(-\cos\left(\frac{n\pi}{2}\right) \right) + \frac{l^2}{n^2\pi^2} \sin\left(\frac{n\pi}{2}\right) \right] \\
 &\quad - \frac{4h}{l^2} \left[\frac{l^2}{n\pi} \left(-\cos(n\pi) \right) - \frac{l^2}{2n\pi} \left(-\cos\left(\frac{n\pi}{2}\right) \right) + \frac{l^2}{n^2\pi^2} \left[\sin(n\pi) - \sin\left(\frac{n\pi}{2}\right) \right] \right] \\
 &\quad - \frac{4h}{n\pi} \left[\cos(n\pi) - \cos\left(\frac{n\pi}{2}\right) \right] \\
 &= -\frac{2h}{n\pi} \cos\left(\frac{n\pi}{2}\right) + \frac{4h}{n^2\pi^2} \sin\left(\frac{n\pi}{2}\right) + \frac{4h}{n\pi} \cos(n\pi) - \frac{2h}{n\pi} \cos\left(\frac{n\pi}{2}\right) + \frac{4h}{n^2\pi^2} \sin\left(\frac{n\pi}{2}\right) \\
 &\quad - \frac{4h}{n\pi} \cos(n\pi) + \frac{4h}{n\pi} \cos\left(\frac{n\pi}{2}\right)
 \end{aligned}$$

$$\Rightarrow b_n = \frac{8h}{n\pi^2} \sin\left(\frac{n\pi}{2}\right) \checkmark$$

Therefore,

$$f(x) = \frac{8h}{\pi^2} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)^2} \sin\left(\frac{(2n-1)\pi x}{l}\right).$$