

Name _____

Midterm Exam
March 4, 2010

This test consists of three parts. For the first part, you may write your answers directly on the exam, if you wish. For the other parts, use separate sheets of paper. Useful equations can be found at the start of part 3. The total test is worth 200 points.

Part I: Multiple Choice [40 points]

For each question, choose the best answer (2 points each)

- Observations indicate that the Sun is currently moving vertically at about 7 km/s northwards, away from the galactic disk. What is the long term effect of this motion?
 - It will move to a positive z -value, and then orbit indefinitely at that z -value
 - It will oscillate up and down, but never straying very far from the disk
 - It will follow a large orbit which takes it well out into the halo
 - It will settle back down into the plane of the galaxy over time; this is a temporary phenomenon
 - It will leave the galaxy in this direction
- Which of the following is *not* a typical characteristic of a thick disk star, compared to a thin disk star?
 - It will tend to have a larger radius, and hence is thicker
 - It will tend to be older
 - It will have more random motions; *i.e.* less perfectly circular
 - It will tend to have lower metallicity
 - Actually, all of these *are* characteristics of thick disk stars
- Most of the mass of our galaxy is probably in the form of
 - Black holes in the nucleus
 - Stars, gas, and dust in the disk
 - Globular clusters out in the halo
 - Old stars in the bulge
 - Unknown dark matter spread throughout the halo
- Although the bulge of our galaxy is obscured by dust, we are still confident it is there because
 - We can detect its gravitational contribution to the orbits of stars
 - Neutrinos from it are observed at Earth
 - We can see stars from it that have been expelled to our neighborhood
 - We can see it with infrared light, which penetrates the gas and dust
 - We simply infer its presence since other spiral galaxies like ours have bulges

5. If the surface temperature of a star is doubled, but its radius remains the same, its luminosity would increase by a factor of
A) 2 B) 4 C) 8 D) 16 E) 256
6. Spectroscopic parallax is a technique for measuring distance based on the idea that
A) For main sequence stars, the spectral class more or less tells you the luminosity (absolute magnitude) , and combined with the apparent magnitude you can get the distance
B) The spectrum of a star tells you its velocity; combined with other information, this gives you the distance
C) There is a relationship between the spectrum of a variable star and its period
D) The spectrum of a star tells you its approximate age; this can be combined with how long it took light to get here to get the actual distance
E) The apparent angle of the spectrum as viewed from different points in Earth's orbit tells you the distance to the star
7. Suppose one had a catalog of the apparent magnitude of all the brightest planetary nebulas in a nearby galaxy. What *additional* information would we need to find the distance to this galaxy? Assume there is no dust obscuring the galaxy.
A) The period of at least one Cepheid variable star
B) The parallax to at least one of the planetary nebulas in that galaxy
C) The physical size of at least one of the planetary nebulas
D) The spectral class of one or more planetary nebulas
E) None of the above; you already have sufficient information to find the distance
8. List the types of electromagnetic radiation below in order from longest wavelength to shortest
A) X-rays, infrared, visible, ultraviolet, radio
B) Radio, ultraviolet, visible, infrared, X-rays
C) X-rays, ultraviolet, visible, infrared, radio
D) Radio, infrared, visible, ultraviolet, X-rays
E) None of the above are correct
9. Which of the following is most likely to be found very close to the galactic plane?
A) Thin disk stars
B) Halo stars
C) Cool molecular clouds and star forming regions
D) Hot thin hydrogen clouds
E) Globular clusters
10. The oldest stars can be found in the
A) Thin disk B) Thick disk C) Bulge D) Globular Clusters E) Nucleus

11. In the moving cluster method, how does one determine the actual direction that a cluster of stars is moving?
- A) One calculates the transverse and radial velocities, then you project the resulting vector
 - B) The angular motion of all the stars in the cluster are extended to the vanishing point, or convergent point, where they all come together; this is the direction of motion
 - C) Differential Doppler shifts on different stars throughout the cluster are used to determine the direction of overall motion
 - D) The ages of the stars is estimated, and based on the location of the molecular cloud from which it was born, one can then determine the direction of motion
 - E) The speed of the stars in the cluster is compared with the mass and size, and the Virial theorem then tells you the direction of motion
12. The distance to the center of our galaxy is probably closest to
- A) 80 pc
 - B) 800 pc
 - C) 8 Mpc
 - D) 80 Mpc
 - E) 8 kpc
13. The abundance of which of the following elements would be included when computing the metallicity of a star?
- A) Iron, but not neon
 - B) Iron and neon, but not carbon
 - C) Iron, neon, and carbon, but not helium nor hydrogen
 - D) Iron, neon, carbon, and helium, but not hydrogen
 - E) Iron, neon, carbon, and hydrogen, but not helium
14. The hottest spectral type of those listed is
- A) A
 - B) B
 - C) K
 - D) M
 - E) F
15. The O and B main sequence stars are the first to die because
- A) They have the least mass, and therefore run out of fuel the fastest
 - B) They burn their fuel the fastest, and therefore run out of fuel the fastest
 - C) They are the coolest, and therefore they soon lose temperatures sufficient to sustain fusion
 - D) They have the most mass, and attract more mass making them unstable
 - E) The question is false; these stars live the longest, not the shortest
16. Which of the following is probably closest to the speed of the Sun around the center of the galaxy?
- A) 200 km/s
 - B) 600 km/s
 - C) 2,000 km/s
 - D) 6,000 km/s
 - E) 20,000 km/s
17. The mass of the black hole at the center of our galaxy was estimated by
- A) Observing the Schwarzschild radius
 - B) Sending a probe and measuring the gravitational pull
 - C) Studying the orbit of some stars that were very close to it
 - D) Measuring the Doppler shift of gas falling into it
 - E) Studying the bending of light from objects behind it

18. In galaxies like ours, how does the approximate rotational velocity depend on distance as you move away from the center of the galaxy?
- A) At large radii, it falls off as about $1/\sqrt{R}$
 - B) At large radii, it falls off as about $1/R$
 - C) At large radii, it rises as about \sqrt{R}
 - D) At large radii, it rises as about R
 - E) It is roughly constant at large radii
19. The disk of our galaxy is in some ways more difficult to study than that of other galaxies, primarily because
- A) Our galaxy's disk is very dim, making it difficult to see
 - B) The motion of our disk is exceptionally chaotic, making patterns hard to spot
 - C) We are in the plane of our galaxy, where the dust blocks our view of much of the disk
 - D) The stars in our galaxy are so close they overexpose in most telescopes
 - E) The stars are so close that we can't use the usual assumption that the distance to all the stars in the galaxy is identical
20. The spiral arms of a galaxy, according to current theory,
- A) Are self-propagating waves of higher density
 - B) Contains a fixed set of stars and gas that orbits together
 - C) Is actually a magnetic phenomenon, connected by magnetic field lines
 - D) Represents a series of exploding stars, which set off formation of new stars in a spiral pattern
 - E) Is actually a giant pinwheel toy made by ultrapowerful aliens for their amusement

Part II: Short essays [60 points]

Write approximately a paragraph or two explaining the following (15 points each).

21. Explain what a Cepheid Variable star is, and how it can be used to determine the distance to an object containing one. Include all appropriate equations, explaining how they are relevant.
22. Two clusters have identical metallicity, but one is much older than the other. Explain how one might observationally determine which cluster is older.
23. Most of the stars in our neighborhood are members of the thin or thick disk, but a very small number are halo stars. Explain how these stars might noticeably differ from ordinary stars in our neighborhood. Give at least two ways they generally differ.
24. What is the source of the 21 cm radio line? What type of gas does it come from? Why is it important in understanding the rotation of the galaxy?

Part III: Calculation: [100 points]

For each of the following problems, give the answer, explaining your work. The value of each portion appears in square brackets. Some possibly useful equations appear below. [20 points each]

<p align="center">Units</p> <p>1 AU = 1.496×10^{11} m 1 pc = 3.086×10^{16} m 1 eV = 1.602×10^{-19} J 1 y = 3.155×10^7 s $R_{\odot} = 6.955 \times 10^8$ m $M_{\odot} = 1.989 \times 10^{30}$ kg $L_{\odot} = 3.839 \times 10^{26}$ W $T_{\odot} = 5777$ K</p>	<p align="center">Physical Constant</p> <p>$k_B = 1.381 \times 10^{-23}$ J/K $k_B = 8.671 \times 10^{-5}$ eV/K $\sigma = 5.670 \times 10^{-8}$ W/m²/K⁴ $\hbar = 1.055 \times 10^{-34}$ J·s $\hbar = 6.582 \times 10^{-16}$ eV·s $G = 6.673 \times 10^{-11}$ m³/kg/s²</p>	<p align="center">Gravitational Potential Energy</p> $E_p = -\frac{1}{2} \int d^3\mathbf{r} \int d^3\mathbf{r}' \frac{G\rho(\mathbf{r})\rho(\mathbf{r}')}{ \mathbf{r}-\mathbf{r}' }$ $E_p = \frac{1}{2} \int d^3\mathbf{r} \rho(\mathbf{r}) \Phi(\mathbf{r})$	
<p align="center">Brightness/Magnitude</p> $b = 2.52 \times 10^{-8} \text{ W/m}^2 (10^{-\frac{2}{5}m})$		<p align="center">Black Body Radiation</p> $u = \frac{\pi^2 (k_B T)^4}{15 (\hbar c)^3}$ $\lambda_{\text{max}} T = 2.8978 \times 10^{-3} \text{ m} \cdot \text{K}$	<p align="center">Black Hole</p> $R_s = \frac{2GM}{c^2}$ <p align="center">Relative speed, Circular Orbits</p> $v_r = R_0 \sin l \left(\frac{V}{R} - \frac{V_0}{R_0} \right)$ <p align="center">Doppler Shift</p> $1+z = \frac{\lambda}{\lambda_0} = \sqrt{\frac{1+v_r/c}{1-v_r/c}}$
<p align="center">Moving Cluster</p> $d = \frac{v_r \tan \theta}{\mu}$	<p align="center">Planetary Nebula</p> $M^* = -4.47$	<p align="center">Type Ia SN</p> $M_{\text{max}} = -19.3$	<p align="center">Cepheid Period/Luminosity</p> $M = -2.81 \log(P) - 1.43$


25. A certain galaxy has an unusual distribution of matter. The density of matter takes the form

$$\rho(r) = \frac{A}{r}$$

at a distance r from the center of the galaxy.


- (a) [6] What is the total amount of mass contained within a radius R from the center of the galaxy?
- (b) [8] What is the magnitude of the gravitational acceleration g as a function of R ?
- (c) [6] What is the velocity V for circular orbits as a function of R ?

26. A distant star is a recurrent nova. By sheer luck, there is a circular ring of gas surrounding the star, as sketched at right, exactly perpendicular to our line of sight, with 1 cm representing 1 arc-second. Exactly 30 days after the nova goes off, the ring is suddenly illuminated. (1 d = 86,400 s)



- (a) [8] What is the radius of the ring, in AU?
- (b) [12] What is the distance to the nova/ring, in pc?

27. A type Ia supernova goes off in a distant galaxy! The apparent magnitude as a function of time is sketched at right.



- (a) [5] What is the approximate distance to this galaxy?
- (b) [7] With a good telescope, one can observe planetary nebulas that are as dim as about $m = 30$. Would planetary nebulas be a good way to check the distance to this galaxy? Why or why not?
- (c) [8] A study of the spectrum from this galaxy shows that the Lyman-alpha line, normally at a wavelength of 121.57 nm, has been shifted to a wavelength of 153.28 nm. What is the speed of this galaxy, and is it towards us or away from us? You may leave your answer as a fraction of c if you wish.

28. A cluster of stars has total mass M and radius R . The mass distribution is such that the gravitational potential energy is given by

$$E_p = -\frac{2GM^2}{R}$$

- (a) [8] According to the Virial theorem, what is the kinetic energy of this cluster? What is the total energy?
- (b) [8] Due to an amazing coincidence, half of the stars suddenly explode, causing them to completely expel their mass from the cluster. Hence the mass suddenly is reduced, so $M \rightarrow \frac{1}{2}M$. The speed and position of all remaining stars are unchanged. Immediately after the explosion, what is the new potential, kinetic, and total energy of the cluster?
- (c) [4] Does the cluster now have enough gravity to hold it together? Or is it too little energy? Or is it right at the boundary?

29. A star at $\beta = 30^\circ$ is tracked over the course of four years. Plotted at right is the position of the star in mas, with x denoting the motion parallel to the plane of the Earth's orbit around the Sun, and y denoting the apparent motion perpendicular to that orbit.

- (a) [7] Explain qualitatively the two types of motion seen; *i.e.*, what is causing the oscillatory motion, and what is causing the gradual drift?
- (b) [5] What is the angular velocity μ_x and μ_y for this star?
- (c) [8] What is the approximate parallax p and distance d in parsecs for this star?

