

Physics 742 – Graduate Quantum Mechanics 1
Solutions to Chapter 13

2. [15] Complete the computation of the spin-orbit splitting for hydrogen for the $2p$, $3p$, and $3d$ states of hydrogen. Write your answers as multiples of $\alpha^2 |E_n|$, where α is the fine structure constant and $|E_n|$ is the unperturbed binding energy of this state.

Starting from equation (13.8) in the class notes, we have

$$\Delta E = \frac{\hbar^2 (2l+1)}{4m^2 c^2} \int_0^\infty r^2 dr \left(\frac{1}{r} \frac{dV_c(r)}{dr} \right) R_{nl}^2(r) = \frac{\hbar^2 (2l+1) k_e e^2}{4m^2 c^2} \int_0^\infty \frac{dr}{r} R_{nl}^2(r).$$

Recalling that the energy of the n 'th state is $|E_n| = k_e e^2 / 2n^2 a_0$, this can be rewritten as

$$\Delta E = |E_n| \frac{\hbar^2 (2l+1) a_0 n^2}{2m^2 c^2} \int_0^\infty \frac{dr}{r} R_{nl}^2(r) = |E_n| \frac{\hbar^2 (2l+1) n^2}{2m^2 c^2 a_0} \int_0^\infty \frac{dr}{r} a_0^3 R_{nl}^2(r).$$

Substituting $a_0 = \hbar^2 / k_e e^2 m$, and then using $\alpha = k_e e^2 / \hbar c$, this can be rewritten as

$$\Delta E = |E_n| \frac{k_e^2 e^4 (2l+1) n^2}{2c^2 \hbar^2} \int_0^\infty \frac{dr}{r} a_0^3 R_{nl}^2(r) = \frac{1}{2} \alpha^2 |E_n| n^2 (2l+1) \int_0^\infty \frac{dr}{r} a_0^3 R_{nl}^2(r).$$

Only the final integral remains. We find

$$\begin{aligned} \int_0^\infty \frac{dr}{r} a_0^3 R_{21}^2(r) &= \frac{1}{a_0^2} \int_0^\infty \frac{dr}{r} \frac{r^2}{24} e^{-r/a_0} = \frac{1}{24} \int_0^\infty x e^{-x} dx = \frac{1}{24}, \\ \int_0^\infty \frac{dr}{r} a_0^3 R_{31}^2(r) &= \frac{1}{a_0^2} \int_0^\infty \frac{dr}{r} \frac{2^5 r^2}{3^7} \left(1 - \frac{r}{6a_0}\right)^2 e^{-2r/3a_0} = \frac{2^5}{3^7} \left(\frac{3}{2}\right)^2 \int_0^\infty x \left(1 - \frac{x}{4}\right)^2 e^{-x} dx \\ &= \frac{8}{243} \cdot \left(1 - \frac{2 \cdot 2}{4} + \frac{6}{16}\right) = \frac{1}{81}, \\ \int_0^\infty \frac{dr}{r} a_0^3 R_{32}^2(r) &= \frac{1}{a_0^4} \int_0^\infty \frac{dr}{r} \frac{2^3 r^4}{3^9 \cdot 5} e^{-2r/3a_0} = \frac{2^3}{3^9 \cdot 5} \left(\frac{3}{2}\right)^4 \int_0^\infty x^3 e^{-x} dx = \frac{6}{3^5 \cdot 10} = \frac{1}{405}. \end{aligned}$$

We now just substitute these numbers into our previous equations, to find

$$\begin{aligned} \Delta E_{2p} &= \frac{1}{2} \alpha^2 |E_2| 2^2 (3) \frac{1}{24} = \frac{1}{4} \alpha^2 |E_2|, \\ \Delta E_{3p} &= \frac{1}{2} \alpha^2 |E_3| 3^2 (3) \frac{1}{81} = \frac{1}{6} \alpha^2 |E_3|, \\ \Delta E_{3d} &= \frac{1}{2} \alpha^2 |E_3| 3^2 (5) \frac{1}{405} = \frac{1}{18} \alpha^2 |E_3|. \end{aligned}$$

3. [5] Prove, as asserted in section C, that $\int (3\hat{r}_j\hat{r}_k - \delta_{jk}) d\Omega = 0$. This is actually nine formulas, but only six of them are independent.

We simply write $\hat{\mathbf{r}} = \mathbf{r}/r = (\sin\theta\cos\phi, \sin\theta\sin\phi, \cos\theta)$ and work out all the components. We'll do it as a matrix for fun.

$$\int (3\hat{r}_j\hat{r}_k - \delta_{jk}) d\Omega = \int \begin{pmatrix} 3\sin^2\theta\cos^2\phi - 1 & 3\sin^2\theta\cos\phi\sin\phi & 3\sin\theta\cos\theta\cos\phi \\ 3\sin^2\theta\cos\phi\sin\phi & 3\sin^2\theta\sin^2\phi - 1 & 3\sin\theta\cos\theta\sin\phi \\ 3\sin\theta\cos\theta\cos\phi & 3\sin\theta\cos\theta\sin\phi & 3\cos^2\theta - 1 \end{pmatrix} d\Omega$$

We start with the integral over ϕ . It is easy to see that the integral of $\cos\phi$ or $\sin\phi$ is just zero, and since $\cos\phi\sin\phi = \frac{1}{2}\sin(2\phi)$, this can be seen to integrate to zero. The integral of 1 is trivial, and we can use Carlson's rule on the $\cos^2\phi$ and $\sin^2\phi$ (they are each half of the size of the interval), so we have

$$\begin{aligned} \int (3\hat{r}_j\hat{r}_k - \delta_{jk}) d\Omega &= 2\pi \int_0^\pi \begin{pmatrix} \frac{3}{2}\sin^2\theta - 1 & 0 & 0 \\ 0 & \frac{3}{2}\sin^2\theta - 1 & 0 \\ 0 & 0 & 3\cos^2\theta - 1 \end{pmatrix} \sin\theta d\theta \\ &= \begin{pmatrix} \pi & 0 & 0 \\ 0 & \pi & 0 \\ 0 & 0 & -2\pi \end{pmatrix} \int_{-1}^1 (1 - 3\cos^2\theta) d(\cos\theta) = 0. \end{aligned}$$