

MTH 225: Homework #8

Due Date: April 12, 2024

1. Let $A \in M_{2 \times 2}(\mathbb{C})$ be given by

$$A = \begin{bmatrix} 5/2 & i/2 \\ -i/2 & 5/2 \end{bmatrix}.$$

- Show that A is Hermitian.
- Find the eigenvalues and eigenvectors of A .
- Show that A is positive semidefinite.
- Find the square root of A .

2. Let $A \in M_{2 \times 2}(\mathbb{C})$ be given by

$$A = \begin{bmatrix} 3 & -3 \\ 1 & -1 \end{bmatrix}.$$

- Find an orthonormal basis for $\ker(A)$.
- Find an orthonormal basis for $\text{im}(A)$.
- Find the SVD of A .
- Find the polar form of A .

3. Find $\exp(A)$ when A is given by

(a) $A = \begin{bmatrix} 2 & 1 \\ 0 & -1 \end{bmatrix}$

(b) $A = \begin{bmatrix} 0 & -2 \\ 2 & 0 \end{bmatrix}$

(c) $A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$

Hint: The third matrix is not diagonalizable. You will have to use the definition of the $\exp(A)$ and find a pattern for the powers of A .

- Prove for all $A \in M_{n \times n}(\mathbb{C})$ that $A \exp(A) = \exp(A)A$.
- Prove that if O is the $n \times n$ zero matrix then $\exp(O) = I$.
- Prove for all $A \in M_{n \times n}(\mathbb{C})$ and $a, b \in \mathbb{R}$ that $\exp((a+b)A) = \exp(aA) \exp(bA)$.
- Prove for all $A \in M_{n \times n}(\mathbb{C})$ that $\exp(-A) = \exp(A)^{-1}$.
- Prove for all $A \in M_{n \times n}(\mathbb{C})$ that $\exp(A^*) = \exp(A)^*$.
- Prove that if $A \in M_{n \times n}(\mathbb{C})$ satisfies $A^* = -A$ then $\exp(A)$ is unitary.
- Prove that if $A \in M_{n \times n}(\mathbb{C})$ is diagonalizable then

$$\det(\exp(A)) = e^{\text{Tr}(A)}.$$

Hint: Recall that if $\lambda_1, \dots, \lambda_n$ are the eigenvalues of A then $\text{Tr}(A) = \lambda_1 + \dots + \lambda_n$.

11. Consider the inconsistent system $A\vec{x} = \vec{b}$ when

$$A = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & -1 \\ 0 & 1 & 1 \\ -1 & 1 & -1 \end{bmatrix} \text{ and } \vec{b} = \begin{bmatrix} 2 \\ 5 \\ 6 \\ 6 \end{bmatrix}.$$

- (a) Find the associated normal equations.
 - (b) Find the least square solution to this system.
12. The median price (in thousands of dollars) of existing homes in a certain metropolitan area from 1989 to 1999 is summarized in the following table:

year	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999,
price	86.4	89.8	92.8	96.0	99.6	103.1	106.3	109.5	113.3	120.0	129.5 .

- (a) Set up the normal equations to find the least squares line for these data. **Hint:** Use a calculator or any kind of software to simplify the matrix computations.
 - (b) Solve the normal equations to find the least squares line for these data.
 - (c) Use your least squares line to estimate the median price of a house in 2005 and 2010.
13. A 20 pound turkey that is at the room temperature of 72° is placed in the oven at 1 : 00 PM. The temperature of the turkey is observed in 20 minute intervals to be 79° , 88° , and 96° . A turkey is cooked when its temperature reaches 165° . Using least squares, estimate how much longer you have to wait until the turkey is done cooking.