

**PHY 711 Classical Mechanics and
Mathematical Methods**
10-10:50 AM MWF Olin 103

Plan for Lecture 21:

Read Chapter 7 & Appendices A-D

Generalization of the one dimensional wave equation → various mathematical problems and techniques including:

1. Laplace transforms
2. Complex variables
3. Contour integrals

10/18/2019

PHY 711 Fall 2019 – Lecture 21

1

1

Date	F&W Reading	Topic	Assignment	Due
1 Mon, 8/26/2019	Chap. 1	Introduction	#1	8/30/2019
2 Wed, 8/28/2019	Chap. 1	Scattering theory	#2	9/02/2019
3 Fri, 8/30/2019	Chap. 1	Scattering theory	#3	9/04/2019
4 Mon, 9/02/2019	Chap. 1	Scattering theory	#4	9/06/2019
5 Wed, 9/04/2019	Chap. 2	Non-inertial coordinate systems	#5	9/09/2019
6 Fri, 9/06/2019	Chap. 3	Calculus of Variation	#6	9/11/2019
7 Mon, 9/9/2019	Chap. 3	Calculus of Variation	#7	9/13/2019
8 Wed, 9/11/2019	Chap. 3	Lagrangian Mechanics		
9 Fri, 9/13/2019	Chap. 3	Lagrangian Mechanics	#8	9/16/2019
10 Mon, 9/16/2019	Chap. 3 & 6	Constants of the motion	#9	9/20/2019
11 Wed, 9/18/2019	Chap. 3 & 6	Hamiltonian equations of motion		
12 Fri, 9/20/2019	Chap. 3 & 6	Liouville theorem	#10	9/23/2019
13 Mon, 9/23/2019	Chap. 3 & 6	Canonical transformations		
14 Wed, 9/25/2019	Chap. 4	Small oscillations about equilibrium	#11	9/30/2019
15 Fri, 9/27/2019	Chap. 4	Normal modes of vibration	#12	10/02/2019
16 Mon, 9/30/2019	Chap. 7	Motion of strings	#13	10/04/2019
17 Wed, 10/02/2019	Chap. 7	Sturm-Liouville equations	#14	10/07/2019
18 Fri, 10/04/2019	Chap. 7	Sturm-Liouville equations		
19 Mon, 10/07/2019	Chap. 7	Fourier transform methods		
20 Wed, 10/09/2019	Chap. 1-4,6-7	Review		
<input type="checkbox"/> Fri, 10/11/2019	No class	Fall break		
<input type="checkbox"/> Mon, 10/14/2019	No class	Take-home exam		
<input type="checkbox"/> Wed, 10/16/2019	No class	Take-home exam		
21 Fri, 10/18/2019	Chap. 7	Contour integrals; Exam due	#15	10/23/2019

10/18/2019

PHY 711 Fall 2019 – Lecture 21

2

2

Laplace transforms
Laplace transforms can be used to solve initial value problems. The Laplace transform of a function $\phi(x)$ is defined as
$\mathcal{L}_\phi(p) \equiv \int_0^\infty e^{-px} \phi(x) dx. \quad (24)$
Assuming that $\phi(x)$ is well-behaved in the interval $0 \leq x \leq \infty$, the following properties are useful:
$\mathcal{L}_{d\phi/dx}(p) = -\phi(0) + p\mathcal{L}_\phi(p), \quad (25)$
and
$\mathcal{L}_{d^2\phi/dx^2}(p) = -\frac{d\phi(0)}{dx} - p\phi(0) + p^2\mathcal{L}_\phi(p). \quad (26)$

10/18/2019

PHY 711 Fall 2019 – Lecture 21

3

3

These identities allow us to turn a differential equation for $\phi(x)$ into an algebraic equation for $\mathcal{L}_\phi(p)$. We then need to perform an inverse Laplace transform to find $\phi(x)$. For illustration, we will consider a simple example with $\tau(x) = 1$, $\sigma(x) = 1$, $\lambda = 0$. The differential equation then becomes

$$-\frac{d^2\phi(x)}{dx^2} = F(x), \quad (27)$$

where we will take the initial conditions to be $\phi(0) = 0$ and $d\phi(0)/dx = 0$. For our example, we will also take $F(x) = F_0 e^{-\gamma x}$. Multiplying both sides of the equation by e^{-px} and integrating $0 \leq x \leq \infty$, we find

$$\mathcal{L}_\phi(p) = -\frac{F_0}{p^2(\gamma + p)}. \quad (28)$$

10/18/2019

PHY 711 Fall 2019 – Lecture 21

4

4

In general the inverse Laplace transform involves performing a contour integral, but we can use the following simple relations

$$\mathcal{L}_1 = \int_0^\infty e^{-px} dx = \frac{1}{p}. \quad (29)$$

$$\mathcal{L}_x = \int_0^\infty x e^{-px} dx = \frac{1}{p^2}. \quad (30)$$

$$\mathcal{L}_{e^{-\gamma x}} = \int_0^\infty e^{-\gamma x} e^{-px} dx = \frac{1}{p + \gamma}. \quad (31)$$

Noting that

$$-\frac{F_0}{p^2(\gamma + p)} = -\frac{F_0}{\gamma^2} \left(\frac{1}{\gamma + p} - \frac{1}{p} + \frac{\gamma}{p^2} \right), \quad (32)$$

we see that the inverse Laplace transform gives us

$$\phi(x) = \frac{F_0}{\gamma^2} \left(1 - e^{-\gamma x} - \gamma x \right). \quad (33)$$

We can check that this is a solution to the differential equation

$$-\frac{d^2\phi}{dx^2} = F_0 e^{-\gamma x} \quad \text{for } \phi(0) = 0 \quad \text{and } \frac{d\phi}{dx}(0) = 0$$

10/18/2019

PHY 711 Fall 2019 – Lecture 21

5

5

Using Laplace transforms to solve equation :

$$\left(-\frac{d^2}{dx^2} - 1 \right) \phi(x) = F_0 \sin\left(\frac{\pi x}{L}\right) \quad \text{with } \phi(0) = 0, \frac{d\phi(0)}{dx} = 0$$

$$\mathcal{L}_\phi(p) = -\left(\frac{\pi}{L}\right) \frac{F_0}{(p^2 + 1) \left(p^2 + \left(\frac{\pi}{L}\right)^2 \right)}$$

$$= -F_0 \left(\frac{\pi/L}{(\pi/L)^2 - 1} \right) \left(\frac{1}{p^2 + 1} - \frac{1}{p^2 + (\pi/L)^2} \right)$$

$$\text{Note that : } \int_0^\infty \sin(at) e^{-pt} dt = \frac{a}{a^2 + p^2}$$

$$\Rightarrow \phi(x) = \frac{F_0}{(\pi/L)^2 - 1} \left(\sin\left(\frac{\pi x}{L}\right) - \frac{\pi}{L} \sin(x) \right)$$

10/18/2019

PHY 711 Fall 2019 – Lecture 21

6

6

Inverse Laplace transform :

$$\mathcal{L}_\phi(p) = \int_0^\infty e^{-pt} \phi(t) dt$$

$$\phi(t) = \frac{1}{2\pi i} \int_{\lambda-i\infty}^{\lambda+i\infty} e^{pt} \mathcal{L}_\phi(p) dp$$

$$\text{Check: } \frac{1}{2\pi i} \int_{\lambda-i\infty}^{\lambda+i\infty} e^{pt} \mathcal{L}_\phi(p) dp = \frac{1}{2\pi i} \int_{\lambda-i\infty}^{\lambda+i\infty} e^{pt} dp \int_0^\infty e^{-pu} \varphi(u) du$$

$$\frac{1}{2\pi i} \int_0^\infty \varphi(u) du \int_{\lambda-i\infty}^{\lambda+i\infty} e^{p(t-u)} dp = \frac{1}{2\pi i} \int_0^\infty \varphi(u) du \int_{-\infty}^\infty e^{\lambda(t-u)} e^{i\pi(t-u)} i ds$$

$$= \frac{1}{2\pi i} \int_0^\infty \varphi(u) du (e^{\lambda(t-u)} 2\pi i \delta(t-u))$$

$$= \begin{cases} \varphi(t) & \text{if } t \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

10/18/2019

PHY 711 Fall 2019 – Lecture 21

7

Complex numbers

$$i \equiv \sqrt{-1} \quad i^2 = -1$$

$$\text{Define } z = x + iy$$

$$|z|^2 = zz^* = (x+iy)(x-iy) = x^2 + y^2$$

Polar representation

$$z = \rho(\cos\phi + i\sin\phi) = \rho e^{i\phi}$$

Functions of complex variables

$$f(z) = \Re(f(z)) + i\Im(f(z)) \equiv u(x, y) + iv(x, y)$$

Derivatives: Cauchy-Riemann equations

$$\frac{\partial f(z)}{\partial x} = \frac{\partial u(z)}{\partial x} + i \frac{\partial v(z)}{\partial x} \quad \frac{\partial f(z)}{\partial y} = \frac{\partial u(z)}{\partial y} + i \frac{\partial v(z)}{\partial y} = \frac{\partial v(z)}{\partial y} - i \frac{\partial u(z)}{\partial y}$$

$$\text{Argue that } \frac{df}{dz} = \frac{\partial f(z)}{\partial z} = \frac{\partial f(z)}{\partial y} \Rightarrow \frac{\partial u(z)}{\partial x} = \frac{\partial v(z)}{\partial y} \quad \text{and} \quad \frac{\partial v(z)}{\partial x} = -\frac{\partial u(z)}{\partial y}$$

10/18/2019

PHY 711 Fall 2019 – Lecture 21

8

Analytic function

$f(z)$ is analytic if it is:

- continuous
- single valued
- its first derivative satisfies Cauchy-Riemann conditions

Which of the following functions are analytic?

$$f(z) = e^z$$

$$f(z) = z^n$$

$$f(z) = \ln z$$

$$f(z) = z^\alpha$$

10/18/2019

PHY 711 Fall 2019 – Lecture 21

9

9

Some details

$$e^z = e^{x+iy} = e^x \cos(y) + ie^x \sin(y)$$

$$\frac{\partial u}{\partial x} = e^x \cos(y) = \frac{\partial v}{\partial y} \quad \frac{\partial v}{\partial x} = e^x \sin(y) = -\frac{\partial u}{\partial y}$$

$$z^2 = (x+iy)^2 = (x^2 - y^2) + 2ixy$$

$$\frac{\partial u}{\partial x} = 2x = \frac{\partial v}{\partial y} \quad \frac{\partial v}{\partial x} = 2y = -\frac{\partial u}{\partial y}$$

10/18/2019

PHY 711 Fall 2019 -- Lecture 21

10

10

PHY 711 – Contour Integration

These notes summarize some basic properties of complex functions and their integrals. An *analytic* function $f(z)$ in a certain region of the complex plane z is one which takes a single (non-infinite) value and is differentiable within that region. Cauchy's theorem states that a closed contour integral of the function within that region has the value

$$\oint_C f(z) dz = 0. \quad (1)$$

As an example, functions composed of integer powers of z –

$$f(z) = z^n, \quad \text{for } n = 0, 1, \pm 2, \pm 3, \dots \quad (2)$$

fall in this category. Notice that non-integral powers are generally not analytic and that $n = -1$ is also special. In fact, we can show that

$$\oint_C \frac{dz}{z} = 2\pi i. \quad (3)$$

10/18/2019

PHY 711 Fall 2019 -- Lecture 21

11

11

$$\oint_C \frac{dz}{z} = 2\pi i. \quad (3)$$

This result follows from the fact that we can deform the contour to a unit circle about the origin so that $z = e^{i\theta}$. Then

$$\oint_C \frac{dz}{z} = \int_0^{2\pi} \frac{e^{i\theta}}{e^{i\theta}} id\theta = 2\pi i. \quad (4)$$

10/18/2019

PHY 711 Fall 2019 -- Lecture 21

12

12

Another result of this analysis is the Residue Theorem which states that if the complex function $g(z)$ has poles at a finite number of points z_p within a region C but is otherwise analytic, the contour integral can be evaluated according to

$$\oint_C g(z) dz = 2\pi i \sum_p \text{Res}(g_p), \quad (6)$$

where the residue is given by

$$\text{Res}(g_p) \equiv \lim_{z \rightarrow z_p} \left\{ \frac{1}{(m-1)!} \frac{d^{m-1}}{dz^{m-1}} ((z - z_p)^m g(z)) \right\}, \quad (7)$$

where m denotes the order of the pole.

10/18/2019

PHY 711 Fall 2019 – Lecture 21

13

Example:

$$\int_{-\infty}^{\infty} \frac{x^2}{1+x^4} dx = \oint \frac{z^2}{1+z^4} dz$$

$$1 + z^4 = (z - e^{i\pi/4})(z - e^{3i\pi/4})(z - e^{-i\pi/4})(z - e^{-3i\pi/4})$$

$$\oint \frac{z^2}{1+z^4} dz = 2\pi i (\text{Res}(z_p = e^{i\pi/4}) + \text{Res}(z_p = e^{3i\pi/4}))$$

$$\text{Res}(z_p = e^{i\pi/4}) = \frac{e^{i\pi/4}}{4i} \quad \text{Res}(z_p = e^{3i\pi/4}) = -\frac{e^{3i\pi/4}}{4i}$$

$$\oint \frac{z^2}{1+z^4} dz = 2\pi i \left(\frac{e^{i\pi/4}}{4i} - \frac{e^{3i\pi/4}}{4i} \right) = \frac{\pi}{\sqrt{2}}$$

10/18/2019

PHY 711 Fall 2019 – Lecture 21

14

Another example:

$$I = \int_0^{\infty} \frac{\cos(ax)}{4x^4 + 5x^2 + 1} dx.$$

$$\int_0^{\infty} \frac{\cos(ax)}{4x^4 + 5x^2 + 1} dx = \frac{1}{2} \int_{-\infty}^{\infty} \frac{e^{iax}}{4x^4 + 5x^2 + 1} dx = \frac{1}{2} \oint \frac{e^{iaz}}{4z^4 + 5z^2 + 1} dz$$

$$4z^4 + 5z^2 + 1 = 4(z-i)(z-\frac{1}{2})(z+i)(z+\frac{1}{2})$$

$$I = 2\pi i (\text{Res}(z_p = i) + \text{Res}(z_p = -\frac{1}{2}))$$

15

$$\begin{aligned}
 \int_0^\infty \frac{\cos(ax)}{4x^4 + 5x^2 + 1} dx &= \frac{1}{2} \oint \frac{e^{iaz}}{4z^4 + 5z^2 + 1} dz \\
 &= 2\pi i (\text{Res}(z_p = i) + \text{Res}(z_p = \frac{i}{2})) \\
 &= \frac{\pi}{6} (-e^{-a} + 2e^{-a/2})
 \end{aligned}$$

10/18/2019

PHY 711 Fall 2019 -- Lecture 21

16

16

Cauchy integral theorem for analytic function $f(z)$:

$$f(z) = \frac{1}{2\pi i} \oint_C \frac{f(z')}{z' - z} dz'.$$

10/18/2019

PHY 711 Fall 2019 -- Lecture 21


17

17

Example

Suppose $f(|z| \rightarrow \infty) = 0$ and for $z = x$:

$$f(x) = a(x) + ib(x)$$



10/18/2019


PHY 711 Fall 2019 -- Lecture 21

18

18

Example -- continued

$$f(z) = \frac{1}{2\pi i} \oint \frac{f(z')}{z'-z} dz' \quad \text{where } f(x) = a(x) + ib(x)$$



$$a(x) + ib(x) = \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{a(x') + ib(x')}{x' - x} dx'$$

10/18/2019

PHY 711 Fall 2019 -- Lecture 21

19

19

Example -- continued

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{f(x')}{x' - x} dx' &= \int_{-\infty}^{x-\varepsilon} \frac{f(x')}{x' - x} dx' + \int_{x+\varepsilon}^{\infty} \frac{f(x')}{x' - x} dx' + \int_{x-\varepsilon}^{x+\varepsilon} \frac{f(x')}{x' - x} dx' \\ &= P \int_{-\infty}^{\infty} \frac{f(x')}{x' - x} dx' + i\pi f(x) \end{aligned}$$

10/18/2019

PHY 711 Fall 2019 -- Lecture 21

20

20

Example -- continued

$$\begin{aligned} \int_{-\infty}^{\infty} \frac{f(x')}{x' - x} dx' &= \int_{-\infty}^{x-\varepsilon} \frac{f(x')}{x' - x} dx' + \int_{x+\varepsilon}^{\infty} \frac{f(x')}{x' - x} dx' + \int_{x-\varepsilon}^{x+\varepsilon} \frac{f(x')}{x' - x} dx' \\ &= P \int_{-\infty}^{\infty} \frac{f(x')}{x' - x} dx' + i\pi f(x) \end{aligned}$$

$$\begin{aligned} a(x) + ib(x) &= \frac{P}{2\pi i} \int_{-\infty}^{\infty} \frac{a(x') + ib(x')}{x' - x} dx' + \frac{\pi i}{2\pi i} (a(x) + ib(x)) \\ \Rightarrow a(x) &= \frac{P}{\pi} \int_{-\infty}^{\infty} \frac{b(x')}{x' - x} dx' \quad b(x) = -\frac{P}{\pi} \int_{-\infty}^{\infty} \frac{a(x')}{x' - x} dx' \end{aligned}$$

Kramers-Kronig relationships


10/18/2019

PHY 711 Fall 2019 -- Lecture 21

21

Comment on evaluating principal parts integrals

$$a(x) = \frac{P}{\pi} \int_{-\infty}^{\infty} \frac{b(x')}{x' - x} dx' = \lim_{\epsilon \rightarrow 0} \left(\frac{1}{\pi} \int_{-\infty}^{x'-\epsilon} \frac{b(x')}{x' - x} dx' + \frac{1}{\pi} \int_{x'+\epsilon}^{\infty} \frac{b(x')}{x' - x} dx' \right)$$



10/18/2019


PHY 711 Fall 2019 -- Lecture 21

22

22

Example:

$$b(x') = \begin{cases} 0 & \text{for } x' < -2L, -L < x' < L, x' > 2L \\ B_0 & \text{for } L < x' < 2L \\ -B_0 & \text{for } -2L < x' < -L \end{cases}$$



10/18/2019


PHY 711 Fall 2019 -- Lecture 21

23

23

$$a(x) = \frac{P}{\pi} \int_{-\infty}^{\infty} \frac{b(x')}{x' - x} dx' = \lim_{\epsilon \rightarrow 0} \left(\frac{1}{\pi} \int_{-\infty}^{x'-\epsilon} \frac{b(x')}{x' - x} dx' + \frac{1}{\pi} \int_{x'+\epsilon}^{\infty} \frac{b(x')}{x' - x} dx' \right)$$

$$\text{For our example: } a(x) = \frac{B_0}{\pi} \ln \left(\left| \frac{4L^2 - x^2}{L^2 - x^2} \right| \right)$$



10/18/2019

PHY 711 Fall 2019 -- Lecture 21

24

24