



PHY 711 Classical Mechanics and Mathematical Methods

10-10:50 AM MWF in Olin 103

Discussion of Lecture 6 -- Chap. 3 & 6 (F &W)

Details and extensions of Lagrangian mechanics

- 1. More about constraints and Lagrange multipliers.**
- 2. Constants of the motion**
- 3. Conserved quantities**
- 4. Legendre transformations**

Your questions

From Thomas –

On slide 15 should it be p^2/mr^4 ? On slide 31 was defining the Hamiltonian using that summation a guess or how did we come up with that?

From Julia –

Are there any tips for deciding when to apply the change of variables?

Course schedule


(Preliminary schedule -- subject to frequent adjustment.)

	Date	F&W	Topic	HW
1	Mon, 8/26/2024		Introduction and overview	#1
2	Wed, 8/28/2024	Chap. 3(17)	Calculus of variation	#2
3	Fri, 8/30/2024	Chap. 3(17)	Calculus of variation	#3
4	Mon, 9/02/2024	Chap. 3	Lagrangian equations of motion	#4
5	Wed, 9/04/2024	Chap. 3 & 6	Lagrangian equations of motion	#5
6	Fri, 9/06/2024	Chap. 3 & 6	Lagrangian equations of motion	#6
7	Mon, 9/09/2024	Chap. 3 & 6	Lagrangian to Hamiltonian formalism	
8	Wed, 9/11/2024	Chap. 3 & 6	Phase space	
9	Fri, 9/13/2024	Chap. 3 & 6	Canonical Transformations	



PHY 711 -- Assignment #6

Assigned: 9/06/2024 Due: 9/09/2024



- The figure above shows a box of mass m sliding on the frictionless surface of an inclined plane (angle θ). The inclined plane itself has a mass M and is supported on a horizontal frictionless surface. Write down the Lagrangian for this system in terms of the generalized coordinates X and s and the fixed constants of the system (θ , m , M , and g) and determine the equations of motion. (Note that X and s represent the lengths of vectors whose directions are related by the angle θ .)
- Assume that initially positions $X(t=0)=0$, $s(t=0)=0$ and that their corresponding velocities are also 0. Find the trajectories $X(t)$ and $s(t)$ for $t > 0$.

From previous lecture -- motion with constraints --

Comments on generalized coordinates: $L = L(\{q_\sigma(t)\}, \{\dot{q}_\sigma(t)\}, t)$

Here we have assumed that the generalized coordinates

q_σ are independent. Now consider the possibility that the coordinates are related through constraint equations of the form:

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}_\sigma} - \frac{\partial L}{\partial q_\sigma} = 0$$

Lagrangian: $L = L(\{q_\sigma(t)\}, \{\dot{q}_\sigma(t)\}, t)$

Constraints: $f_j = f_j(\{q_\sigma(t)\}, t) = 0$

Modified Euler - Lagrange equations: $\frac{d}{dt} \frac{\partial L}{\partial \dot{q}_\sigma} - \frac{\partial L}{\partial q_\sigma} + \sum_j \lambda_j \frac{\partial f_j}{\partial q_\sigma} = 0$

Lagrange
multipliers



Some details --

Lagrangian : $L = L(\{q_\sigma(t)\}, \{\dot{q}_\sigma(t)\}, t)$

Constraints : $f_j = f_j(\{q_\sigma(t)\}, t) = 0$


Modified Euler - Lagrange equations : $\frac{d}{dt} \frac{\partial L}{\partial \dot{q}_\sigma} - \frac{\partial L}{\partial q_\sigma} + \sum_j \lambda_j \frac{\partial f_j}{\partial q_\sigma} = 0$

This amounts to modifying our optimization problem --


$\delta S = 0$ and for each i : $\delta f_i = 0$

$\Rightarrow \delta W \equiv \delta(S + \sum_i \lambda_i f_i) = 0$, introducing the new constants λ_i .

Simple example:



$$L(u(t), \dot{u}(t)) = \frac{1}{2} m \dot{u}^2 + m g u \sin \theta$$



$$L(x, y, \dot{x}, \dot{y}) = \frac{1}{2} m (\dot{x}^2 + \dot{y}^2) - m g y$$

$$f(x, y) = \sin \theta x + \cos \theta y = 0$$

Note that: $u = x \cos \theta - y \sin \theta$

Case 1:

$$L(u(t), \dot{u}(t)) = \frac{1}{2} m \dot{u}^2 + m g u \sin \theta$$

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{u}} - \frac{\partial L}{\partial u} = 0 = m \ddot{u} - m g \sin \theta = 0$$

$$\Rightarrow \ddot{u} = g \sin \theta$$

Case 2:

$$L(x, y, \dot{x}, \dot{y}) = \frac{1}{2} m (\dot{x}^2 + \dot{y}^2) - m g y$$

$$f(x, y) = \sin \theta x + \cos \theta y = 0$$

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{x}} - \frac{\partial L}{\partial x} + \lambda \frac{\partial f}{\partial x} = 0 = m \ddot{x} + \lambda \sin \theta$$

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{y}} - \frac{\partial L}{\partial y} + \lambda \frac{\partial f}{\partial y} = 0 = m \ddot{y} + m g + \lambda \cos \theta$$

$$\sin \theta \ddot{x} + \cos \theta \ddot{y} = 0$$

$$\Rightarrow \lambda = -m g \cos \theta$$

$$(\cos \theta \ddot{x} - \sin \theta \ddot{y}) = g \sin \theta$$

Which method would you use to solve the problem?

Case 1

Case 2

Force of constraint;
normal to incline

Rational for Lagrange multipliers

Recall Hamilton's principle:

$$S = \int_{t_i}^{t_f} L(\{q_\sigma(t)\}, \{\dot{q}_\sigma(t)\}, t) dt$$

$$\delta S = 0 = \int_{t_i}^{t_f} \left(\sum_{\sigma} \left(\frac{d}{dt} \frac{\partial L}{\partial \dot{q}_\sigma} - \frac{\partial L}{\partial q_\sigma} \right) \delta q_\sigma \right) dt$$

With constraints: $f_j = f_j(\{q_\sigma(t)\}, t) = 0$

Variations δq_σ are no longer independent.

$$\delta f_j = 0 = \sum_{\sigma} \frac{\partial f_j}{\partial q_\sigma} \delta q_\sigma \quad \text{at each } t$$

\Rightarrow Add 0 to Euler-Lagrange equations in the form:

$$\sum_j \lambda_j \sum_{\sigma} \frac{\partial f_j}{\partial q_\sigma} \delta q_\sigma$$

Example --


Euler-Lagrange equations with constraints:

$$\text{Lagrangian: } L = L(\{q_\sigma(t)\}, \{\dot{q}_\sigma(t)\}, t)$$

$$\text{Constraints: } f_j = f_j(\{q_\sigma(t)\}, t) = 0$$

$$\text{Modified Euler - Lagrange equations: } \frac{d}{dt} \frac{\partial L}{\partial \dot{q}_\sigma} - \frac{\partial L}{\partial q_\sigma} + \sum_j \lambda_j \frac{\partial f_j}{\partial q_\sigma} = 0$$

Example:



$$\text{Lagrangian: } L = \frac{1}{2} m (\dot{r}^2 + r^2 \dot{\theta}^2) + mgr \cos \theta$$

$$\text{Constraints: } f = r - \ell = 0$$

Example continued:

$$\text{Lagrangian: } L = \frac{1}{2} m (\dot{r}^2 + r^2 \dot{\theta}^2) + mgr \cos \theta$$

$$\text{Constraints: } f = r - \ell = 0$$


$$\frac{d}{dt} m \dot{r} - m r \dot{\theta}^2 - mg \cos \theta + \lambda = 0$$

$$\frac{d}{dt} m r^2 \dot{\theta} + mgr \sin \theta = 0$$


$$\dot{r} = 0 = \ddot{r} \quad r = \ell$$

$$\Rightarrow \ddot{\theta} = -\frac{g}{\ell} \sin \theta$$

$$\Rightarrow \lambda = m \ell \dot{\theta}^2 + mg \cos \theta$$



Another example:



Lagrangian : $L = \frac{1}{2} m_1 \dot{\ell}_1^2 + \frac{1}{2} m_2 \dot{\ell}_2^2 + m_1 g \ell_1 + m_2 g \ell_2$

Constraints : $f = \ell_1 + \ell_2 - \ell = 0$

$$\frac{d}{dt} m_1 \dot{\ell}_1 - m_1 g + \lambda = 0$$

$$\frac{d}{dt} m_2 \dot{\ell}_2 - m_2 g + \lambda = 0$$

$$\dot{\ell}_1 + \dot{\ell}_2 = 0 = \ddot{\ell}_1 + \ddot{\ell}_2$$

Figure 19.1 Atwood's machine.

$$\Rightarrow \lambda = \frac{2m_1 m_2}{m_1 + m_2} g$$

$$\ddot{\ell}_1 = -\ddot{\ell}_2 = \frac{m_1 - m_2}{m_1 + m_2} g$$

Summary of Lagrangian formalism (without constraints)

For independent generalized coordinates $q_\sigma(t)$:

$$L = L(\{q_\sigma(t)\}, \{\dot{q}_\sigma(t)\}, t)$$

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}_\sigma} - \frac{\partial L}{\partial q_\sigma} = 0$$

Note that if $\frac{\partial L}{\partial q_\sigma} = 0$, then $\frac{d}{dt} \frac{\partial L}{\partial \dot{q}_\sigma} = 0$

$$\Rightarrow \frac{\partial L}{\partial \dot{q}_\sigma} = (\text{constant})$$

Comment -- Note that in deriving these equations we have assumed that there are only conservative forces (no friction) acting on this system. The equations are easily modified to take constraints, including those from static friction, into account. What is harder to treat is dynamic friction which is typically modeled by velocity dependent dissipative forces. However, some tricks for this have been developed such as described in the textbook by Herbert Goldstein.

Examples of constants of the motion:

Example 1: one-dimensional potential:

$$L = \frac{1}{2} m (\dot{x}^2 + \dot{y}^2 + \dot{z}^2) - V(z)$$

$$\Rightarrow \frac{d}{dt} m\dot{x} = 0 \quad \Rightarrow m\dot{x} \equiv p_x \text{ (constant)}$$

$$\Rightarrow \frac{d}{dt} m\dot{y} = 0 \quad \Rightarrow m\dot{y} \equiv p_y \text{ (constant)}$$

$$\Rightarrow \frac{d}{dt} m\dot{z} = -\frac{\partial V}{\partial z}$$

Examples of constants of the motion:

Example 2: Motion in a central potential

$$L = \frac{1}{2} m \left(\dot{r}^2 + r^2 \dot{\theta}^2 \right) - V(r)$$

$$\Rightarrow \frac{d}{dt} m r^2 \dot{\theta} = 0 \quad \Rightarrow m r^2 \dot{\theta} \equiv p_{\theta} \quad (\text{constant})$$

$$\Rightarrow \frac{d}{dt} m \dot{r} = m r \dot{\theta}^2 - \frac{\partial V}{\partial r} = \frac{p_{\theta}^2}{m r^3} - \frac{\partial V}{\partial r}$$

Note that

$$\dot{\theta} = \frac{p_{\theta}}{m r^2} \quad m r \dot{\theta}^2 = m r \left(\frac{p_{\theta}}{m r^2} \right)^2$$

Recall alternative form of Euler-Lagrange equations:

Starting from:

$$L = L(\{q_\sigma(t)\}, \{\dot{q}_\sigma(t)\}, t)$$

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}_\sigma} - \frac{\partial L}{\partial q_\sigma} = 0$$

Also note that:

$$\begin{aligned} \frac{dL}{dt} &= \sum_\sigma \frac{\partial L}{\partial q_\sigma} \dot{q}_\sigma + \sum_\sigma \frac{\partial L}{\partial \dot{q}_\sigma} \ddot{q}_\sigma + \frac{\partial L}{\partial t} \\ &= \sum_\sigma \frac{d}{dt} \frac{\partial L}{\partial \dot{q}_\sigma} \dot{q}_\sigma + \sum_\sigma \frac{\partial L}{\partial \dot{q}_\sigma} \ddot{q}_\sigma + \frac{\partial L}{\partial t} \\ &= \frac{d}{dt} \left(\sum_\sigma \frac{\partial L}{\partial \dot{q}_\sigma} \dot{q}_\sigma \right) + \frac{\partial L}{\partial t} \\ &\Rightarrow \frac{d}{dt} \left(L - \sum_\sigma \frac{\partial L}{\partial \dot{q}_\sigma} \dot{q}_\sigma \right) = \frac{\partial L}{\partial t} \end{aligned}$$

Additional constant of the motion:

$$\text{If } \frac{\partial L}{\partial t} = 0;$$

$$\text{then: } \frac{d}{dt} \left(L - \sum_{\sigma} \frac{\partial L}{\partial \dot{q}_{\sigma}} \dot{q}_{\sigma} \right) = \frac{\partial L}{\partial t} = 0$$

$$\Rightarrow L - \sum_{\sigma} \frac{\partial L}{\partial \dot{q}_{\sigma}} \dot{q}_{\sigma} = -E \quad (\text{constant})$$

Example 1: one - dimensional potential :

$$L = \frac{1}{2} m (\dot{x}^2 + \dot{y}^2 + \dot{z}^2) - V(z)$$

$$\Rightarrow \frac{d}{dt} \left(\frac{1}{2} m (\dot{x}^2 + \dot{y}^2 + \dot{z}^2) - V(z) - m\dot{x}^2 - m\dot{y}^2 - m\dot{z}^2 \right) = 0$$

$$\Rightarrow - \left(\frac{1}{2} m (\dot{x}^2 + \dot{y}^2 + \dot{z}^2) + V(z) \right) = -E \quad (\text{constant})$$

For this case, we also have $m\dot{x} \equiv p_x$ and $m\dot{y} \equiv p_y$

$$\Rightarrow E = \frac{p_x^2}{2m} + \frac{p_y^2}{2m} + \frac{1}{2} m \dot{z}^2 + V(z)$$

Summary from previous slide

$$L = \frac{1}{2}m(\dot{x}^2 + \dot{y}^2 + \dot{z}^2) - V(z) \quad \rightarrow 3 \text{ variable functions}$$

$$E = \frac{p_x^2}{2m} + \frac{p_y^2}{2m} + \frac{1}{2}m\dot{z}^2 + V(z) \quad p_x, p_y, E \text{ constant}$$

\rightarrow 1 variable function

Why might this be useful?

Additional constant of the motion -- continued:

$$\text{If } \frac{\partial L}{\partial t} = 0;$$

$$\text{then : } \frac{d}{dt} \left(L - \sum_{\sigma} \frac{\partial L}{\partial \dot{q}_{\sigma}} \dot{q}_{\sigma} \right) = \frac{\partial L}{\partial t} = 0$$

$$\Rightarrow L - \sum_{\sigma} \frac{\partial L}{\partial \dot{q}_{\sigma}} \dot{q}_{\sigma} = -E \quad (\text{constant})$$

Example 2: Motion in a central potential

$$L = \frac{1}{2} m (\dot{r}^2 + r^2 \dot{\theta}^2) - V(r) \quad \rightarrow 2 \text{ variable functions}$$

$$\Rightarrow \frac{d}{dt} \left(\frac{1}{2} m (\dot{r}^2 + r^2 \dot{\theta}^2) - V(r) - m\dot{r}^2 - mr^2 \dot{\theta}^2 \right) = 0$$

$$\Rightarrow - \left(\frac{1}{2} m (\dot{r}^2 + r^2 \dot{\theta}^2) + V(r) \right) = -E \quad (\text{constant})$$

For this case, we also have $mr^2 \dot{\theta} \equiv p_{\theta}$

$$\Rightarrow E = \frac{p_{\theta}^2}{2mr^2} + \frac{1}{2} m \dot{r}^2 + V(r) \quad \rightarrow 1 \text{ variable function}$$

Other examples

$$L = \frac{1}{2}m(\dot{x}^2 + \dot{y}^2 + \dot{z}^2) + \frac{q}{2c}B_0(-\dot{x}y + \dot{y}x)$$

$$\frac{\partial L}{\partial z} = 0 \quad \Rightarrow \quad m\dot{z} = p_z \quad (\text{constant})$$

$$E = \sum_{\sigma} \frac{\partial L}{\partial \dot{q}_{\sigma}} \dot{q}_{\sigma} - L$$

$$= m(\dot{x}^2 + \dot{y}^2 + \dot{z}^2) + \frac{q}{2c}B_0(-\dot{x}y + \dot{y}x)$$

$$- \frac{1}{2}m(\dot{x}^2 + \dot{y}^2 + \dot{z}^2) - \frac{q}{2c}B_0(-\dot{x}y + \dot{y}x)$$

$$\Rightarrow E = \frac{1}{2}m(\dot{x}^2 + \dot{y}^2) + \frac{p_z^2}{2m}$$

Other examples

$$L = \frac{1}{2}m(\dot{x}^2 + \dot{y}^2 + \dot{z}^2) - \frac{q}{c}B_0\dot{x}y$$

$$\frac{\partial L}{\partial z} = 0 \quad \Rightarrow \quad m\dot{z} = p_z \quad (\text{constant})$$

$$\frac{\partial L}{\partial x} = 0 \quad \Rightarrow \quad m\dot{x} = p_x \quad (\text{constant})$$

$$E = \sum_{\sigma} \frac{\partial L}{\partial \dot{q}_{\sigma}} \dot{q}_{\sigma} - L$$

$$= m(\dot{x}^2 + \dot{y}^2 + \dot{z}^2) - \frac{q}{c}B_0\dot{x}y$$

$$- \frac{1}{2}m(\dot{x}^2 + \dot{y}^2 + \dot{z}^2) + \frac{q}{c}B_0\dot{x}y$$

$$\Rightarrow E = \frac{1}{2}m\dot{y}^2 + \frac{p_x^2}{2m} + \frac{p_z^2}{2m}$$

Lagrangian picture

For independent generalized coordinates $q_\sigma(t)$:

$$L = L(\{q_\sigma(t)\}, \{\dot{q}_\sigma(t)\}, t)$$

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}_\sigma} - \frac{\partial L}{\partial q_\sigma} = 0$$

\Rightarrow Second order differential equations for $q_\sigma(t)$

Switching variables – Legendre transformation

Mathematical transformations for continuous functions of several variables & Legendre transforms:

Simple change of variables:


$$z(x, y) \Leftrightarrow x(y, z) ???$$

$$z(x, y) \Rightarrow dz = \left(\frac{\partial z}{\partial x} \right)_y dx + \left(\frac{\partial z}{\partial y} \right)_x dy$$

$$x(y, z) \Rightarrow dx = \left(\frac{\partial x}{\partial y} \right)_z dy + \left(\frac{\partial x}{\partial z} \right)_y dz$$

But : $\left(\frac{\partial x}{\partial y} \right)_z = - \frac{\left(\frac{\partial z}{\partial y} \right)_x}{\left(\frac{\partial z}{\partial x} \right)_y}$ Assuming $dz=0$.

Note on notation for partial derivatives

$$z(x, y) \Rightarrow dz = \left(\frac{\partial z}{\partial x} \right)_y dx + \left(\frac{\partial z}{\partial y} \right)_x dy$$



hold y fixed.

hold x fixed.

Simple change of variables -- continued:

$$z(x, y) \Rightarrow dz = \left(\frac{\partial z}{\partial x} \right)_y dx + \left(\frac{\partial z}{\partial y} \right)_x dy$$

$$x(y, z) \Rightarrow dx = \left(\frac{\partial x}{\partial y} \right)_z dy + \left(\frac{\partial x}{\partial z} \right)_y dz$$


$$\Rightarrow \left(\frac{\partial x}{\partial y} \right)_z = - \frac{(\partial z / \partial y)_x}{(\partial z / \partial x)_y} \quad \Rightarrow \left(\frac{\partial x}{\partial z} \right)_y = \frac{1}{(\partial z / \partial x)_y}$$

Simple change of variables -- continued:

Example:

$$z(x, y) = e^{x^2 + y} \quad z(x, y) \Rightarrow dz = \left(\frac{\partial z}{\partial x} \right)_y dx + \left(\frac{\partial z}{\partial y} \right)_x dy$$
$$x(y, z) = (\ln z - y)^{1/2} \quad x(y, z) \Rightarrow dx = \left(\frac{\partial x}{\partial y} \right)_z dy + \left(\frac{\partial x}{\partial z} \right)_y dz$$

$$\left(\frac{\partial x}{\partial y} \right)_z \stackrel{?}{=} - \frac{(\partial z / \partial y)_x}{(\partial z / \partial x)_y}$$

$$\left(\frac{\partial x}{\partial z} \right)_y \stackrel{?}{=} \frac{1}{(\partial z / \partial x)_y}$$

$$- \frac{1}{2(\ln z - y)^{1/2}} \stackrel{\checkmark}{=} - \frac{e^{x^2 + y}}{2xe^{x^2 + y}}$$

$$\frac{1}{2z(\ln z - y)^{1/2}} \stackrel{\checkmark}{=} \frac{1}{2xe^{x^2 + y}}$$

Now that we see that these transformations are possible, we should ask the question why we might want to do this?

An example comes from thermodynamics where we have various interdependent variables such as temperature T , pressure P , volume V , etc. etc. Often a measurable property can be specified as a function of two of those, while the other variables are also dependent on those two. For example we might specify T and P while the volume will be $V(T,P)$. Or we might specify T and V while the pressure will be $P(T,V)$.

Other examples from thermo --
For thermodynamic functions:

Internal energy: $U = U(S, V)$

$$dU = TdS - PdV$$

$$dU = \left(\frac{\partial U}{\partial S} \right)_V dS + \left(\frac{\partial U}{\partial V} \right)_S dV$$

$$\Rightarrow T = \left(\frac{\partial U}{\partial S} \right)_V \quad P = - \left(\frac{\partial U}{\partial V} \right)_S$$

Enthalpy: $H = H(S, P) = U + PV$

$$dH = dU + PdV + VdP = TdS + VdP = \left(\frac{\partial H}{\partial S} \right)_P dS + \left(\frac{\partial H}{\partial P} \right)_S dP$$

$$\Rightarrow T = \left(\frac{\partial H}{\partial S} \right)_P \quad V = \left(\frac{\partial H}{\partial P} \right)_S$$



Name	Potential	Differential Form
Internal energy	$E(S, V, N)$	$dE = TdS - PdV + \mu dN$
Entropy	$S(E, V, N)$	$dS = \frac{1}{T}dE + \frac{P}{T}dV - \frac{\mu}{T}dN$
Enthalpy	$H(S, P, N) = E + PV$	$dH = TdS + VdP + \mu dN$
Helmholtz free energy	$F(T, V, N) = E - TS$	$dF = -SdT - PdV + \mu dN$
Gibbs free energy	$G(T, P, N) = F + PV$	$dG = -SdT + VdP + \mu dN$
Landau potential	$\Omega(T, V, \mu) = F - \mu N$	$d\Omega = -SdT - PdV - Nd\mu$

Mathematical transformations for continuous functions of several variables & Legendre transforms continued:

$$z(x, y) \Rightarrow dz = \left(\frac{\partial z}{\partial x} \right)_y dx + \left(\frac{\partial z}{\partial y} \right)_x dy$$

Let $u \equiv \left(\frac{\partial z}{\partial x} \right)_y$ and $v \equiv \left(\frac{\partial z}{\partial y} \right)_x$

Define new function

$$w(u, y) \Rightarrow dw = \left(\frac{\partial w}{\partial u} \right)_y du + \left(\frac{\partial w}{\partial y} \right)_u dy$$

For $w = z - ux$, $dw = dz - udx - xdu = \cancel{udx} + vdy - \cancel{udx} - xdu$

$$dw = -xdu + vdy$$

$$\Rightarrow \left(\frac{\partial w}{\partial u} \right)_y = -x \quad \left(\frac{\partial w}{\partial y} \right)_u = \left(\frac{\partial z}{\partial y} \right)_x = v$$

Lagrangian picture

For independent generalized coordinates $q_\sigma(t)$:

$$L = L(\{q_\sigma(t)\}, \{\dot{q}_\sigma(t)\}, t)$$

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}_\sigma} - \frac{\partial L}{\partial q_\sigma} = 0$$

\Rightarrow Second order differential equations for $q_\sigma(t)$

Switching variables – Legendre transformation

Define: $H = H(\{q_\sigma(t)\}, \{p_\sigma(t)\}, t)$

$$H = \sum_{\sigma} \dot{q}_\sigma p_\sigma - L \quad \text{where } p_\sigma = \frac{\partial L}{\partial \dot{q}_\sigma}$$

$$dH = \sum_{\sigma} \left(\dot{q}_\sigma dp_\sigma + p_\sigma d\dot{q}_\sigma - \frac{\partial L}{\partial q_\sigma} dq_\sigma - \frac{\partial L}{\partial \dot{q}_\sigma} d\dot{q}_\sigma \right) - \frac{\partial L}{\partial t} dt$$

Hamiltonian picture – continued

$$H = H(\{q_\sigma(t)\}, \{p_\sigma(t)\}, t)$$

$$H = \sum_{\sigma} \dot{q}_{\sigma} p_{\sigma} - L \quad \text{where} \quad p_{\sigma} = \frac{\partial L}{\partial \dot{q}_{\sigma}}$$

$$dH = \sum_{\sigma} \left(\dot{q}_{\sigma} dp_{\sigma} + p_{\sigma} d\dot{q}_{\sigma} - \frac{\partial L}{\partial q_{\sigma}} dq_{\sigma} - \frac{\partial L}{\partial \dot{q}_{\sigma}} d\dot{q}_{\sigma} \right) - \frac{\partial L}{\partial t} dt$$

$$= \sum_{\sigma} \left(\frac{\partial H}{\partial q_{\sigma}} dq_{\sigma} + \frac{\partial H}{\partial p_{\sigma}} dp_{\sigma} \right) + \frac{\partial H}{\partial t} dt$$

$$\Rightarrow \dot{q}_{\sigma} = \frac{\partial H}{\partial p_{\sigma}} \quad \frac{\partial L}{\partial q_{\sigma}} = \frac{d}{dt} \frac{\partial L}{\partial \dot{q}_{\sigma}} \equiv \dot{p}_{\sigma} = -\frac{\partial H}{\partial q_{\sigma}} \quad \frac{\partial L}{\partial t} = -\frac{\partial H}{\partial t}$$